







## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> :  <b>C07D 223/18, A61K 31/55, C07D 471/06, 243/14, 409/12, 243/12, 223/16, 267/14; 281/10, 471/04, 471/08, 401/04, 417/04, 409/04, 243/10, 487/04 // (C07D 471/06, 223:00, 221:00) (C07D 471/04, 223:00, 221:00) (C07D 471/08, 223:00, 221:00) (C07D 487/04, 249:00, 243:00)</b></p>	<b>A1</b>	<p>(11) International Publication Number: <b>WO 99/67219</b></p> <p>(43) International Publication Date: 29 December 1999 (29.12.99)</p>
<p>(21) International Application Number: <b>PCT/US99/14096</b></p> <p>(22) International Filing Date: <b>22 June 1999 (22.06.99)</b></p> <p>(30) Priority Data:              09/102,507                      22 June 1998 (22.06.98)      US              09/162,757                      30 September 1998 (30.09.98)      US</p> <p>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications              US    09/102,507 (CIP)              Filed on                                      22 June 1998 (22.06.98)              US    09/162,757 (CIP)              Filed on                                      30 September 1998 (30.09.98)</p> <p>(71) Applicants (for all designated States except US): <b>ELAN PHARMACEUTICALS, INC. [US/US]; 800 Gateway Boulevard, South San Francisco, CA 94080 (US). ELI LILLY &amp; COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).</b></p>	<p>(72) Inventors; and          (75) Inventors/Applicants (for US only): <b>AUDIA, James, E. [US/US]; 6449 Lakeside Woods Circle, Indianapolis, IN 46278 (US). PORTER, Warren, J. [US/US]; 8037 Lieber Road, Indianapolis, IN 46260 (US). THOMPSON, Richard, C. [US/US]; 763 N. County Road, 900 West, Frankfort, IN 46041 (US). WILKIE, Stephen, C. [US/US]; 8229 Quetico Drive, Indianapolis, IN 46268 (US). STACK, Douglas, R. [US/US]; 9755 Overcrest Drive, Fishers, IN 46038 (US). SHI, Qing [CN/US]; 13138 Penneagle Drive, Carmel, IN 46033 (US).</b></p> <p>(74) Agents: <b>SWISS, Gerald, F. et al.; Burns, Doane, Swecker &amp; Mathis, L.L.P., P.O. Box 1404, Alexandria, VA 22313-1404 (US).</b></p> <p>(81) Designated States: <b>AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</b></p> <p><b>Published</b>  <i>With international search report.</i>  <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: <b>COMPOUNDS FOR INHIBITING BETA-AMYLOID PEPTIDE RELEASE AND/OR ITS SYNTHESIS</b></p> <div style="text-align: center; margin: 20px 0;">   </div> <div style="display: flex; justify-content: space-around; margin-top: 20px;"> <div style="text-align: center;">  <p>(2)</p> </div> <div style="text-align: center;">  <p>(3)</p> </div> </div> <p>(57) Abstract</p> <p>Compounds of formula (I), wherein W is a cyclic group of the type (2); (3); Y is represented by the formula (II); these compounds inhibit <math>\beta</math>-amyloid peptide release and/or its synthesis, and, accordingly, have utility in treating Alzheimer's disease. Also disclosed are pharmaceutical compositions comprising a compound which inhibits <math>\beta</math>-amyloid peptide release and/or its synthesis as well as methods for treating Alzheimer's disease both prophylactically and therapeutically with such pharmaceutical compositions.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-- 1 --

## COMPOUNDS FOR INHIBITING BETA-AMYLOID PEPTIDE RELEASE AND/OR ITS SYNTHESIS

## CROSS REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application No. 60/\_\_\_\_, which was converted pursuant to 37 C.F.R. §1.53(b) from U.S. Patent Application No. 09/102,507, filed June 22, 1998 and U.S. Provisional  
5 Application No. 60/\_\_\_\_, which was converted pursuant to 37 C.F.R. §1.53(b) from U.S. Patent Application No. 09/162,757, filed September 30, 1998, the disclosures of which is incorporated herein by reference in its entirety.

## BACKGROUND OF THE INVENTION

10 Field of the Invention

This invention relates to compounds which inhibit  $\beta$ -amyloid peptide release and/or its synthesis, and, accordingly, have utility in treating Alzheimer's disease.

15 References

The following publications, patents and patent applications are cited in this application as superscript numbers:

- 1 Glenner, et al., *Biochem. Biophys. Res. Commun.* (1984) 120:885-890.
- 20 2 U.S. Patent No. 4,666,829, issued May 19, 1987, to G. G. Glenner et al., entitled "Polypeptide Marker for Alzheimer's Disease and Its Use for Diagnosis."
- 25 3 Selkoe, *Neuron*. (1991) 6:487-498.

-- 2 --

- 4 Goate, et al., *Nature* (1990) **349**:704-706.
- 5 Chartier Harlan, et al., *Nature* (1989) **353**:844-846.
- 5 6 Murrell, et al., *Science* (1991) **254**:97-99.
- 7 Mullan, et al., *Nature Genet.* (1992) **1**:345-347.
- 8 Schenk, et al., International Patent Application Publication No. WO  
10 94/10569, "*Methods and Compositions for the Detection of Soluble  $\beta$ -Amyloid Peptide*", published 11 May 1994.
- 9 Selkoe, Scientific American, "Amyloid Protein and Alzheimer's  
15 10 Disease", pp. 2-8, November, 1991.  
Yates et al., U.S. Patent No. 3,598,859.
- 11 *Tetrahedron Letters* **1993**, *34*(48), 7685.
- 12 R. F. C. Brown et al., *Tetrahedron Letters* **1971**, *8*, 667-670.
- 20 13 A. O. King et al., *J. Org. Chem.* **1993**, *58*, 3384-3386.
- 14 U.S. Provisional Application Serial No. 60/019,790, filed June 14,  
25 1996.
- 15 R. D. Clark et al., *Tetrahedron* **1993**, *49*(7), 1351-1356.
- 16 Citron, et al., *Nature* (1992) **360**:672-674.
- 30 17 P. Seubert, *Nature* (1992) **359**:325-327.
- 18 Hansen, et al., *J. Immun. Meth.*(1989) **119**:203-210.
- 19 Games et al., *Nature* (1995) **373**:523-527.
- 35 20 Johnson-Wood et al., *PNAS USA* (1997) **94**:1550-1555.

All of the above publications, patents and patent applications are herein  
incorporated by reference in their entirety to the same extent as if each individual  
40 publication, patent or patent application was specifically and individually indicated  
to be incorporated by reference in its entirety.



-- 3 --

State of the Art

Alzheimer's Disease (AD) is a degenerative brain disorder characterized clinically by progressive loss of memory, cognition, reasoning, judgment and emotional stability that gradually leads to profound mental deterioration and ultimately death. AD is a very common cause of progressive mental failure (dementia) in aged humans and is believed to represent the fourth most common medical cause of death in the United States. AD has been observed in races and ethnic groups worldwide and presents a major present and future public health problem. The disease is currently estimated to affect about two to three million individuals in the United States alone. AD is at present incurable. No treatment that effectively prevents AD or reverses its symptoms and course is currently known.

The brains of individuals with AD exhibit characteristic lesions termed senile (or amyloid) plaques, amyloid angiopathy (amyloid deposits in blood vessels) and neurofibrillary tangles. Large numbers of these lesions, particularly amyloid plaques and neurofibrillary tangles, are generally found in several areas of the human brain important for memory and cognitive function in patients with AD. Smaller numbers of these lesions in a more restrictive anatomical distribution are also found in the brains of most aged humans who do not have clinical AD. Amyloid plaques and amyloid angiopathy also characterize the brains of individuals with Trisomy 21 (Down's Syndrome) and Hereditary Cerebral Hemorrhage with Amyloidosis of the Dutch Type (HCHWA-D). At present, a definitive diagnosis of AD usually requires observing the aforementioned lesions in the brain tissue of patients who have died with the disease or, rarely, in small biopsied samples of brain tissue taken during an invasive neurosurgical procedure.

The principal chemical constituent of the amyloid plaques and vascular amyloid deposits (amyloid angiopathy) characteristic of AD and the other

-- 4 --

disorders mentioned above is an approximately 4.2 kilodalton (kD) protein of about 39-43 amino acids designated the  $\beta$ -amyloid peptide ( $\beta$ AP) or sometimes  $A\beta$ ,  $A\beta P$  or  $\beta/A4$ .  $\beta$ -Amyloid peptide was first purified and a partial amino acid sequence was provided by Glenner, et al.<sup>1</sup> The isolation procedure and the  
5 sequence data for the first 28 amino acids are described in U.S. Patent No. 4,666,829<sup>2</sup>.

Molecular biological and protein chemical analyzes have shown that the  $\beta$ -amyloid peptide is a small fragment of a much larger precursor protein termed the  
10 amyloid precursor protein (APP), that is normally produced by cells in many tissues of various animals, including humans. Knowledge of the structure of the gene encoding APP has demonstrated that  $\beta$ -amyloid peptide arises as a peptide fragment that is cleaved from APP by protease enzyme(s). The precise biochemical mechanism by which the  $\beta$ -amyloid peptide fragment is cleaved from  
15 APP and subsequently deposited as amyloid plaques in the cerebral tissue and in the walls of the cerebral and meningeal blood vessels is currently unknown.

Several lines of evidence indicate that progressive cerebral deposition of  $\beta$ -amyloid peptide plays a seminal role in the pathogenesis of AD and can precede  
20 cognitive symptoms by years or decades. See, for example, Selkoe<sup>3</sup>. The most important line of evidence is the discovery that missense DNA mutations at amino acid 717 of the 770-amino acid isoform of APP can be found in affected members but not unaffected members of several families with a genetically determined (familial) form of AD (Goate, et al.<sup>4</sup>; Chartier Harlan, et al.<sup>5</sup>; and Murrell, et  
25 al.<sup>6</sup>) and is referred to as the Swedish variant. A double mutation changing lysine<sup>595</sup>-methionine<sup>596</sup> to asparagine<sup>595</sup>-leucine<sup>596</sup> (with reference to the 695 isoform) found in a Swedish family was reported in 1992 (Mullan, et al.<sup>7</sup>). Genetic linkage analyses have demonstrated that these mutations, as well as certain other mutations in the APP gene, are the specific molecular cause of AD in the

-- 5 --

affected members of such families. In addition, a mutation at amino acid 693 of the 770-amino acid isoform of APP has been identified as the cause of the  $\beta$ -amyloid peptide deposition disease, HCHWA-D, and a change from alanine to glycine at amino acid 692 appears to cause a phenotype that resembles AD in some patients but HCHWA-D in others. The discovery of these and other mutations in APP in genetically based cases of AD prove that alteration of APP and subsequent deposition of its  $\beta$ -amyloid peptide fragment can cause AD.

Despite the progress which has been made in understanding the underlying mechanisms of AD and other  $\beta$ -amyloid peptide related diseases, there remains a need to develop methods and compositions for treatment of the disease(s). Ideally, the treatment methods would advantageously be based on drugs which are capable of inhibiting  $\beta$ -amyloid peptide release and/or its synthesis *in vivo*.


Compounds which inhibit  $\beta$ -amyloid peptide release and/or its synthesis *in vivo* are disclosed in U.S. Patent Application No. 08/996,422, filed December 22, 1997, and entitled "Cycloalkyl, Lactam, Lactone and Related Compounds, Pharmaceutical Compositions Comprising Same, and Methods for Inhibiting  $\beta$ -Amyloid Peptide Release, and/or its Synthesis by Use of Such Compounds," the disclosure of which is incorporated herein by reference in its entirety. The present invention is directed to deoxy derivatives of such compounds.

#### SUMMARY OF THE INVENTION

This invention is directed to the discovery of a class of compounds which inhibit  $\beta$ -amyloid peptide release and/or its synthesis and, therefore, are useful in the prevention of AD in patients susceptible to AD and/or in the treatment of patients with AD in order to inhibit further deterioration in their condition.

-- 6 --


Accordingly, in one of its composition aspects, the present invention provides compounds of formula I:




wherein:

*W* is a cyclic group selected from the group consisting of:


5



-- 7 --



-- 8 --



wherein:

- ring A, together with the atoms to which it is attached, forms a carbocyclic or heterocyclic ring selected from the group consisting of aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;
- 5

-- 9 --

ring *B*, together with the atoms to which it is attached, forms a carbocyclic or heterocyclic ring selected from the group consisting of aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

5 ring *C*, together with the atoms to which it is attached, forms a heteroaryl or heterocyclic ring;

Y is represented by the formula:



$\text{R}^1$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkenyl, substituted alkynyl, substituted cycloalkyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic;

15

$\text{R}^2$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclic;

each  $\text{R}^3$  is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, acyl, aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

20

each  $\text{R}^4$  is independently selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

25

$\text{R}^5$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, aryl, aryloxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted

-- 10 --

cycloalkenyl, substituted amino, heteroaryl, heterocyclic, thioalkoxy and substituted thioalkoxy;

Q is oxygen, sulfur, -S(O)-, -S(O)<sub>2</sub>-, -C(O)- or -C(S)- ;

5 Z is represented by the formula -T-CX'X''V-, wherein T is selected from the group consisting of a bond covalently linking R<sup>1</sup> to -CX'X''-, oxygen, sulfur and -NR<sup>6</sup>-, wherein R<sup>6</sup> is hydrogen, acyl, alkyl, aryl or heteroaryl;

X' is hydrogen, hydroxy or fluoro;

X'' is hydrogen, hydroxy or fluoro, or X' and X'' together form an oxo group; and

10 V is selected from the group consisting of alkylene or substituted alkylene, or R<sup>1</sup> and Z together form an aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic or substituted heterocyclic;

X is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, acyl, aryl, 15 cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted alkenyl, heteroaryl and heterocyclic; or X and one of R<sup>4</sup> and the atoms to which they are attached form a double bond;

t is an integer from 0 to 2;

f is an integer from 0 to 2;

20 n is an integer equal to 1 or 2;

and pharmaceutically acceptable salts thereof provided that neither of X' and X'' can be hydroxy or fluoro when T is other than a covalent bond linking R<sup>1</sup> to -CX'X''-.

25 This invention also provides for novel pharmaceutical compositions comprising a pharmaceutically acceptable carrier and a compound of the formula I above.



-- 11 --

Additionally, in one of its method aspects, this invention is directed to a method for inhibiting  $\beta$ -amyloid peptide release and/or its synthesis in a cell which method comprises administering to such a cell an amount of a compound or a mixture of compounds of formula I above effective in inhibiting the cellular  
5 release and/or synthesis of  $\beta$ -amyloid peptide.

Because the *in vivo* generation of  $\beta$ -amyloid peptide is associated with the pathogenesis of AD<sup>8,9</sup>, the compounds of formula I can also be employed in conjunction with a pharmaceutical composition to prophylactically and/or  
10 therapeutically prevent and/or treat AD. Accordingly, in another of its method aspects, this invention is directed to a prophylactic method for preventing the onset of AD in a patient at risk for developing AD which method comprises administering to said patient a pharmaceutical composition comprising a pharmaceutically inert carrier and an effective amount of a compound or a mixture  
15 of compounds of formula I above.

In yet another of its method aspects, this invention is directed to a therapeutic method for treating a patient with AD in order to inhibit further deterioration in the condition of that patient which method comprises  
20 administering to said patient a pharmaceutical composition comprising a pharmaceutically inert carrier and an effective amount of a compound or a mixture of compounds of formula I above.

In formula I above, rings *A* and *B* may be the same or different and are  
25 preferably independently selected from the group consisting of aryl, cycloalkyl, cycloalkenyl, heteroaryl and heterocyclic. More preferably, rings *A* and *B* are independently selected from the group consisting of aryl and cycloalkyl. Still more preferably, rings *A* and *B* are independently aryl.

-- 12 --

Particularly preferred *A* and *B* rings include, by way of example, phenyl, substituted phenyl, including fluoro-substituted phenyl, cyclohexyl and the like.

Preferred *C* rings include, by way of example, pyrrolidinyl, piperidinyl,  
5 morpholino and the like.

Preferred  $R^1$  groups include unsubstituted aryl groups such as phenyl, 1-naphthyl, 2-naphthyl, etc.; substituted aryl groups such as monosubstituted phenyls (preferably substituents at 3 or 5 positions); disubstituted phenyls  
10 (preferably substituents at 3 and 5 positions); and trisubstituted phenyls (preferably substituents at the 3,4,5 positions). Preferably, the substituted phenyl groups do not include more than 3 substituents. Examples of substituted phenyls include, for instance, 2-chlorophenyl, 2-fluorophenyl, 2-bromophenyl, 2-hydroxyphenyl, 2-nitrophenyl, 2-methylphenyl, 2-methoxyphenyl, 2-phenoxyphenyl, 2-  
15 trifluoromethylphenyl, 4-fluorophenyl, 4-chlorophenyl, 4-bromophenyl, 4-nitrophenyl, 4-methylphenyl, 4-hydroxyphenyl, 4-methoxyphenyl, 4-ethoxyphenyl, 4-butoxyphenyl, 4-*iso*-propylphenyl, 4-phenoxyphenyl, 4-trifluoromethylphenyl, 4-hydroxymethylphenyl, 3-methoxyphenyl, 3-hydroxyphenyl, 3-nitrophenyl, 3-fluorophenyl, 3-chlorophenyl, 3-bromophenyl, 3-  
20 phenoxyphenyl, 3-thiomethoxyphenyl, 3-methylphenyl, 3-trifluoromethylphenyl, 2,3-dichlorophenyl, 2,3-difluorophenyl, 2,4-dichlorophenyl, 2,5-dimethoxyphenyl, 3,4-dichlorophenyl, 3,4-difluorophenyl, 3,4-methylenedioxyphenyl, 3,4-dimethoxyphenyl, 3,5-difluorophenyl, 3,5-dichlorophenyl, 3,5-di-(trifluoromethyl)phenyl, 3,5-dimethoxyphenyl, 2,4-difluorophenyl, 2,6-difluorophenyl, 3,4,5-trifluorophenyl, 3,4,5-  
25 trimethoxyphenyl, 3,4,5-tri-(trifluoromethyl)phenyl, 2,4,6-trifluorophenyl, 2,4,6-trimethylphenyl, 2,4,6-tri-(trifluoromethyl)phenyl, 2,3,5-trifluorophenyl, 2,4,5-trifluorophenyl, 2,5-difluorophenyl, 4-fluoro-2-trifluoromethylphenyl, 2-fluoro-4-trifluoromethylphenyl, 4-benzyloxyphenyl, 2-chloro-6-fluorophenyl, 2-fluoro-6-

-- 13 --

chlorophenyl, 2,3,4,5,6-pentafluorophenyl, 2,5-dimethylphenyl, 4-phenylphenyl and 2-fluoro-3-trifluoromethylphenyl.

Other preferred R<sup>1</sup> groups include, by way of example, adamantyl, benzyl,

5 2-phenylethyl, 3-phenyl-*n*-propyl, 4-phenyl-*n*-butyl, methyl, ethyl, *n*-propyl, *iso*-propyl, *iso*-butyl, *sec*-butyl, *tert*-butyl, *n*-pentyl, *iso*-valeryl, *n*-hexyl, cyclopropyl, cyclobutyl, cyclohexyl, cyclopentyl, cyclopenten-1-yl, cyclopenten-2-yl, cyclohex-

en-1-yl, -CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>-cyclobutyl, -CH<sub>2</sub>-cyclohexyl, -CH<sub>2</sub>-cyclopentyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclobutyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclohexyl, -CH<sub>2</sub>CH<sub>2</sub>-

10 cyclopentyl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, fluoropyridyls (including 5-fluoropyridin-3-yl), chloropyridyls (including 5-chloropyridin-3-yl), thien-2-yl, thien-3-yl, benzothiazol-4-yl, 2-phenylbenzoxazol-5-yl, furan-2-yl, benzofuran-2-yl, thionaphthen-2-yl, thionaphthen-3-yl, thionaphthen-4-yl, 2-chlorothiophen-5-yl, 3-methylisoxazol-5-yl, 2-(thiophenyl)thien-5-yl, 6-methoxythionaphthen-2-yl, 3-

15 phenyl-1,2,4-thioxadiazol-5-yl, 2-phenyloxazol-4-yl, indol-3-yl, 1-phenyl-tetrazol-5-yl, allyl, 2-(cyclohexyl)ethyl, (CH<sub>3</sub>)CH=CHCH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)-, (CH<sub>3</sub>)<sub>2</sub>C=CHCH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)-,  $\phi$ C(O)CH<sub>2</sub>-, thien-2-ylmethyl, 2-(thien-2-yl)ethyl, 3-(thien-2-yl)-*n*-propyl, 2-(4-nitrophenyl)ethyl, 2-(4-methoxyphenyl)-ethyl, norboran-2-yl, (4-methoxyphenyl)-methyl, (2-methoxyphenyl)methyl, (3-

20 methoxyphenyl)methyl, (3-hydroxyphenyl)-methyl, (4-hydroxyphenyl)methyl, (4-methoxyphenyl)methyl, (4-methylphenyl)-methyl, (4-fluorophenyl)methyl, (4-fluorophenoxy)methyl, (2,4-dichlorophenoxy)-ethyl, (4-chlorophenyl)methyl, (2-chlorophenyl)methyl, (1-phenyl)ethyl, (1-(*p*-chlorophenyl)ethyl, (1-trifluoromethyl)ethyl, (4-methoxyphenyl)ethyl, CH<sub>3</sub>OC(O)CH<sub>2</sub>-,

25 benzylthiomethyl, 5-(methoxycarbonyl)-*n*-pentyl, 3-(methoxycarbonyl)-*n*-propyl, indan-2-yl, (2-methylbenzofuran-3-yl), methoxymethyl, CH<sub>3</sub>CH=CH-, CH<sub>3</sub>CH<sub>2</sub>CH=CH-, (4-chlorophenyl)C(O)CH<sub>2</sub>-, (4-fluorophenyl)C(O)CH<sub>2</sub>-, (4-methoxyphenyl)C(O)CH<sub>2</sub>-, 4-(fluorophenyl)-NHC(O)CH<sub>2</sub>-, 1-phenyl-*n*-butyl, ( $\phi$ )<sub>2</sub>CHNHC(O)CH<sub>2</sub>CH<sub>2</sub>-, (CH<sub>3</sub>)<sub>2</sub>NC(O)CH<sub>2</sub>-, ( $\phi$ )<sub>2</sub>CHNHC(O)CH<sub>2</sub>CH<sub>2</sub>-,

-- 14 --

methylcarbonylmethyl, (2,4-dimethylphenyl)-C(O)CH<sub>2</sub>-, phenylC(O)CH<sub>2</sub>-,  
 CH<sub>3</sub>C(O)N(φ)-, ethenyl, methylthiomethyl, (CH<sub>3</sub>)<sub>3</sub>CNHC(O)CH<sub>2</sub>-,  
 diphenylmethyl, phenoxymethyl, 3,4-methylenedioxy-phenyl-CH<sub>2</sub>-,  
 benzo[b]thiophen-3-yl, (CH<sub>3</sub>)<sub>3</sub>COC(O)NHCH<sub>2</sub>-, *trans*-styryl, H<sub>2</sub>NC(O)CH<sub>2</sub>CH<sub>2</sub>-,  
 5 2-trifluoromethylphenyl-C(O)CH<sub>2</sub>, φC(O)NHCH(φ)CH<sub>2</sub>-, mesityl,  
 CH<sub>3</sub>C(=NHOH)CH<sub>2</sub>-, 4-CH<sub>3</sub>-φ-NHC(O)CH<sub>2</sub>CH<sub>2</sub>-, φC(O)CH(φ)CH<sub>2</sub>-,  
 (CH<sub>3</sub>)<sub>2</sub>CHC(O)NHCH(φ)-, CH<sub>3</sub>CH<sub>2</sub>OCH<sub>2</sub>-, CH<sub>3</sub>OC(O)CH(CH<sub>3</sub>)(CH<sub>2</sub>)<sub>3</sub>-, 2,2,2-  
 trifluoroethyl, 1-(trifluoromethyl)ethyl, 2-CH<sub>3</sub>-benzofuran-3-yl, 2-(2,4-  
 dichlorophenoxy)ethyl, φSO<sub>2</sub>CH<sub>2</sub>-, 3-cyclohexyl-*n*-propyl, CF<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>- and  
 10 N-pyrrolidinyl.

R<sup>2</sup> is preferably selected from the group consisting of alkyl, substituted  
 alkyl, alkenyl, cycloalkyl, aryl, heteroaryl and heterocyclic.

15 Particularly preferred R<sup>2</sup> substituents include, by way of example, methyl,  
 ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *sec*-butyl, *tert*-butyl,  
 -CH<sub>2</sub>CH(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, 2-methyl-*n*-butyl, 6-fluoro-*n*-hexyl, phenyl, benzyl,  
 cyclohexyl, cyclopentyl, cycloheptyl, allyl, *iso*-buten-2-yl, 3-methylpentyl, -CH<sub>2</sub>-  
 cyclopropyl, -CH<sub>2</sub>-cyclohexyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclohexyl,  
 20 -CH<sub>2</sub>-indol-3-yl, *p*-(phenyl)phenyl, *o*-fluorophenyl, *m*-fluorophenyl, *p*-  
 fluorophenyl, *m*-methoxyphenyl, *p*-methoxyphenyl, phenethyl, benzyl, *m*-  
 hydroxybenzyl, *p*-hydroxybenzyl, *p*-nitrobenzyl, *m*-trifluoromethylphenyl, *p*-  
 (CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O-benzyl, *p*-(CH<sub>3</sub>)<sub>3</sub>COC(O)CH<sub>2</sub>O-benzyl, *p*-  
 (HOOCCH<sub>2</sub>O)-benzyl, 2-aminopyridin-6-yl, *p*-(N-morpholino-CH<sub>2</sub>CH<sub>2</sub>O)-benzyl,  
 25 -CH<sub>2</sub>CH<sub>2</sub>C(O)NH<sub>2</sub>, -CH<sub>2</sub>-imidazol-4-yl, -CH<sub>2</sub>-(3-tetrahydrofuranyl), -CH<sub>2</sub>-  
 thiophen-2-yl, -CH<sub>2</sub>(1-methyl)cyclopropyl, -CH<sub>2</sub>-thiophen-3-yl, thiophen-3-yl,  
 thiophen-2-yl, -CH<sub>2</sub>-C(O)O-*t*-butyl, -CH<sub>2</sub>-C(CH<sub>3</sub>)<sub>3</sub>, -CH<sub>2</sub>CH(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>,  
 -2-methylcyclopentyl, -cyclohexen-2-yl, -CH[CH(CH<sub>3</sub>)<sub>2</sub>]COOCH<sub>3</sub>,  
 -CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>C(CH<sub>3</sub>)=CH<sub>2</sub>, -CH<sub>2</sub>CH=CHCH<sub>3</sub> (*cis* and *trans*),

-- 15 --

-CH<sub>2</sub>OH, -CH(OH)CH<sub>3</sub>, -CH(O-*i*-butyl)CH<sub>3</sub>, -CH<sub>2</sub>OCH<sub>3</sub>, -(CH<sub>2</sub>)<sub>4</sub>NH-Boc,  
-(CH<sub>2</sub>)<sub>4</sub>NH<sub>2</sub>, -CH<sub>2</sub>-pyridyl (e.g., 2-pyridyl, 3-pyridyl and 4-pyridyl), pyridyl (2-  
pyridyl, 3-pyridyl and 4-pyridyl), -CH<sub>2</sub>-naphthyl (e.g., 1-naphthyl and 2-  
naphthyl), -CH<sub>2</sub>-(N-morpholino), *p*-(N-morpholino-CH<sub>2</sub>CH<sub>2</sub>O)-benzyl,  
5 benzo[b]thiophen-2-yl, 5-chlorobenzo[b]thiophen-2-yl, 4,5,6,7-  
tetrahydrobenzo[b]thiophen-2-yl, benzo[b]thiophen-3-yl, 5-  
chlorobenzo[b]thiophen-3-yl, benzo[b]thiophen-5-yl, 6-methoxynaphth-2-yl,  
-CH<sub>2</sub>CH<sub>2</sub>SCH<sub>3</sub>, thien-2-yl, thien-3-yl, and the like.

10 Preferably, R<sup>3</sup> is selected from the group consisting of hydrogen, alkyl,  
substituted alkyl and cycloalkyl.

Particularly preferred R<sup>3</sup> substituents include, by way of example,  
hydrogen, methyl, 2-methylpropyl, hexyl, methoxycarbonylmethyl, 3,3-dimethyl-  
15 2-oxobutyl, 4-phenylbutyl, cyclopropylmethyl, 2,2,2-trifluoroethyl, cyclohexyl,  
and the like.

When present, R<sup>4</sup> is preferably alkyl or substituted alkyl.


20 R<sup>5</sup> is preferably alkyl; substituted alkyl; aryl; substituted aryl, such as  
2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2-chlorophenyl and the like;  
cycloalkyl, such as cyclohexyl and the like; or heteroaryl or heterocyclic, such as  
piperidin-1-yl, 2-pyridyl, 2-thiazolyl, 2-thienyl and the like.

25 Preferably, *f* is 0 or 1. More preferably, *f* is 0.

Preferably, *n* is 1.

-- 16 --

In one preferred embodiment of this invention, *W* is a cyclic group of the formula:



wherein

each  $R^6$  is independently selected from the group consisting of acyl,  
 5 acylamino, acyloxy, alkenyl, substituted alkenyl, alkoxy, substituted alkoxy,  
 alkyl, substituted alkyl, alkynyl, substituted alkynyl, amino, substituted amino,  
 aminoacyl, aryl, aryloxy, carboxyl, carboxyalkyl, cyano, cycloalkyl, substituted  
 cycloalkyl, halo, heteroaryl, heterocyclic, nitro, thioalkoxy, substituted  
 thioalkoxy, thioaryloxy, thioheteroaryloxy, -SO-alkyl, -SO-substituted alkyl, -SO-  
 10 aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl, and -SO<sub>2</sub>-  
 heteroaryl;

each  $R^7$  is independently selected from the group consisting of acyl,  
 acylamino, acyloxy, alkenyl, substituted alkenyl, alkoxy, substituted alkoxy,  
 alkyl, substituted alkyl, alkynyl, substituted alkynyl, amino, substituted amino,  
 15 aminoacyl, aryl, aryloxy, carboxyl, carboxyalkyl, cyano, cycloalkyl, substituted  
 cycloalkyl, halo, heteroaryl, heterocyclic, nitro, thioalkoxy, substituted  
 thioalkoxy, thioaryloxy, thioheteroaryloxy, -SO-alkyl, -SO-substituted alkyl, -SO-  
 aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl, and -SO<sub>2</sub>-  
 heteroaryl;

20  $R^8$  is selected from the group consisting of hydrogen, alkyl, substituted  
 alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, acyl, aryl,  
 cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl,  
 heteroaryl and heterocyclic;

-- 17 --

$p$  is an integer from 0 to 4; and


$q$  is an integer from 0 to 4.

Preferably,  $R^6$  and  $R^7$  are independently selected from the group consisting  
 5 of alkoxy, substituted alkoxy, alkyl, substituted alkyl, amino, substituted amino,  
 carboxyl, carboxyalkyl, cyano, halo, nitro, thioalkoxy and substituted thioalkoxy.  
 More preferably, when present,  $R^6$  and  $R^7$  are fluoro.

$R^8$  is preferably selected from the group consisting of hydrogen, alkyl,  
 10 substituted alkyl, acyl, aryl, cycloalkyl and substituted cycloalkyl. More  
 preferably,  $R^8$  is selected from the group consisting of hydrogen, alkyl,  
 substituted alkyl and cycloalkyl.

Particularly preferred  $R^8$  substituents include, by way of example,  
 15 hydrogen, methyl, 2-methylpropyl, hexyl, methoxycarbonylmethyl, 3,3-dimethyl-  
 2-oxobutyl, 4-phenylbutyl, cyclopropylmethyl, 2,2,2-trifluoroethyl, cyclohexyl,  
 and the like.


In another preferred embodiment of this invention,  $W$  is a cyclic group of  
 20 the formula:



wherein  $R^6$ ,  $R^7$ , and  $p$  are as defined herein and  $r$  is an integer from 0 to 3.


-- 18 --

In still another preferred embodiment of this invention, *W* is a cyclic group of the formula:




wherein  $R^6$  and  $p$  are as defined herein.

- 5 In yet another preferred embodiment of this invention, *W* is a cyclic ring of the formula:



wherein  $R^6$  and  $p$  are as defined herein.

- 10 In still another preferred embodiment of this invention, *W* is a cyclic ring of the formula:



wherein  $R^6$ ,  $R^8$  and  $p$  are as defined herein; and

each  $R^9$  is independently selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl,




-- 19 --

cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl,  
heteroaryl and heterocyclic; and  
 $g$  is an integer from 0 to 2.

5 When present,  $R^9$  is preferably alkyl or substituted alkyl.


In another preferred embodiment of this invention,  $W$  is a cyclic ring of the  
formula:



wherein  $R^6$ ,  $R^8$ ,  $R^9$ ,  $g$  and  $p$  are as defined herein.

10

In yet another preferred embodiment of this invention,  $W$  is a cyclic ring of  
the formula:




wherein  $R^6$ ,  $R^8$ ,  $R^9$ ,  $g$  and  $p$  are as defined herein.

15


In still another preferred embodiment of this invention,  $W$  is a cyclic ring  
of the formula:

-- 20 --




wherein  $R^6$ , each  $R^8$  and  $p$  are as defined herein.

In another preferred embodiment of this invention,  $W$  is a cyclic ring of the formula:



5 wherein  $R^6$ , each  $R^8$ ,  $R^9$ ,  $g$  and  $p$  are as defined herein.

In another preferred embodiment of this invention,  $W$  is a cyclic ring of the formula:




wherein  $R^6$ ,  $R^8$  and  $p$  are as defined herein; and

10  $R^{10}$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, cycloalkyl,

-- 21 --

substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic.


In another preferred embodiment of this invention, *W* is a cyclic ring of the  
5 formula:



wherein  $R^6$ ,  $R^{10}$  and  $p$  are as defined herein; and

D-E is selected from the group consisting of alkylene, alkenylene, substituted alkylene, substituted alkenylene and  $-N=CH-$ .

10 In another preferred embodiment of this invention, *W* is a cyclic ring of the formula:




wherein  $R^6$ ,  $R^8$ ,  $R^9$ ,  $g$  and  $p$  are as defined herein; and

Q is oxygen, sulfur,  $-S(O)-$  or  $-S(O)_2-$ .


15 In another preferred embodiment of this invention, *W* is a cyclic ring of the formula:

-- 22 --




wherein  $R^6$ ,  $R^8$  and  $p$  are as defined herein.

In another preferred embodiment of this invention,  $W$  is a cyclic ring of the formula:



5 wherein  $R^8$  is as defined herein.

In another preferred embodiment of this invention,  $W$  is a cyclic ring of the formula:




wherein  $R^8$  is as defined herein.

10


In another preferred embodiment of this invention,  $W$  is a cyclic ring of the formula:

-- 23 --




wherein  $R^8$  is as defined herein.

In another preferred embodiment of this invention,  $W$  is a cyclic ring of the formula:



5 wherein  $R^4$  and  $R^8$  are as defined herein.


In another preferred embodiment of this invention,  $W$  is a cyclic ring of the formula:



10 wherein  $R^4$  is as defined herein.


-- 24 --

In another preferred embodiment of this invention, *W* is a cyclic ring of the formula:



wherein  $R^4$ ,  $R^6$ ,  $R^8$  and  $p$  are as defined herein.

5 In another preferred embodiment of this invention, *W* is a cyclic ring of the formula:



wherein  $R^4$ ,  $R^6$ ,  $R^8$  and  $p$  are as defined herein.

Compounds of this invention include, by way of example, the following:

- 10 5-(S)-(N'-(2-(3,5-difluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(R)-hydroxy-2-(3,5-difluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 15 5(S)-(N'-(2(S)-hydroxy-2-(3,5-difluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 20 5(S)-(N'-(2(R/S)-hydroxy-3-isopropoxypropyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(R/S)-hydroxybutyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

-- 25 --

- 5(S)-(N'-(2(R/S)-3-dihydroxypropyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5 5(S)-(N'-(2(R/S)-hydroxy-3-morpholinopropyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(R/S)-hydroxytetradecyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 10 5(S)-(N'-(2(R/S)-hydroxyoctyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(R)-hydroxy-2-(3,4,5-trifluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 15 5(S)-(N'-(2(S)-hydroxy-2-(3,4,5-trifluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(R)-hydroxy-2-(4-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 20 5(S)-(N'-(2(S)-hydroxy-2-(4-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(R)-hydroxy-2-(3,5-bis-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 25 5(S)-(N'-(2(S)-hydroxy-2-(3,5-bis-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 30 5(S)-(N'-(2(R)-hydroxy-2-(3,3,3-trifluoropropyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(S)-hydroxy-2-(3,3,3-trifluoropropyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 35 5(S)-(N'-(3-methyl-2-butanone)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 40 5(S)-(N'-(3-methyl-2-(S)-hydroxybutyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 3-(N'-(3-methyl-2-(S)-2-hydroxybutyl)-L-Alaninyl)-amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

-- 26 --


- 3-(N'-(2-(R/S)-3,5-difluorophenyl-2-hydroxyethyl)-L-alaninyl)-amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1.5-benzodiazepine
- 5 5-[N'-(S)-2-(4-methylpentyl)amino-3-methylbutyryl-L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5-(S)-{N'-[(1RS, 2SR)-1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl]alaninyl}-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 10 5-(S)-{N'-[(1RS, 2RS)-1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl]alaninyl}-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 15 5-(S)-[N'-(2- $\alpha$ -tetralone)-L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one hydrochloride
- 5-(S)-[N'-(1,2,3,4-tetrahydro-2-naphthyl)-L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 20 5-(S)-{N'-[(1RS, 2SR)-1-hydroxy-2-cyclohexyl]alaninyl}-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5-(S)-[N'-(4-methylpentyl)-L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 25 Other examples are set forth in the Table below:



-- 27 --


TABLE

The compounds listed below conform to formula I as shown below:



I

W is the following structure, unless otherwise indicated:



5









10

15




20

R <sup>1</sup>	Z	Y	n	W
3,5-di-fluoro-Φ-	-CH <sub>2</sub> -CH <sub>2</sub> -	-HC(CH <sub>3</sub> )-C(O)-NH-	1	as above
3,5-di-fluoro-Φ-	CH(OH)-CH <sub>2</sub> -	-HC(CH <sub>3</sub> )-C(O)-NH-	1	as above
(CH <sub>3</sub> ) <sub>2</sub> CH-	-O-CH <sub>2</sub> -CH(OH)-CH <sub>2</sub> -	-HC(CH <sub>3</sub> )-C(O)-NH-	1	as above
CH <sub>3</sub> -	-CH <sub>2</sub> -CH(OH)-CH <sub>2</sub> -	-HC(CH <sub>3</sub> )-C(O)-NH-	1	as above
HO-CH <sub>2</sub> -	-CH(OH)-CH <sub>2</sub> -	-HC(CH <sub>3</sub> )-C(O)-NH-	1	as above
morpholine-	-CH <sub>2</sub> -CH(OH)-CH <sub>2</sub> -	-HC(CH <sub>3</sub> )-C(O)-NH-	1	as above
CH <sub>3</sub> -(CH <sub>2</sub> ) <sub>5</sub> -	-(CH <sub>2</sub> ) <sub>6</sub> -CH(OH)-CH <sub>2</sub> -	-HC(CH <sub>3</sub> )-C(O)-NH-	1	as above
CH <sub>3</sub> -	-(CH <sub>2</sub> ) <sub>5</sub> -CH(OH)-CH <sub>2</sub> -	-HC(CH <sub>3</sub> )-C(O)-NH-	1	as above
3,4,5-tri-fluoro-Φ-	-CH(OH)-CH <sub>2</sub> -	-HC(CH <sub>3</sub> )-C(O)-NH-	1	as above
p-F <sub>3</sub> C-Φ-	-CH(OH)-CH <sub>2</sub> -	-HC(CH <sub>3</sub> )-C(O)-NH-	1	as above
3,5-di-F <sub>3</sub> C-Φ-	-CH(OH)-CH <sub>2</sub> -	-HC(CH <sub>3</sub> )-C(O)-NH-	1	as above
F <sub>3</sub> C-	-CH(OH)-CH <sub>2</sub> -	-HC(CH <sub>3</sub> )-C(O)-NH-	1	as above
(CH <sub>3</sub> ) <sub>2</sub> -CH-	-C(O)-CH <sub>2</sub> -	-HC(CH <sub>3</sub> )-C(O)-NH-	1	as above
(CH <sub>3</sub> ) <sub>2</sub> -CH-	-CH(OH)-CH <sub>2</sub> -	-HC(CH <sub>3</sub> )-C(O)-NH-	1	as above

-- 28 --

$(\text{CH}_3)_2\text{-CH-}$	$\text{-CH(OH)-CH}_2\text{-}$	$\text{-HC(CH}_3\text{)-C(O)-NH-}$	1	
3,5-di-fluoro- $\Phi$ -	$\text{-CH(OH)-CH}_2\text{-}$	$\text{-HC(CH}_3\text{)-C(O)-NH-}$	1	
$\text{R}'\text{Z=}$ 		$\text{-HC(CH}_3\text{)-C(O)-NH-}$	1	
$\text{R}'\text{Z=}$ 		$\text{-HC(CH}_3\text{)-C(O)-NH-}$	1	
$\text{R}'\text{Z=}$ 		$\text{-HC(CH}_3\text{)-C(O)-NH-}$	1	

-- 29 --

		$\text{-HC(CH}_3\text{)-C(O)-NH-}$	1	
$(\text{CH}_3)_2\text{-CH-}$	$\text{-CH}_2\text{-CH}_2\text{-CH}_2\text{-}$	$\text{-HC(CH}_3\text{)-C(O)-NH-}$	1	

As is appreciated by the skilled person, compounds of the present invention exist as isomers. Herein, the Cahn-Prelog-Ingold designations of (R)- and (S)- and, for amino acid derived portions of the compounds, the L- and D- designations of stereochemistry relative to the isomers of glyceraldehyde are used to refer to specific isomers where designated. The specific isomers can be prepared by stereospecific synthesis or can be resolved and recovered by techniques known in the art, such as, chromatography on chiral stationary phases, and fractional recrystallization of addition salts formed by reagents used for that purpose. Useful methods of resolving and recovering specific stereoisomers are known in the art and described, for example, in Stereochemistry of Organic Compounds, E.L. Eliel and S.H. Wilen (Wiley-Interscience 1994), Enantiomers, Racemates and Resolutions, J. Jacques, A. Collet and S.J. Wilen (Wiley-Interscience 1981), and European Patent Application No. EP-A-838448, published April 29, 1998. It is to be understood that the invention extends to all of the isomeric forms of the compounds of the present invention, including the diastereomeric, enantiomeric and racemic forms of the compounds.

-- 30 --

Also included within the scope of this invention are prodrugs of the compounds of formula I above including acylated forms of alcohols and thiols, aminals of one or more amines, and the like.

## 5 DETAILED DESCRIPTION OF THE INVENTION

As above, this invention relates to compounds which inhibit  $\beta$ -amyloid peptide release and/or its synthesis, and, accordingly, have utility in treating Alzheimer's disease. However, prior to describing this invention in further detail, the following terms will first be defined.

10

### Definitions

The term " $\beta$ -amyloid peptide" refers to a 39-43 amino acid peptide having a molecular weight of about 4.2 kD, which peptide is substantially homologous to the form of the protein described by Glenner, et al.<sup>1</sup> including mutations and post-translational modifications of the normal  $\beta$ -amyloid peptide. In whatever form, the  $\beta$ -amyloid peptide is an approximate 39-43 amino acid fragment of a large membrane-spanning glycoprotein, referred to as the  $\beta$ -amyloid precursor protein (APP). Its 43-amino acid sequence is:

20

1  
Asp Ala Glu Phe Arg His Asp Ser Gly Tyr

11  
Glu Val His His Gln Lys Leu Val Phe Phe

25

21  
Ala Glu Asp Val Gly Ser Asn Lys Gly Ala

30

31  
Ile Ile Gly Leu Met Val Gly Gly Val Val

41  
Ile Ala Thr (SEQ ID NO: 1)

35 or a sequence which is substantially homologous thereto.

-- 31 --

"Alkyl" refers to monovalent alkyl groups preferably having from 1 to 20 carbon atoms and more preferably 1 to 6 carbon atoms. This term is exemplified by groups such as methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *n*-hexyl, and the like.

5

"Substituted alkyl" refers to an alkyl group, preferably of from 1 to 10 carbon atoms, having from 1 to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyacylamino, cyano, halogen, hydroxyl, carboxyl, carboxylalkyl, keto, thioketo, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl and -SO<sub>2</sub>-heteroaryl.

10  
15

"Alkylene" refers to divalent alkylene groups preferably having from 1 to 10 carbon atoms and more preferably 1 to 6 carbon atoms. This term is exemplified by groups such as methylene (-CH<sub>2</sub>-), ethylene (-CH<sub>2</sub>CH<sub>2</sub>-), the propylene isomers (e.g., -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>- and -CH(CH<sub>3</sub>)CH<sub>2</sub>-) and the like.

20

"Substituted alkylene" refers to an alkylene group, preferably of from 1 to 10 carbon atoms, having from 1 to 3 substituents selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkoxy, substituted cycloalkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, cyano, halogen, hydroxyl, carboxyl, carboxylalkyl, keto, thioketo, thiol, thioalkoxy, substituted thioalkoxy, aryl, heteroaryl, heterocyclic, heterocycloxy, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl, and -SO<sub>2</sub>-

25

-- 32 --

heteroaryl. Additionally, such substituted alkylene groups include those where 2 substituents on the alkylene group are fused to form one or more cycloalkyl, aryl, heterocyclic or heteroaryl groups fused to the alkylene group. Preferably, such fused cycloalkyl groups contain from 1 to 3 fused ring structures.

5

"Alkenylene" refers to divalent alkenylene groups preferably having from 2 to 10 carbon atoms and more preferably 2 to 6 carbon atoms. This term is exemplified by groups such as ethenylene ( $-\text{CH}=\text{CH}-$ ), the propenylene isomers (e.g.,  $-\text{CH}_2\text{CH}=\text{CH}-$  and  $-\text{C}(\text{CH}_3)=\text{CH}-$ ), and the like.

10

"Substituted alkenylene" refers to an alkenylene group, preferably of from 2 to 10 carbon atoms, having from 1 to 3 substituents selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkoxy, substituted cycloalkoxyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, cyano, halogen, hydroxyl, carboxyl, carboxylalkyl, keto, thioketo, thiol, thioalkoxy, substituted thioalkoxy, aryl, heteroaryl, heterocyclic, heterocycloxy, nitro -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl, and -SO<sub>2</sub>-heteroaryl. Additionally, such substituted alkylene groups include those where 2 substituents on the alkylene group are fused to form one or more cycloalkyl, aryl, heterocyclic or heteroaryl groups fused to the alkylene group.

15

20

"Alkaryl" refers to -alkylene-aryl groups preferably having from 1 to 8 carbon atoms in the alkylene moiety and from 6 to 10 carbon atoms in the aryl moiety. Such alkaryl groups are exemplified by benzyl, phenethyl and the like.

25

"Alkoxy" refers to the group "alkyl-O-". Preferred alkoxy groups include, by way of example, methoxy, ethoxy, *n*-propoxy, *iso*-propoxy,

-- 33 --

*n*-butoxy, *tert*-butoxy, *sec*-butoxy, *n*-pentoxy, *n*-hexoxy, 1,2-dimethylbutoxy, and the like.

"Substituted alkoxy" refers to the group "substituted alkyl-O-" where  
5 substituted alkyl is as defined above.

"Alkylalkoxy" refers to the group "-alkylene-O-alkyl" which includes by way of example, methylenemethoxy ( $-\text{CH}_2\text{OCH}_3$ ), ethylenemethoxy ( $-\text{CH}_2\text{CH}_2\text{OCH}_3$ ), *n*-propylene-*iso*-propoxy ( $-\text{CH}_2\text{CH}_2\text{CH}_2\text{OCH}(\text{CH}_3)_2$ ),  
10 methylene-*t*-butoxy ( $-\text{CH}_2-\text{O}-\text{C}(\text{CH}_3)_3$ ) and the like.

"Alkylthioalkoxy" refers to the group "-alkylene-S-alkyl" which includes by way of example, methylenethiomethoxy ( $-\text{CH}_2\text{SCH}_3$ ), ethylenethiomethoxy ( $-\text{CH}_2\text{CH}_2\text{SCH}_3$ ), *n*-propylene-thio-*iso*-propoxy ( $-\text{CH}_2\text{CH}_2\text{CH}_2\text{SCH}(\text{CH}_3)_2$ ),  
15 methylenethio-*t*-butoxy ( $-\text{CH}_2\text{SC}(\text{CH}_3)_3$ ) and the like.

"Alkenyl" refers to alkenyl groups preferably having from 2 to 10 carbon atoms and more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-2 sites of alkenyl unsaturation. Preferred alkenyl groups  
20 include ethenyl ( $-\text{CH}=\text{CH}_2$ ), *n*-propenyl ( $-\text{CH}_2\text{CH}=\text{CH}_2$ ), *iso*-propenyl ( $-\text{C}(\text{CH}_3)=\text{CH}_2$ ), and the like.

"Substituted alkenyl" refers to an alkenyl group as defined above having from 1 to 3 substituents selected from the group consisting of alkoxy, substituted  
25 alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkoxy, substituted cycloalkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, cyano, halogen, hydroxyl, carboxyl, carboxylalkyl, keto, thioketo, thiol, thioalkoxy, substituted thioalkoxy, aryl, heteroaryl, heterocyclic, heterocycloxy,

-- 34 --

nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl, and -SO<sub>2</sub>-heteroaryl.

"Alkynyl" refers to alkynyl groups preferably having from 2 to 10 carbon atoms and more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-2 sites of alkynyl unsaturation. Preferred alkynyl groups include ethynyl (-C≡CH), propargyl (-CH<sub>2</sub>C≡CH) and the like.

"Substituted alkynyl" refers to an alkynyl group as defined above having from 1 to 3 substituents selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkoxy, substituted cycloalkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, cyano, halogen, hydroxyl, carboxyl, carboxylalkyl, keto, thioketo, thiol, thioalkoxy, substituted thioalkoxy, aryl, heteroaryl, heterocyclic, heterocycloxy, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl, and -SO<sub>2</sub>-heteroaryl.

"Acyl" refers to the groups alkyl-C(O)-, substituted alkyl-C(O)-, cycloalkyl-C(O)-, substituted cycloalkyl-C(O)-, aryl-C(O)-, heteroaryl-C(O)- and heterocyclic-C(O)- where alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, heteroaryl and heterocyclic are as defined herein.

"Acylamino" refers to the group -C(O)NRR where each R is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, heteroaryl, heterocyclic and where both R groups are joined to form a heterocyclic group, wherein alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

"Amino" refers to the group -NH<sub>2</sub>.



-- 35 --

“Substituted amino” refers to the group  $-N(R)_2$  where each R is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, heterocyclic and where both R groups are joined to form a heterocyclic group. When both R groups are hydrogen,  $-N(R)_2$  is an amino group. Examples of substituted amino groups include, by way of illustration, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-heteroarylamino, mono- and di-heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic, and the like.

The term “amino-blocking group” or “amino-protecting group” refers to any group which, when bound to an amino group, prevents undesired reactions from occurring at the amino group and which may be removed by conventional chemical and/or enzymatic procedures to reestablish the amino group. Any known amino-blocking group may be used in this invention. Typically, the amino-blocking group is selected so as to render the resulting blocked-amino group unreactive to the particular reagents and reaction conditions employed in a subsequent pre-determined chemical reaction or series of reactions. After completion of the reaction(s), the amino-blocking group is selectively removed to regenerate the amino group. Examples of suitable amino-blocking groups include, by way of illustration, *tert*-butoxycarbonyl (Boc), benzyloxycarbonyl (Cbz), acetyl, 1-(1'-adamantyl)-1-methylethoxycarbonyl (Acm), allyloxycarbonyl (Aloc); benzyloxymethyl (Bom), 2-*p*-biphenylisopropylloxycarbonyl (Bpoc), *tert*-butyldimethylsilyl (Bsi), benzoyl (Bz), benzyl (Bn), 9-fluorenyl-methyloxycarbonyl (Fmoc), 4-methylbenzyl, 4-methoxybenzyl, 2-nitrophenylsulfenyl (Nps), 3-nitro-2-pyridinesulfenyl (NPys), trifluoroacetyl (Tfa),

-- 36 --

2,4,6-trimethoxybenzyl (Tmob), trityl (Trt), and the like. If desired, amino-blocking groups covalently attached to a solid support may also be employed.

"Aminoacyl" refers to the group -NRC(O)R where each R is independently  
5 hydrogen, alkyl, substituted alkyl, aryl, heteroaryl, or heterocyclic wherein alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

"Aminoacyloxy" refers to the group -NRC(O)OR where each R is  
independently hydrogen, alkyl, substituted alkyl, aryl, heteroaryl, or heterocyclic  
10 wherein alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

"Acyloxy" refers to the groups alkyl-C(O)O-, substituted alkyl-C(O)O-,  
cycloalkyl-C(O)O-, substituted cycloalkyl-C(O)O-, aryl-C(O)O-, heteroaryl-  
15 C(O)O-, and heterocyclic-C(O)O- wherein alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, heteroaryl, and heterocyclic are as defined herein.

"Aryl" refers to an unsaturated aromatic carbocyclic group of from 6 to 14  
carbon atoms having a single ring (e.g., phenyl) or multiple condensed (fused)  
20 rings (e.g., naphthyl or anthryl). Preferred aryls include phenyl, naphthyl and the like.

Unless otherwise constrained by the definition for the aryl substituent, such  
aryl groups can optionally be substituted with from 1 to 5 substituents selected  
25 from the group consisting of acyloxy, hydroxy, acyl, alkyl, alkoxy, alkenyl, alkynyl, substituted alkyl, substituted alkoxy, substituted alkenyl, substituted alkynyl, amino, substituted amino, aminoacyl, acylamino, alkaryl, aryl, aryloxy, azido, carboxyl, carboxylalkyl, cyano, halo, nitro, heteroaryl, heterocyclic,

-- 37 --

aminoacyloxy, oxyacylamino, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioheteroaryloxy, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-heteroaryl and trihalomethyl. Preferred substituents include alkyl, alkoxy, halo, cyano, nitro, trihalomethyl, and thioalkoxy.

"Aryloxy" refers to the group aryl-O- wherein the aryl group is as defined above including optionally substituted aryl groups as also defined above.

10 "Carboxyalkyl" refers to the groups "-C(O)O-alkyl" and "-C(O)O-substituted alkyl" where alkyl is as defined above.

"Cycloalkyl" refers to cyclic alkyl groups of from 3 to 12 carbon atoms having a single cyclic ring or multiple condensed rings, including bridged, fused and spiro rings and combinations thereof. The cycloalkyl ring may optionally be fused to an aryl, heteroaryl, or heterocycle ring. Such cycloalkyl groups include, by way of example, single ring structures such as cyclopropyl, cyclobutyl, cyclopentyl, cyclooctyl, and the like, or multiple ring structures such as adamantanyl, and the like.

20

"Substituted cycloalkyl" refers to cycloalkyl groups having from 1 to 5 (preferably 1 to 3) substituents selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyacylamino, cyano, halogen, hydroxyl, carboxyl, carboxylalkyl, keto, thioketo, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl, and -SO<sub>2</sub>-heteroaryl.

25

-- 38 --

"Cycloalkenyl" refers to cyclic alkenyl groups of from 4 to 8 carbon atoms having a single cyclic ring and at least one point of internal unsaturation. Examples of suitable cycloalkenyl groups include, for instance, cyclobut-2-enyl, cyclopent-3-enyl, cyclooct-3-enyl and the like.

5

"Substituted cycloalkenyl" refers to cycloalkenyl groups having from 1 to 5 substituents selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyacylamino, cyano, halogen, hydroxyl, carboxyl, carboxylalkyl, keto, thioketo, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl, and -SO<sub>2</sub>-heteroaryl.

15

"Halo" or "halogen" refers to fluoro, chloro, bromo and iodo and preferably is either fluoro or chloro.

"Heteroaryl" refers to an aromatic group of from 1 to 15 carbon atoms and 1 to 4 heteroatoms selected from oxygen, nitrogen and sulfur within at least one ring (if there is more than one ring).

Unless otherwise constrained by the definition for the heteroaryl substituent, such heteroaryl groups can be optionally substituted with 1 to 5 substituents selected from the group consisting of acyloxy, hydroxy, acyl, alkyl, alkoxy, alkenyl, alkynyl, substituted alkyl, substituted alkoxy, substituted alkenyl, substituted alkynyl, amino, substituted amino, aminoacyl, acylamino, alkaryl, aryl, aryloxy, azido, carboxyl, carboxylalkyl, cyano, halo, nitro, heteroaryl, heterocyclic, aminoacyloxy, oxyacylamino, thioalkoxy, substituted

25

-- 39 --

thioalkoxy, thioaryloxy, thioheteroaryloxy, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-heteroaryl and trihalomethyl. Such heteroaryl groups can have a single ring (e.g., pyridyl or furyl) or multiple condensed rings (e.g., indolizinyl or benzothienyl). Preferred heteroaryls include pyridyl, pyrrolyl and furyl.

"Heteroaryloxy" refers to the group "-O-heteroaryl".

"Heterocycle" or "heterocyclic" refers to a monovalent saturated or unsaturated group having a single ring or multiple condensed rings having from 1 to 15 carbon atoms and from 1 to 4 hetero atoms selected from nitrogen, sulfur or oxygen within at least one ring.

Unless otherwise constrained by the definition for the heterocyclic substituent, such heterocyclic groups can be optionally substituted with 1 to 5 substituents selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyacylamino, cyano, halogen, hydroxyl, carboxyl, carboxylalkyl, keto, thioketo, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-aryl, and -SO<sub>2</sub>-heteroaryl. Such heterocyclic groups can have a single ring or multiple condensed rings. Preferred heterocyclics include morpholino, piperidinyl, and the like.

"Monocyclic heterocyclics" refer to single ring heterocycle groups which are exemplified by, for example, pyrrolidinyl, morpholino, and the like.

-- 40 --

"Bicyclic heterocyclics" refer to heterocyclic groups comprised of two ring systems which may be fused, spiro or bridged wherein at least one of the rings contains a heteroatom and the other ring is selected from the group consisting of cycloalkyl, cycloalkenyl, aryl, heteroaryl and heterocyclic.

- 5 Examples of fused bicyclic heterocyclic ring systems include, for instance, 3-(1,2,3,4-tetrahydro-isoquinoliny) and the like.

"Tricyclic heterocyclics" refer to heterocyclic groups comprised of three ring systems wherein each of the ring systems is independently fused, spiro or  
10 bridged wherein at least one of the rings contains a heteroatom and the remaining two rings are selected from the group consisting of cycloalkyl, cycloalkenyl, aryl, heteroaryl and heterocyclic. When the remaining two rings are cycloalkyl, cycloalkenyl or heterocyclic, these rings may optionally be spiro linked.

- 15 Examples of heterocycles and heteroaryls include, but are not limited to, pyrrole, furan, imidazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, indazole, purine, quinolizine, isoquinoline, quinoline, phthalazine, naphthylpyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, phenanthroline,  
20 isothiazole, phenazine, isoxazole, phenoxazine, phenothiazine, imidazolidine, imidazoline, piperidine, piperazine, indoline, morpholino, piperidiny, tetrahydrofuranyl, and the like as well as N-alkoxy-nitrogen containing heterocycles.

- 25 "Heterocyclooxy" refers to the group "-O-heterocycle".

"Keto" or "oxo" refers to the group "=O".

-- 41 --

"Oxyacylamino" refers to the group  $-OC(O)NRR$  where each R is independently hydrogen, alkyl, substituted alkyl, aryl, heteroaryl, or heterocyclic wherein alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

5

"Thiol" refers to the group  $-SH$ .

"Thioalkoxy" refers to the group  $-S$ -alkyl.

10

"Substituted thioalkoxy" refers to the group  $-S$ -substituted alkyl.

"Thioaryloxy" refers to the group aryl-S- wherein the aryl group is as defined above including optionally substituted aryl groups also defined above.


15

"Thioheteroaryloxy" refers to the group heteroaryl-S- wherein the heteroaryl group is as defined above including optionally substituted aryl groups as also defined above.

"Thioketo" refers to the group  $=S$ .

20


The term "5,7-dihydro-6H-dibenz[b,d]azepin-6-one" refers to a polycyclic  $\epsilon$ -caprolactam ring system having the formula:



-- 42 --


wherein, for nomenclature purposes, the atoms and bonds are numbered and lettered, respectively, as shown.

The term "5,6-dihydro-4H-quinolizin-8(9H)-one" refers  
 5 to a polycyclic  $\epsilon$ -caprolactam ring system having the formula:



wherein, for nomenclature purposes, the atoms and bonds are numbered and lettered, respectively, as shown.

The term "1,3,4,7,12,12a-hexahydropyrido[2,1-b][3]benzazepin-6(2H)-  
 10 one" refers to a polycyclic  $\epsilon$ -caprolactam ring system having the formula:




wherein, for nomenclature purposes, the atoms and bonds are numbered and lettered, respectively, as shown.

The term "4,5,6,7-tetrahydro-3,7-methano-3H-benzazonin-2(1H)-one" refers to a polycyclic  $\epsilon$ -caprolactam ring system having the formula:  
 15



-- 43 --



wherein, for nomenclature purposes, the atoms and bonds are numbered and lettered, respectively, as shown.

As to any of the above groups which contain 1 or more substituents, it is understood, of course, that such groups do not contain any substitution or substitution patterns which are sterically impractical and/or synthetically non-feasible.

"Pharmaceutically acceptable salt" refers to pharmaceutically acceptable salts of a compound of formula I which salts are derived from a variety of organic and inorganic counter ions well known in the art and include, by way of example only, sodium, potassium, calcium, magnesium, ammonium, tetraalkylammonium, and the like; and when the molecule contains a basic functionality, salts of organic or inorganic acids, such as hydrochloride, hydrobromide, tartrate, mesylate, acetate, maleate, oxalate and the like can be used as the pharmaceutically acceptable salt.

The term "protecting group" or "blocking group" refers to any group which when bound to one or more hydroxyl, thiol, carboxyl groups or other protectable functional group of the compounds prevents reactions from occurring at these groups and which protecting group can be removed by conventional chemical or enzymatic steps to reestablish the unprotected functional group. The

-- 44 --

particular removable blocking group employed is not critical and preferred removable hydroxyl blocking groups include conventional substituents such as allyl, benzyl, acetyl, chloroacetyl, thiobenzyl, benzyldine, phenacyl, *t*-butyl-diphenylsilyl and any other group that can be introduced chemically onto a hydroxyl functionality and later selectively removed either by chemical or enzymatic methods in mild conditions compatible with the nature of the product.


Preferred carboxyl protecting groups include esters such as methyl, ethyl, propyl, *t*-butyl, etc., which can be removed by mild hydrolysis conditions compatible with the nature of the product.

#### Compound Preparation

When *n* is one, compounds of formula I are readily prepared by conventional acylation, followed by reductive amination as illustrated in Scheme 1.

-- 45 --

Scheme 1



-- 46 --

As shown in Scheme 1, a protected amino acid **1** (where B is a protecting group) can be coupled with an amine compound, such as **2** (where R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, p and q are as defined herein), by conventional acylation reactions to provide, after deprotection, intermediate **3**. It should be recognized by one skilled in the art that amine **2** is merely representative and those skilled in the art will recognize that amino derivatives of any of the other ring systems described herein may be employed in this reaction. Typically, this reaction is conducted using conventional coupling reagents and procedures and at least a stoichiometric amount of amino acid **1** and amine **2**. For example, well known coupling reagents such as carbodiimides with or without the use of well known additives such as N-hydroxysuccinimide, 1-hydroxybenzotriazole, etc. can be used to facilitate coupling. The reaction is conventionally conducted in an inert aprotic polar diluent such as dimethylformamide, dichloromethane, chloroform, acetonitrile, tetrahydrofuran and the like.

Acid **1** can also be coupled to amine **2** prepared by use of polymer supported forms of carbodiimide peptide coupling reagents. A polymer supported form of EDC, for example, has been described (*Tetrahedron Letters*, 34(48), 7685 (1993))<sup>11</sup>. Additionally, a new carbodiimide coupling reagent, PEPC, and its corresponding polymer supported forms have been discovered and are very useful for the preparation of such compounds.

Polymers suitable for use in making a polymer supported coupling reagent are either commercially available or may be prepared by methods well known to the artisan skilled in the polymer arts. A suitable polymer must possess pendant sidechains bearing moieties reactive with the terminal amine of the carbodiimide. Such reactive moieties include chloro, bromo, iodo and methanesulfonyl. Preferably, the reactive moiety is a chloromethyl group. Additionally, the polymer's backbone must be inert to both the carbodiimide and

-- 47 --

reaction conditions under which the ultimate polymer bound coupling reagents will be used.


Certain hydroxymethylated resins may be converted into chloromethylated  
5 resins useful for the preparation of polymer supported coupling reagents.  
Examples of these hydroxylated resins include the 4-hydroxymethylphenyl-  
acetamidomethyl resin (Pam Resin) and 4-benzyloxybenzyl alcohol resin (Wang  
Resin) available from Advanced Chemtech of Louisville, Kentucky, USA (see  
Advanced Chemtech 1993-1994 catalog, page 115). The hydroxymethyl groups  
10 of these resins may be converted into the desired chloromethyl groups by any of  
a number of methods well known to the skilled artisan.

Preferred resins are the chloromethylated styrene/divinylbenzene resins  
because of their ready commercial availability. As the name suggests, these  
15 resins are already chloromethylated and require no chemical modification prior to  
use. These resins are commercially known as Merrifield's resins and are  
available from Aldrich Chemical Company of Milwaukee, Wisconsin, USA (see  
Aldrich 1994-1995 catalog, page 899). Methods for the preparation of PEPC  
and its polymer supported forms are outlined in Scheme 1A below.

20

25

-- 48 --

Scheme 1A

Such methods are described more fully in U.S. Patent Application Serial No. 60/019,790<sup>14</sup> filed June 14, 1996 which application is incorporated herein by reference in its entirety. Briefly, PEPC is prepared by first reacting ethyl isocyanate with 1-(3-aminopropyl)pyrrolidine. The resulting urea is treated with 4-toluenesulfonyl chloride to provide PEPC. The polymer supported form is prepared by reaction of PEPC with an appropriate resin under standard conditions to give the desired reagent.


The carboxylic acid coupling reactions employing these reagents are performed at about ambient to about 45°C, for from about 3 to 120 hours. Typically, the product may be isolated by washing the reaction with CHCl<sub>3</sub> and concentrating the remaining organics under reduced pressure. As discussed *supra*, isolation of products from reactions where a polymer bound reagent has been used is greatly simplified, requiring only filtration of the reaction mixture and then concentration of the filtrate under reduced pressure.

-- 49 --

Alternatively, the acid halide of compound 1 can be employed in reaction (1) and, when so employed, it is typically employed in the presence of a suitable base to scavenge the acid generated during the reaction. Suitable bases include, by way of example, triethylamine, diisopropylethylamine, N-methylmorpholine and the like. Removal of the amine protecting group using conventional procedures and reagents then affords intermediate 3.

Intermediate 3 is then coupled with an excess of aldehyde 4, preferably with 1.1 to 2 equivalents of 4 and an excess, preferably 1.1 to 1.5 equivalents, of a reducing agent, such as sodium cyanoborohydride, to provide for compound of Formula I. Generally, this reaction is conducted in an essentially inert diluent, such as methanol, at a temperature ranging from about 0°C to about 50°C, preferably at ambient temperature, for about 0.5 to 3 hours.


Alternatively, a compound of Formula I can be prepared by reacting an acid of formula 5:



where  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{B}^1$ , T,  $\text{X}'$  and  $\text{X}''$  are as defined herein, with amine 2 under acylating reaction conditions described above, followed by deprotection of the amino protecting group ( $\text{B}^1$ ) using conventional reagents and procedures known in the art.

-- 50 --

Compounds of formula I where n is 2, can be prepared as illustrated in Scheme 2 below, by first coupling an amine, such as **2**, with a protected amino acid **9**, and then, after deprotection, converting the resulting intermediate **10** as described in Scheme 1 above.






-- 51 --

Alternatively, when  $n$  is 2, the aldehyde, or carboxylic acid units may also be coupled together prior to reaction with an amine using the reductive amination or acylation procedures described above, as appropriate. The resulting intermediate is then coupled to the amine, such as 2, to afford precursors of the compounds of formula I.

The compounds of formula I may also be prepared by any one of the following methods by conventional means as known to one of ordinary skill in the art. Some such methods are described below


(a) reductive alkylation, as shown in Scheme 3 below:



SCHEME 3

wherein  $\text{R}^1$ , T,  $\text{X}'$ ,  $\text{X}''$ , Y, W and Z are as defined herein and is described in detail in Scheme 1 above:

(b) epoxide opening, as shown in Scheme 4 below:




SCHEME 4

wherein  $\text{R}^1$ , Y and W are as defined herein;

(c) or nucleophilic displacement, as shown in Scheme 5 below:

-- 52 --



SCHEME 5

wherein  $R^1$ , Y, W and Z are as defined herein, and X is a leaving group such as halogen, tosylate, mesylate, triflate, and the like, as known to one of ordinary skill in the art.

5           Epoxide ring opening reactions provide for  $\beta$ -hydroxyl groups ( $\beta$  to the amine). The conditions employed for such reactions are well known in the art. Similarly, nucleophilic displacement reactions provide for facile methods for reacting, e.g., a primary amine with, for example, an  $\alpha$ -haloacetic acid derivative, to provide for an amino acid derivative which reaction is described in  
10           detail in International Patent Application Publication No. WO98/22441.

In any of the above reaction schemes 3-5, substituents may be protected as necessary by protecting groups as known to one of ordinary skill in the art in order to prevent reaction with the reactive group.

15

#### Synthesis of Aldehyde and Carboxylic Acid Starting materials

The aldehyde and carboxylic acids employed in the above reactions can be readily prepared by several divergent synthetic routes with the particular route selected relative to the ease of compound preparation, commercial availability of  
20           starting materials, whether  $n$  is one or two, etc.

#### A. Synthesis of Aldehydes

The aldehyde of formula 4, employed in this invention can be readily prepared by oxidizing the corresponding alcohol using conventional oxidizing

-- 53 --

agents. For example, Swern oxidation of primary alcohols affords the corresponding aldehyde. Typically, this reaction is conducted by contacting the alcohol with a mixture of oxalyl chloride and dimethyl sulfoxide in the presence of a tertiary amine, such as triethylamine. Generally, this reaction is conducted in an inert diluent, such as dichloromethane, at an initial temperature of about -78°C and then at ambient temperature for about 0.25 to 2 hours to afford the aldehyde. The alcohols employed in this reaction are either commercially available or can be prepared using conventional reagents and procedures. For example, suitable alcohols can be prepared by reduction of the corresponding amino acids or amino acid esters using conventional reducing agents such as lithium aluminum hydride and the like.

#### B. Synthesis of Carboxylic Acids

The carboxylic acids of formula **1** are commercially available or they can be prepared by esterification of corresponding alpha amino acids by methods well known in the art. The acid of formula **5**, can be prepared by conventional coupling of an aldehyde of formula **4** with the amino group of an esterified alpha-amino acid under reductive amination reaction conditions described above.

#### 20 Preparation of Cyclic Compounds (e.g. Benzaepinones, Dibenzazepinones, Benzodiazepines and Related Compounds)

The cyclic compounds and amino-substituted derivatives thereof, such as **2**, employed in the reactions described above are either known in the art or can be prepared by art-recognized procedures using commercially available starting materials and reagents.


For example, 5,7-dihydro-6H-dibenz[b,d]azepin-6-one may be prepared by cyclizing a chloromethyl amide intermediate using the procedures set forth in

-- 54 --

R. F. C. Brown et al., *Tetrahedron Letters* 1971, 8, 667-670<sup>12</sup> and references cited therein.

5        Additionally, the synthesis of a representative cyclic compound, i.e., a 5,7-dihydro-6H-dibenz[b,d]azepin-6-one, is illustrated in Scheme 6. As will be readily apparent to those of ordinary skill in the art, the synthetic procedure illustrated in Scheme 6 and the reaction conditions described below can be modified by selecting the appropriate starting materials and reagents to allow the preparation of other cyclic amines suitable for use in this invention.

-- 55 --

Scheme 6

-- 56 --

As shown in Scheme 6, 5,7-dihydro-6H-dibenz[b,d]azepin-6-one derivatives, **23**, wherein  $R^6$ ,  $R^7$ ,  $p$  and  $q$  are as defined above, can be readily prepared in several steps from a 2-bromotoluene derivative **18** and a 2-bromoaniline derivative **20**. In this synthetic procedure, the 2-bromotoluene derivative, **18**, is first converted into the corresponding 2-methylphenylboronate ester, **19**. This reaction is typically conducted by treating **18** with about 1.0 to about 2.1 equivalents of an alkyl lithium reagent, preferably *sec*-butyl lithium or *tert*-butyl lithium, in an inert diluent, such as THF, at a temperature ranging from about -80°C to about -60°C for about 0.25 to about 1 hour. The resulting lithium anion is then treated *in situ* with an excess, preferably 1.5 equivalents, of a trialkylborate, such as trimethylborate. This reaction is initially conducted at -80°C to about -60°C and then allowed to warm to about 0°C to about 30°C for about 0.5 to about 3 hours. The resulting methyl boronate ester is typically not isolated, but is preferably converted *in situ* into the pinacol ester by treating the reaction mixture with an excess, preferably about 2.0 equivalents, of pinacol. This reaction is typically conducted at ambient temperature for about 12 to about 24 hours to afford the 2-methylphenylboronate ester, **19**, in which both  $R^a$  groups are preferably joined together to form  $-C(CH_3)_2C(CH_3)_2-$ .

20

In a separate reaction, the amino group of a 2-bromoaniline derivative, **20**, is converted into the *N*-Boc derivative **21** by treating **20** with about 1.0 to about 1.5 equivalents of di-*tert*-butyl-dicarbonate. Typically, this reaction is conducted at a temperature ranging from 25°C to about 100°C for about 12 to 48 hours to afford the *N*-Boc-2-bromoaniline derivative **21**.

25

As further illustrated in Scheme 6, the 2-methylphenylboronate ester, **19**, and the *N*-Boc-2-bromoaniline derivative **21** can then be coupled to form the biphenyl derivative **22**. This reaction is typically conducted by contacting **21**

-- 57 --

with about 1.0 to about 1.2 equivalents of **19** and about 1.0 to about 1.2 equivalents of potassium carbonate in the presence of a palladium catalyst, preferably tetrakis(triphenylphosphine)palladium(0). Generally, this coupling reaction is conducted in a diluent, preferably 20% water/dioxane, under an inert atmosphere at a temperature ranging from about 50°C to about 100°C for about 6 to 24 hours.

Biphenyl derivative **22** is then readily converted into the 5,7-dihydro-6H-dibenz[b,d]azepin-6-one **23** by carboxylation of the 2-methyl group, followed by cyclization to form the  $\epsilon$ -caprolactam. The carboxylation reaction is typically conducted by contacting **22** with about 2.0 to about 2.5 equivalents of a suitable base, such as *sec*-butyllithium, *tert*-butyllithium and the like, in an inert diluent, such as THF, at a temperature ranging from about -100°C to about -20°C for about 0.5 to 6 hours. The resulting dianion is then treated with excess anhydrous carbon dioxide to form the carboxylate. Treatment of the carboxylate with excess hydrogen chloride in a suitable diluent, such as methanol, at a temperature ranging from about 25°C to about 100°C then affords the 5,7-dihydro-6H-dibenz[b,d]azepin-6-one **23**. Various other cyclic compounds can be prepared by routine modifications of the above described procedures.


20

Preferred synthetic procedures for aminating a representative compound are illustrated in Scheme 7. It will be readily apparent to those of ordinary skill in the art that the synthetic procedure illustrated in Scheme 7 and the following reaction conditions can be modified by selecting the appropriate starting materials and reagents to allow the preparation of other amino compounds suitable for use in this invention.

25

-- 58 --

Scheme 7





-- 59 --

As shown in Scheme 7, 5,7-dihydro-6H-dibenz[b,d]azepin-6-one, **23**, is optionally *N*-alkylated using conventional reagents and conditions to provide a 7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one derivative, **24**. Typically, this reaction is conducted by first contacting **23** with about 1.0 to 1.5 equivalents of a suitable base, such as sodium hydride, sodium bis(trimethylsilyl)amide and the like, in an inert diluent, such as DMF, THF and the like, at a temperature ranging from about -78°C to about 50°C for about 0.25 to about 6 hours. The resulting anion is then treated *in situ* with an excess, preferably about 1.1 to about 2.0 equivalents, of an alkyl, substituted alkyl, cycloalkyl halide, etc., typically a chloride, bromide or iodide. This reaction is typically conducted at a temperature ranging from about 0°C to about 60°C for about 1.0 to about 48 hours to afford the 7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one derivative, **24**.

15

The 7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one **24** is then oximated by contacting **24** with an excess, preferably with about 1.0 to 1.5 equivalents of a suitable base, such as sodium bis(trimethylsilyl)amide and the like, in the presence of about 1.0 to about 2.0 equivalents of an alkyl nitrite. Suitable alkyl nitrites for use in this reaction include, by way of example, butyl nitrite, isoamyl nitrite and the like. This reaction is typically conducted in an inert diluent, such as THF and the like, at a temperature ranging from about -10°C to about 20°C for about 0.5 to about 6 hours to afford the 7-alkyl-5-oximo-5,7-dihydro-6H-dibenz[b,d]azepin-6-one derivative **25**.

20

Reduction of **25** using conventional reagents and conditions then affords the 5-amino-7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one **26**. Preferably, this reduction reaction is conducted by hydrogenating the oxime **25** in the presence of a catalyst, such as Raney nickel. This reaction is typically conducted under

-- 60 --

about 200 psi to about 600 psi of hydrogen at a temperature of about 70°C to about 120°C for about 8 to 48 hours in a diluent, preferably a mixture of ethanol and ammonia (about 20:1). Alternatively, in another preferred procedure, the oxime may be reduced using 10% Pd/C and between about 30 to about 60 psi of hydrogen at a temperature ranging from about 20°C to about 50°C for about 4 hours. The resulting 5-amino-7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one 26 is generally purified using well known procedures, such as recrystallization and/or chromatography.

Alternatively, 5-amino-7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-ones, 26, can be prepared by first forming the 5-iodo derivative 27 of 5,7-dihydro-6H-dibenz[b,d]azepin-6-one, 23. This reaction is typically conducted as described in A. O. King et al.<sup>13</sup> by treating 23 with an excess, preferably about 1.2 to about 2.5 equivalents, of trimethylsilyl iodide in the presence of an excess of a trialkylamine, such as triethylamine, diisopropylethylamine, TMEDA and the like, at a temperature ranging from about -20°C to about 0°C for about 3 to 30 minutes and then adding about 1.1 to about 2.0 equivalents of iodine (I<sub>2</sub>). Typically, after addition of the iodine, the reaction is stirred at a temperature ranging from about 0°C to about 20°C for about 2 to about 4 hours to afford 5-iodo-5,7-dihydro-6H-dibenz[b,d]azepin-6-one, 27.

Displacement of iodide from 27 using an alkali metal azide then affords 5-azido-5,7-dihydro-6H-dibenz[b,d]azepin-6-one, 28. Typically, this reaction is conducted by contacting 27 with about 1.1 to about 1.5 equivalents of sodium azide in an inert diluent, such as DMF, at a temperature ranging from about 0°C to about 50°C for about 12 to about 48 hours.

The azido derivative 28 is then reduced to the corresponding amino derivative 29 using conventional procedures and reagents. For example, the

-- 61 --

azido group is preferably reduced by contacting 28 with an excess, preferably with about 3 equivalents, of triphenylphosphine in a diluent, preferably a mixture of THF and water. This reduction reaction is typically conducted at a temperature ranging from about 0°C to about 50°C for about 12 to 48 hours to afford 5-amino-5,7-dihydro-6H-dibenz[b,d]azepin-6-one, 29.

The amino group of 29 is then protected or blocked using a conventional amino blocking group. Preferably, compound 29 is treated with about 1.0 to about 1.1 equivalents of di-*tert*-butyl dicarbonate in the presence of an excess, preferably about 2 to about 3 equivalents, of a trialkylamine, such as triethylamine. This reaction is typically conducted in an inert diluent, such as THF, at a temperature ranging from about 0°C to about 50°C for 3 to about 24 hours to provide 5-(*N*-Boc-amino)-5,7-dihydro-6H-dibenz[b,d]azepin-6-one, 30.

Compound 30 is then optionally *N*-alkylated to afford, after de-blocking of the amino group, a 5-amino-7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one, 26. The *N*-alkylation reaction is typically conducted by treating 30 with about 1.0 to 1.5 equivalents of an alkyl halide, a substituted alkyl halide or a cycloalkyl halide in the presence of about 1.0 to about 1.5 equivalents of a suitable base, such as cesium carbonate and the like. This reaction is generally conducted in an inert diluent, such as DMF and the like, at a temperature ranging from about 25°C to about 100°C for about 12 to about 48 hours.


Representative alkyl, substituted alkyl and cycloalkyl halides suitable for use in this *N*-alkylation reaction include, by way of illustration, 1-iodo-2-methylpropane, methyl bromoacetate, 1-chloro-3,3-dimethyl-2-butanone, 1-chloro-4-phenylbutane, bromomethylcyclopropane, 1-bromo-2,2,2-trifluoroethane, bromocyclohexane, 1-bromohexane and the like.

-- 62 --

The *N*-Boc protecting group is then removed using conventional procedures and reagents to afford the 5-amino-7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one, **26**. This deblocking reaction is typically conducted by treating the *N*-Boc compound **30** with anhydrous hydrogen chloride in an inert diluent, such as 1,4-dioxane, at a temperature ranging from about 0°C to about 50°C for about 2 to about 8 hours. The resulting 5-amino-7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one **26** is generally purified using well known procedures, such as recrystallization and/or chromatography.

The 5-amino-7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-ones, **26**, can also be prepared via an azide transfer reaction as illustrated in Scheme 8.

Scheme 8



-- 63 --

As shown in Scheme 8, 5,7-dihydro-6H-dibenz[b,d]azepin-6-one, **23**, is first *N*-alkylated as described above using conventional reagents and conditions to provide a 7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one derivative, **24**.

5           The 7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one **24** is then reacted with an azide transfer reagent to afford 5-azido-7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one **31**. Typically, this reaction is conducted by first contacting **24** with an excess, preferably with about 1.0 to 1.5 equivalents of a suitable base, such as lithium diisopropylamine and the like, in an inert diluent  
10           such as THF, at a temperature ranging from about -90°C to about -60°C for about 0.25 to about 2.0 hours. The resulting anion is then treated with an excess, preferably with about 1.1 to about 1.2 equivalents, of an azide transfer reagent, such as 2,4,6-triisopropylbenzenesulfonyl azide (trisyl azide). This reaction is typically conducted at a temperature ranging from about -90°C to  
15           about -60°C for about 0.25 to about 2.0 hours. The reaction mixture is then typically treated with an excess of glacial acetic acid and the mixture is allowed to warm to ambient temperature and then heated at about 35°C to about 50°C for about 2 to 4 hours to afford the 5-azido-7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one derivative **31**. Reduction of **31** as described above using  
20           conventional reagents and conditions then affords the 5-amino-7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one **26**.

          If desired, the aryl rings of 5-amino-7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-ones, **26**, and similar or related compounds may be partially  
25           or fully saturated by treatment with hydrogen in the presence of a hydrogenation catalyst. Typically, this reaction is conducted by treating **26** with hydrogen at a pressure of about 10 to about 100 psi in the presence of a catalyst, such as rhodium on carbon. This reaction is typically conducted at a temperature

-- 64 --

ranging from about 20°C to about 100°C for about 12 to 96 hours in a suitable diluent, such as ethyl acetate/acetic acid (1:1) and the like.

Other methods for preparing intermediates useful in this invention are described in U.S. Patent Application No. 09/102,726, filed on June 22, 1998, and entitled "Polycyclic  $\alpha$ -Amino- $\epsilon$ -caprolactams and Related Compounds", the disclosure of which is incorporated herein by reference in its entirety.

Additionally, the synthesis of various benzapinones and related compounds are described in Busacca et al., *Tetrahedron Lett.*, **33**, 165-168 (1992); Crosisier et al., U.S. Patent No. 4,080,449; J. A. Robl et al. *Tetrahedron Lett.*, **36(10)**, 1593-1596 (1995); Flynn et al. *J. Med. Chem.* **36**, 2420-2423 (1993); Orito et al. *Tetrahedron*, **36**, 1017-1021 (1980); Kawase et al., *J. Org. Chem.*, **54**, 3394-3403 (1989); Lowe et al., *J. Med. Chem.* **37**, 3789-3811 (1994); Robl et al., *Bioorg. Med. Chem. Lett.*, **4**, 1789-1794 (1994); Skiles et al., *Bioorg. Med. Chem. Lett.*, **3**, 773-778 (1993); Grunewald et al., *J. Med. Chem.*, **39(18)**, 3539 (1996); Warshawsky et al., *Bioorg. Med. Chem. Lett.*, **6**, 957-962 (1996); Ben-Ishai, et al., *Tetrahedron*, **43**, 439-450 (1987); van Neil et al. *Bioorg. Med. Chem.* **5**, 1421-1426 (1995); and references cited therein. These publications and patents are incorporated herein by reference in their entirety.

Similarly, various benzodiazepine derivatives suitable for use in this invention can be prepared using conventional procedures and reagents. For example, a 2-aminobenzophenone can be readily coupled to  $\alpha$ -(isopropylthio)-N-(benzyloxycarbonyl)glycine by first forming the acid chloride of the glycine derivative with oxalyl chloride, and then coupling the acid chloride with the 2-aminobenzophenone in the presence of a base, such as 4-methylmorpholine, to afford the 2-[ $\alpha$ -(isopropylthio)-N-(benzyloxycarbonyl)glyciny]-

-- 65 --

aminobenzophenone. Treatment of this compound with ammonia gas in the presence of an excess, preferably about 1.1 to about 1.5 equivalents, of mercury (II) chloride then affords the 2-[N-( $\alpha$ -amino)-N'-(benzyloxycarbonyl)-glyciny]aminobenzophenone. This intermediate can then be readily cyclized by treatment with glacial acetic acid and ammonium acetate to provide the 3-(benzyloxycarbonyl)amino-2,3-dihydro-5-phenyl-1H-1,4-benzodiazepin-2-one. Subsequent removal of the Cbz group affords the 3-amino-2,3-dihydro-5-phenyl-1H-1,4-benzodiazepin-2-one.

Alternatively, 2,3-dihydro-5-phenyl-1H-1,4-benzodiazepin-2-ones can be readily aminated at the 3-position using conventional azide transfer reactions followed by reduction of the resulting azido group to form the corresponding amino group. The conditions for these and related reactions are described in the examples set forth below. Additionally, 2,3-dihydro-5-phenyl-1H-1,4-benzodiazepin-2-ones are readily alkylated at the 1-position using conventional procedures and reagents. For example, this reaction is typically conducted by first treating the benzodiazepinone with about 1.1 to about 1.5 equivalents of a base, such as sodium hydride, potassium *tert*-butoxide, potassium 1,1,1,3,3,3-hexamethyldisilazane, or cesium carbonate, in an inert diluent, such as DMF. This reaction is typically conducted at a temperature ranging from about -78°C to about 80°C for about 0.5 to about 6 hours. The resulting anion is then contacted with an excess, preferably about 1.1 to about 3.0 equivalents, of an alkyl halide, typically an alkyl chloride, bromide or iodide. Generally, this reaction is conducted at a temperature of about 0°C to about 100°C for about 1 to about 48 hours.

Additionally, the 3-amino-2,4-dioxo-2,3,4,5-tetrahydro-1H-1,5-benzodiazepines employed in this invention are typically prepared by first coupling malonic acid with a 1,2-phenylenediamine. Conditions for this reaction

-- 66 --

are well known in the art and are described, for example, in PCT Application WO 96-US8400 960603. Subsequent alkylation and amination using conventional procedures and reagents affords various 3-amino-1,5-bis(alkyl)-2,4-dioxo-2,3,4,5-tetrahydro-1H-1,5-benzodiazepines. Such procedures are  
5 described in further detail in the examples set forth below.

In the synthesis of compounds of formula I using the synthetic methods described herein, the starting materials can contain a chiral center (e.g., alanine) and, when a racemic starting material is employed, the resulting product is a  
10 mixture of R,S enantiomers. Alternatively, a chiral isomer of the starting material can be employed and, if the reaction protocol employed does not racemize this starting material, a chiral product is obtained. Such reaction protocols can involve inversion of the chiral center during synthesis.

15 Accordingly, unless otherwise indicated, the products of this invention are a mixture of R,S enantiomers. Preferably, however, when a chiral product is desired, the chiral product corresponds to the L-amino acid derivative. Alternatively, chiral products can be obtained via purification techniques which separate enantiomers from a R,S mixture to provide for one or the other  
20 stereoisomer. Such techniques are well known in the art.

#### Pharmaceutical Formulations

When employed as pharmaceuticals, the compounds of formula I are usually administered in the form of pharmaceutical compositions. These  
25 compounds can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous, intravenous, intramuscular, and intranasal. These compounds are effective as both injectable and oral compositions. Such compositions are prepared in a manner well known in the pharmaceutical art and comprise at least one active compound.



-- 67 --

This invention also includes pharmaceutical compositions which contain, as the active ingredient, one or more of the compounds of formula I above associated with pharmaceutically acceptable carriers. In making the compositions of this invention, the active ingredient is usually mixed with an excipient, diluted by an excipient or enclosed within a carrier which can be in the form of a capsule, sachet, paper or other container. When the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material which acts as a vehicle, carrier or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders.

In preparing a formulation, it may be necessary to mill the active compound to provide the appropriate particle size prior to combining with the other ingredients. If the active compound is substantially insoluble, it ordinarily is milled to a particle size of less than 200 mesh. If the active compound is substantially water soluble, the particle size is normally adjusted by milling to provide a substantially uniform distribution in the formulation, e.g. about 40 mesh.

Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, sterile water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxy-benzoates; sweetening

-- 68 --

agents; and flavoring agents. The compositions of the invention can be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures known in the art.

5

The compositions are preferably formulated in a unit dosage form, each dosage containing from about 5 to about 100 mg, more usually about 10 to about 30 mg, of the active ingredient. The term "unit dosage forms" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient. Preferably, the compound of formula I above is employed at no more than about 20 weight percent of the pharmaceutical composition, more preferably no more than about 15 weight percent, with the balance being pharmaceutically inert carrier(s).

10

15

20

The active compound is effective over a wide dosage range and is generally administered in a pharmaceutically effective amount. It, will be understood, however, that the amount of the compound actually administered will be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

25

For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical excipient to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly

-- 69 --

throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation is then subdivided into unit dosage forms of the type described above containing from, for example, 0.1 to about 500 mg of the active  
5 ingredient of the present invention.

The tablets or pills of the present invention may be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer  
10 dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permit the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric  
15 acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol, and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include aqueous  
20 solutions suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil, or peanut oil, as well as elixirs and similar pharmaceutical vehicles.

Compositions for inhalation or insufflation include solutions and  
25 suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as described *supra*. Preferably the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions in preferably pharmaceutically acceptable

-- 70 --

solvents may be nebulized by use of inert gases. Nebulized solutions may be breathed directly from the nebulizing device or the nebulizing device may be attached to a face mask, tent, or intermittent positive pressure breathing machine. Solution, suspension, or powder compositions may be administered, preferably orally or nasally, from devices which deliver the formulation in an appropriate manner.

The following formulation examples illustrate the pharmaceutical compositions of the present invention.

10

Formulation Example 1

Hard gelatin capsules containing the following ingredients are prepared:

	<u>Ingredient</u>	<u>Quantity</u> <u>(mg/capsule)</u>
15	Active Ingredient	30.0
	Starch	305.0
	Magnesium stearate	5.0

20 The above ingredients are mixed and filled into hard gelatin capsules in 340 mg quantities.

Formulation Example 2

A tablet formula is prepared using the ingredients below:

25	<u>Ingredient</u>	<u>Quantity</u> <u>(mg/tablet)</u>
	Active Ingredient	25.0
	Cellulose, microcrystalline	200.0
30	Colloidal silicon dioxide	10.0
	Stearic acid	5.0

The components are blended and compressed to form tablets, each weighing 240 mg.

35

-- 71 --

Formulation Example 3

A dry powder inhaler formulation is prepared containing the following components:

	<u>Ingredient</u>	<u>Weight %</u>
5	Active Ingredient	5
	Lactose	95

10        The active ingredient is mixed with the lactose and the mixture is added to a dry powder inhaling appliance.

Formulation Example 4

15        Tablets, each containing 30 mg of active ingredient, are prepared as follows:

	<u>Ingredient</u>	<u>Quantity (mg/tablet)</u>
	Active Ingredient	30.0 mg
20	Starch	45.0 mg
	Microcrystalline cellulose	35.0 mg
	Polyvinylpyrrolidone (as 10% solution in sterile water)	4.0 mg
	Sodium carboxymethyl starch	4.5 mg
25	Magnesium stearate	0.5 mg
	Talc	<u>1.0 mg</u>
	Total	120.0 mg

30        The active ingredient, starch and cellulose are passed through a No. 20 mesh U.S. sieve and mixed thoroughly. The solution of polyvinyl-pyrrolidone is mixed with the resultant powders, which are then passed through a 16 mesh U.S. sieve. The granules so produced are dried at 50° to 60°C and passed through a 16 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate, and  
35        talc, previously passed through a No. 30 mesh U.S. sieve, are then added to the

-- 72 --

granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

#### Formulation Example 5

5 Capsules, each containing 40 mg of medicament are made as follows:

	<u>Ingredient</u>	<u>Quantity (mg/capsule)</u>
	Active Ingredient	40.0 mg
10	Starch	109.0 mg
	Magnesium stearate	<u>1.0 mg</u>
	Total	150.0 mg

The active ingredient, starch, and magnesium stearate are blended, passed  
15 through a No. 20 mesh U.S. sieve, and filled into hard gelatin capsules in 150 mg quantities.

#### Formulation Example 6

Suppositories, each containing 25 mg of active ingredient are made as  
20 follows:

	<u>Ingredient</u>	<u>Amount</u>
	Active Ingredient	25 mg
25	Saturated fatty acid glycerides to	2.000 mg

The active ingredient is passed through a No. 60 mesh U.S. sieve and  
suspended in the saturated fatty acid glycerides previously melted using the  
minimum heat necessary. The mixture is then poured into a suppository mold of  
30 nominal 2.0 g capacity and allowed to cool.

-- 73 --

Formulation Example 7

Suspensions, each containing 50 mg of medicament per 5.0 mL dose are made as follows:

	<u>Ingredient</u>	<u>Amount</u>
5	Active Ingredient	50.0 mg
	Xanthan gum	4.0 mg
	Sodium carboxymethyl cellulose (11 %)/ Microcrystalline cellulose (89 %)	50.0 mg
10	Sucrose	1.75 g
	Sodium benzoate	10.0 mg
	Flavor and Color	q.v.
	Purified water to	5.0 mL

15 The active ingredient, sucrose and xanthan gum are blended, passed through a No. 10 mesh U.S. sieve, and then mixed with a previously made solution of the microcrystalline cellulose and sodium carboxymethyl cellulose in water. The sodium benzoate, flavor, and color are diluted with some of the water and added with stirring. Sufficient water is then added to produce the required volume.

20

Formulation Example 8

	<u>Ingredient</u>	<u>Quantity (mg/capsule)</u>
25	Active Ingredient	15.0 mg
	Starch	407.0 mg
	Magnesium stearate	<u>3.0 mg</u>
	Total	425.0 mg

30

The active ingredient, starch, and magnesium stearate are blended, passed through a No. 20 mesh U.S. sieve, and filled into hard gelatin capsules in 560 mg quantities.

-- 74 --

Formulation Example 9

A subcutaneous formulation may be prepared as follows:

	<u>Ingredient</u>	<u>Quantity</u>
5	Active Ingredient	1.0 mg
	corn oil	1 mL

(Depending on the solubility of the active ingredient in corn oil, up to about 5.0 mg or more of the active ingredient may be employed in this formulation, if  
10 desired).

Formulation Example 10

A topical formulation may be prepared as follows:

	<u>Ingredient</u>	<u>Quantity</u>
15	Active Ingredient	1-10 g
	Emulsifying Wax	30 g
	Liquid Paraffin	20 g
20	White Soft Paraffin	to 100 g

The white soft paraffin is heated until molten. The liquid paraffin and emulsifying wax are incorporated and stirred until dissolved. The active ingredient is added and stirring is continued until dispersed. The mixture is then cooled until solid.

25  
Another preferred formulation employed in the methods of the present invention employs transdermal delivery devices ("patches"). Such transdermal patches may be used to provide continuous or discontinuous infusion of the compounds of the present invention in controlled amounts. The construction and  
30 use of transdermal patches for the delivery of pharmaceutical agents is well known in the art. See, e.g., U.S. Patent 5,023,252, issued June 11, 1991, herein incorporated by reference. Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.



-- 75 --

Frequently, it will be desirable or necessary to introduce the pharmaceutical composition to the brain, either directly or indirectly. Direct techniques usually involve placement of a drug delivery catheter into the host's ventricular system to bypass the blood-brain barrier. One such implantable delivery system used for the transport of biological factors to specific anatomical regions of the body is described in U.S. Patent 5,011,472 which is herein incorporated by reference.

Indirect techniques, which are generally preferred, usually involve formulating the compositions to provide for drug latentiation by the conversion of hydrophilic drugs into lipid-soluble drugs. Latentiation is generally achieved through blocking of the hydroxy, carbonyl, carboxyl, sulfate, and primary amine groups present on the drug to render the drug more lipid soluble and amenable to transportation across the blood-brain barrier. Alternatively, the delivery of hydrophilic drugs may be enhanced by intra-arterial infusion of hypertonic solutions which can transiently open the blood-brain barrier.

Other suitable formulations for use in the present invention can be found in *Remington's Pharmaceutical Sciences*, Mace Publishing Company, Philadelphia, PA, 17th ed. (1985).

20

### Utility

The compounds and pharmaceutical compositions of the invention are useful in inhibiting  $\beta$ -amyloid peptide release and/or its synthesis; and, accordingly, have utility in diagnosing and treating Alzheimer's disease in mammals including humans.

25

As noted above, the compounds described herein are suitable for use in a variety of drug delivery systems described above. Additionally, in order to enhance the *in vivo* serum half-life of the administered compound, the

-- 76 --

compounds may be encapsulated, introduced into the lumen of liposomes, prepared as a colloid, or other conventional techniques may be employed which provide an extended serum half-life of the compounds. A variety of methods are available for preparing liposomes, as described in, e.g., Szoka, et al., U.S. Patent Nos. 4,235,871, 4,501,728 and 4,837,028, each of which is incorporated herein by reference.

The amount of compound administered to the patient will vary depending upon what is being administered, the purpose of the administration, such as prophylaxis or therapy, the state of the patient, the manner of administration, and the like. In therapeutic applications, compositions are administered to a patient already suffering from AD in an amount sufficient to at least partially arrest further onset of the symptoms of the disease and its complications. An amount adequate to accomplish this is defined as a "therapeutically effective dose." Amounts effective for this use will depend on the judgment of the attending clinician depending upon factors such as the degree or severity of AD in the patient, the age, weight and general condition of the patient, and the like. Preferably, for use as therapeutics, the compounds described herein are administered at dosages ranging from about 1 to about 500 mg/kg/day.

In prophylactic applications, compositions are administered to a patient at risk of developing AD (determined for example by genetic screening or familial trait) in an amount sufficient to inhibit the onset of symptoms of the disease. An amount adequate to accomplish this is defined as "prophylactically effective dose." Amounts effective for this use will depend on the judgment of the attending clinician depending upon factors such as the age, weight and general condition of the patient, and the like. Preferably, for use as prophylactics, the compounds described herein are administered at dosages ranging from about 1 to about 500 mg/kg/day.

-- 77 --

As noted above, the compounds administered to a patient are in the form of pharmaceutical compositions described above. These compositions may be sterilized by conventional sterilization techniques, or may be sterile filtered. The resulting aqueous solutions may be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile aqueous carrier prior to administration. The pH of the compound preparations typically will be between 3 and 11, more preferably from 5 to 9 and most preferably from 7 to 8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of pharmaceutical salts.

10

The compounds described herein are also suitable for use in the administration of the compounds to a cell for diagnostic and drug discovery purposes. Specifically, the compounds may be used in the diagnosis of cells releasing and/or synthesizing  $\beta$ -amyloid peptide. In addition the compounds described herein are useful for the measurement and evaluation of the activity of other candidate drugs on the inhibition of the cellular release and/or synthesis of  $\beta$ -amyloid peptide.

15

The following synthetic and biological examples are offered to illustrate this invention and are not to be construed in any way as limiting the scope of this invention.

20

### EXAMPLES

In the examples below, the following abbreviations have the following meanings. If an abbreviation is not defined, it has its generally accepted meaning.

25

BEMP	=	2-tert-butylimino-2-diethylamino-1,3-dimethylperhydro-1,3,2-diazaphosphorine
Boc	=	<i>t</i> -butoxycarbonyl

-- 78 --

	BOP	=	benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate
	bd	=	broad doublet
	bs	=	broad singlet
5	bt	=	broad triplet
	CBZ	=	benzyloxycarbonyl
	d	=	doublet
	dd	=	doublet of doublets
	DIC	=	diisopropylcarbodiimide
10	DIPEA	=	diisopropylethylamine
	DMF	=	dimethylformamide
	DMAP	=	dimethylaminopyridine
	DMSO	=	dimethylsulfoxide
	EDC	=	ethyl-1-(3-dimethylaminopropyl)carbodiimide
15	ee	=	enantiomeric excess
	eq.	=	equivalents
	EtOAc	=	ethyl acetate
	g	=	grams
	HMDS	=	1,1,1,3,3,3-hexamethyldisilazane
20	HOBT	=	1-hydroxybenzotriazole hydrate
	Hunig's base	=	diisopropylethylamine
	L	=	liter
	LDA	=	lithium diisopropylamide
	m	=	multiplet
25	M	=	molar
	max	=	maximum
	meq	=	milliequivalent
	mg	=	milligram
	mL	=	milliliter

-- 79 --

	mm	=	millimeter
	mmol	=	millimole
	MOC	=	methoxyoxycarbonyl
	N	=	normal
5	N/A	=	not available
	ng	=	nanogram
	nm	=	nanometers
	OD	=	optical density
	PEPC	=	1-(3-(1-pyrrolidiny)propyl)-3-ethylcarbodiimide
10	PP-HOBT	=	piperidine-piperidine-1-hydroxybenzotrizole
	psi	=	pounds per square inch
	φ	=	phenyl
	q	=	quartet
	quint.	=	quintet
15	rpm	=	rotations per minute
	RT	=	room temperature
	s	=	singlet
	sat.	=	saturated
	t	=	triplet
20	t-BuOK	=	potassium <i>tert</i> -butoxide
	TFA	=	trifluoroacetic acid
	THF	=	tetrahydrofuran
	tlc	=	thin layer chromatography
	TMSI	=	trimethylsilyl iodide
25	μL	=	microliter
	UV	=	ultra-violet

-- 80 --

In the examples below, all temperatures are in degrees Celsius (unless otherwise indicated). The compounds set forth in the examples below were prepared using the following general procedures as indicated.

- 5        In the following examples and procedures, the term "Aldrich" indicates that the compound or reagent used in the procedure is commercially available from Aldrich Chemical Company, Inc., 1001 West Saint Paul Avenue, Milwaukee, WI 53233, USA; the term "Fluka" indicates that the compound or reagent is commercially available from Fluka Chemical Corp., 980 South 2nd Street, Ronkonkoma, NY 11779, USA; the term "Lancaster" indicates that the compound or reagent is commercially available from Lancaster Synthesis, Inc., P.O. Box 100 Windham, NH 03087, USA; the term "Sigma" indicates that the compound or reagent is commercially available from Sigma, P.O. Box 14508, St. Louis, MO 63178, USA; the term "Chemservice" indicates that the compound or reagent is commercially available from Chemservice Inc., Westchester, PA, USA; the term "Bachem" indicates that the compound or reagent is commercially available from Bachem Biosciences Inc., 3700 Horizon Drive, Renaissance at Gulph Mills, King of Prussia, PA 19406, USA; the term "Maybridge" indicates that the compound or reagent is commercially available from Maybridge Chemical Co. Trevillet, Tintagel, Cornwall PL34 OHW, United Kingdom; the term "TCI" indicates that the compound or reagent is commercially available from TCI America, 9211 North Harborage Street, Portland OR 97203, USA; the term "Alfa" indicates that the compound or reagent is commercially available from Johnson Matthey Catalog Company, Inc. 30 Bond Street, Ward Hill, MA 01835-0747, USA; the term "Novabiochem" indicates that the compound or reagent is commercially available from Calbiochem-Novabiochem Corp., 10933 North Torrey Pines Road, P.O. Box 12087, La Jolla, CA 92039-2087, USA; the term "Oakwood" indicates that the compound or reagent is commercially available from Oakwood, Columbia, South Carolina, USA; the term "Advanced Chemtech" indicates that the

-- 81 --

compound or reagent is commercially available from Advanced Chemtech, Louisville, KY, USA; and the term "Pfaltz & Bauer" indicates that the compound or reagent is commercially available from Pfaltz & Bauer, Waterbury, CT, USA.

5    **I.    Coupling Procedures**

The following coupling procedures may be used to prepare compounds of this invention:

**GENERAL PROCEDURE A**

10       First EDC Coupling Procedure

To a 1:1 mixture of the corresponding carboxylic acid and the corresponding amino acid ester or amide in  $\text{CH}_2\text{Cl}_2$  at  $0^\circ\text{C}$  was added 1.5 equivalents triethylamine, followed by 2.0 equivalents hydroxybenzotriazole monohydrate and then 1.25 equivalents of ethyl-3-(3-dimethylamino)propyl carbodiimide.HCl. The reaction mixture was stirred overnight at room temperature and then transferred to a separatory funnel. The mixture was washed with water, saturated aqueous  $\text{NaHCO}_3$ , 1N HCl and saturated aqueous NaCl, and then dried over  $\text{MgSO}_4$ . The resulting solution was stripped free of solvent on a rotary evaporator to yield the crude product.

20

**GENERAL PROCEDURE B**

Second EDC Coupling Procedure

A mixture of the corresponding acid (1 eqv), N-1-hydroxybenzotriazole (1.6 eqv), the corresponding amine (1 eqv), N-methylmorpholine ( 3 eqv) and dichloromethane (or DMF for insoluble substrates) was cooled in an ice-water bath and stirred until a clear solution was obtained. EDC (1.3 eqv) was then added to the reaction mixture. The cooling bath was then allowed to warm to ambient temperature over 1-2 h and the reaction mixture was stirred overnight. The reaction mixture was then evaporated to dryness under vacuum. To the

25

-- 82 --

residue was added 20% aqueous potassium carbonate and the mixture was shaken thoroughly and then allowed to stand until the oily product solidified (overnight if necessary). The solid product was then collected by filtration, washed thoroughly with 20% aqueous potassium carbonate, water, 10% HCl, and water to  
5 give the product, usually in pure state. No racemization was observed.

### GENERAL PROCEDURE C

#### Third EDC Coupling Procedure

The carboxylic acid was dissolved in methylene chloride. The  
10 corresponding amino acid ester or amide (1 eq.), N-methylmorpholine (5 eq.) and hydroxybenzotriazole monohydrate (1.2 eq.) were added in sequence. A cooling bath was applied to the round bottomed flask until the solution reached 0°C. At that time, 1.2 eq. of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride was added. The solution was allowed to stir overnight and come to room  
15 temperature under nitrogen pressure. The reaction mixture was worked up by washing the organic phase with saturated aqueous sodium carbonate, 0.1M citric acid, and brine before drying with sodium sulfate. The solvents were then removed to yield crude product.

20

### GENERAL PROCEDURE D

#### Fourth EDC Coupling Procedure

A round bottom flask was charged with the corresponding carboxylic acid (1.0 eq.), hydroxybenzotriazole hydrate (1.1 eq.) and the corresponding amine (1.0 eq.) in THF under nitrogen atmosphere. An appropriate amount (1.1 eq for  
25 free amines and 2.2 eq. for hydrochloride amine salts) of base, such as Hunig's base was added to the well stirred mixture followed by EDC (1.1 eq.). After stirring from 4 to 17 hours at room temperature the solvent was removed at reduced pressure, the residue taken up in ethyl acetate (or similar solvent) and water, washed with saturated aqueous sodium bicarbonate solution, 1 N HCl,



-- 83 --

brine, dried over anhydrous sodium sulfate and the solvent removed at reduced pressure to provide the product.

### GENERAL PROCEDURE E

5

#### BOP Coupling Procedure

To a stirred solution of *N*-(3,5-difluorophenylacetyl)alanine (2 mmol) in DMF, cooled in an ice-water bath, was added BOP (2.4 mmol) and *N*-methylmorpholine (6 mmol). The reaction mixture was stirred for 50 min. and then a solution of  $\alpha$ -amino- $\gamma$ -lactam (2 mmol) in DMF cooled at 0 °C was added.

10 The cooling bath was allowed to warm to ambient temperature over 1-2 h and the reaction mixture was then stirred overnight. A 20% aqueous potassium carbonate solution (60 mL) was added and this mixture shaken thoroughly. No solid formed. The mixture was then washed with ethyl acetate (150 mL) and evaporated to dryness under vacuum to give a white solid. Water (50 mL) was then added and

15 this mixture shaken thoroughly. The precipitate that formed was collected by filtration, then washed thoroughly with water, followed by 1 mL of diethyl ether to give the product (51 mg, 0.16 mmol, 7.8%).

### GENERAL PROCEDURE F

20

#### Coupling of an Acid Chloride with an Amino Acid Ester

To a stirred solution of (D,L)-alanine isobutyl ester hydrochloride (4.6 mmol) in 5 ml of pyridine was added 4.6 mmol of the acid chloride. Precipitation occurred immediately. The mixture was stirred for 3.5 h, dissolved in 100 mL of diethyl ether, washed with 10% HCl three times, brine once, 20% potassium

25 carbonate once and brine once. The solution was dried over magnesium sulfate, filtered, and evaporated to yield the product. Other amino acid esters may also be employed in this procedure.

-- 84 --

**GENERAL PROCEDURE G**Coupling of a Carboxylic Acid with an Amino Acid Ester

A solution of the carboxylic acid (3.3 mmol) and 1,1'-carbodiimidazole (CDI) in 20 mL THF was stirred for 2 h. (D,L)-alanine isobutyl ester  
5 hydrochloride (3.6 mmol) was added, followed by 1.5 mL (10.8 mmol) of triethylamine. The reaction mixture was stirred overnight. The reaction mixture was dissolved in 100 mL of diethyl ether, washed with 10% HCl three times, brine once, 20% potassium carbonate once and brine once. The solution was dried over magnesium sulfate, filtered, and evaporated to yield the product. Other  
10 amino acid esters may also be employed in this procedure.

**GENERAL PROCEDURE H**Fifth EDC Coupling Procedure

In a round bottom flask was added a carboxylic acid (1.1 eq.) in THF, an  
15 amine hydrochloride (1.0 eq.), 1-hydroxybenzotriazole hydrate (1.1 eq.), N,N-diisopropylethylamine (2.1 eq.), followed by 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (1.1 eq.). The reaction mixture was stirred at room temperature for 10-20 hours under an atmosphere of nitrogen. The mixture was diluted with EtOAc and washed with 0.1 M HCl (1 x 10 mL),  
20 saturated NaHCO<sub>3</sub> (1 x 10 mL), H<sub>2</sub>O (1 x 10 mL), and brine and dried over MgSO<sub>4</sub>. The drying agent was removed by filtration and the filtrate was concentrated *in vacuo*. The residue was purified by flash column chromatography on silica gel followed by trituration from EtOAc and hexanes.

25

**GENERAL PROCEDURE I**Sixth EDC Coupling Procedure

To a solution or suspension of the amine or amine hydrochloride (1.0 eq.) in THF (0.05-0.1 M) under N<sub>2</sub> at 0°C was added the carboxylic acid (1.0-1.1 eq.), hydroxybenzotriazole monohydrate (1.1-1.15 eq.), Hunig's base (1.1 eq. for free

-- 85 --

amines and 1.1-2.3 eq. for hydrochloride amine salts), followed by 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.1-1.15 eq.). The cooling bath was removed and the mixture allowed to warm to room temperature for 10-24 hours. The solution or mixture was diluted with EtOAc, in a 3-5  
5 volume multiple of the initial THF volume, and washed with 0.1-1.0 M aq. HCl (1 or 2x), dilute NaHCO<sub>3</sub> (1 or 2x), and brine (1x). Then, the organic phase was dried over either MgSO<sub>4</sub> or Na<sub>2</sub>SO<sub>4</sub>, filtered, concentrated to provide the crude product, which was either further purified or utilized without further purification.

10

## GENERAL PROCEDURE J

### EEDQ Coupling Procedure

To a solution of the amine in THF (1.0 eq., 0.05-0.08 M, final molarity) under N<sub>2</sub> at room temperature was added the N-t-Boc protected amino acid (1.1 eq., either as a solid or in THF via cannula), followed by EEDQ (Aldrich, 1.1  
15 eq.). The pale yellow solution was stirred at room temperature for 16-16.5 hours, then diluted with EtOAc (in a 3-5 volume multiple of the initial THF volume), and washed with 1M aq. HCl (2x), dilute aq. NaHCO<sub>3</sub> (2x), and brine (1x). The organic phase was dried over either Na<sub>2</sub>SO<sub>4</sub> or MgSO<sub>4</sub>, filtered, and concentrated.

20

## II. Carboxylic Acids

The following procedures may be used to prepare carboxylic acid intermediates useful in the present invention:

25

## GENERAL PROCEDURE II-A

### Ester Hydrolysis to Free Acid

Ester hydrolysis to the free acid was conducted by conventional methods. Below are two examples of such conventional de-esterification methods.

#### Method A:

-- 86 --

To a carboxylic ester compound in a 1:1 mixture of CH<sub>3</sub>OH/H<sub>2</sub>O was added 2-5 equivalents of K<sub>2</sub>CO<sub>3</sub>. The mixture was heated to 50°C for 0.5 to 1.5 hours until tlc showed complete reaction. The reaction was cooled to room temperature and the methanol was removed on a rotary evaporator. The pH of the remaining aqueous solution was adjusted to ~2, and ethyl acetate was added to extract the product. The organic phase was then washed with saturated aqueous NaCl and dried over MgSO<sub>4</sub>. The solution was stripped free of solvent on a rotary evaporator to yield the product.

Method B:

The amino acid ester was dissolved in dioxane/water (4:1) to which was added LiOH (~2 eq.) that was dissolved in water such that the total solvent after addition was about 2:1 dioxane:water. The reaction mixture was stirred until reaction completion and the dioxane was removed under reduced pressure. The residue was dissolved in water and washed with ether. The layers were separated and the aqueous layer was acidified to pH 2. The aqueous layer was extracted with ethyl acetate. The ethyl acetate extracts were dried over Na<sub>2</sub>SO<sub>4</sub> and the solvent was removed under reduced pressure after filtration. The residue was purified by conventional methods (e.g., recrystallization).

20

**GENERAL PROCEDURE II-B**

Acid Chloride Preparation

3,5-Difluorophenylacetic acid (30 g, 0.174 mol) (Aldrich) was dissolved in dichloromethane and this solution was cooled to 0°C. DMF (0.5 mL, catalytic) was added followed by the dropwise addition of oxalyl chloride (18 mL, 0.20 mol) over a 5 minute period. The reaction was stirred for 3 h and then rotoevaporated at reduced pressure to give an oil which was placed on a high vacuum pump for 1 h to afford 3,5-difluorophenylacetyl chloride as a thin yellow oil. Other acid chlorides can be prepared in a similar manner.

-- 87 --

**GENERAL PROCEDURE II-C**Schotten-Baumann Procedure

3,5-Difluorophenylacetyl chloride (from General Procedure II-B) was added dropwise to a 0°C solution of L-alanine (Aldrich) (16.7 g, 0.187 mol) in 2 N sodium hydroxide (215 mL, 0.43 mol). The reaction was stirred for 1 h at 0°C and then overnight at room temperature. The reaction was diluted with water (100 mL), then extracted with ethyl acetate (3 x 150 mL). The organic layer was then washed with brine (200 mL), dried over MgSO<sub>4</sub>, and rotoevaporated at reduced pressure to a residue. Recrystallization of the residue from ethyl acetate/hexanes afforded the desired product (34.5 g, 82% yield). Other acid chlorides may be used in this procedure to provide for intermediates useful in this invention.

**GENERAL PROCEDURE II-D**Reductive Amination

To a solution of the arylamine in ethanol in a hydrogenation flask was added 1 equivalent of the 2-oxocarboxylic acid ester (e.g., pyruvate ester), followed by 10% palladium on carbon (25 weight percent based on the arylamine). The reaction was hydrogenated at 20 psi H<sub>2</sub> on a Parr shaker until complete reaction was indicated by tlc (30 minutes to 16 hours). The reaction mixture was then filtered through a pad of Celite 545 (available from Aldrich Chemical Company, Inc.) and stripped free of solvent on a rotary evaporator. The crude product residue was then further purified via chromatography.

**III. Cyclic Compounds**

The following procedures illustrate the synthesis of various cyclic compound intermediates useful for preparing compounds of this invention:

**A. Benzazepinone Derivatives and Related Compounds****GENERAL PROCEDURE 1-A**

-- 88 --

Alkylation of 1-Amino-1,3,4,5-tetrahydro-2H-3-benzazepin-2-oneStep A

1-Ethoxycarbonylamino-1,3,4,5-tetrahydro-2H-3-benzazepin-2-one was prepared according to the procedure of Ben-Ishai et al., *Tetrahedron*, 1987, 43, 430, incorporated herein by reference.

Step B

1-Ethoxycarbonylamino-1,3,4,5-tetrahydro-2H-3-benzazepin-2-one (2.0 g, 100 M%) was dissolved in DMF (30 mL) and NaH (95 %, 0.17 g, 100M %) was added in one portion. The reaction mixture was stirred for 1 hour and then the appropriate alkyl iodide (300M%) was added and the mixture was stirred for 12 hours. The reaction was poured into water and extracted with ethyl acetate (3x). The ethyl acetate extracts were then washed with water (3x) and brine (1x). Treatment with MgSO<sub>4</sub>, rotoevaporation, and chromatography (30% EtOAc/hexanes) yielded 1-ethoxycarbonylamino-3-alkyl-1,3,4,5-tetrahydro-2H-3-benzazepin-2-one in 87 % yield.

Step C

1-Ethoxycarbonylamino-3-alkyl-1,3,4,5-tetrahydro-2H-3-benzazepin-2-one (1.0g, 100M%) was suspended in 30 mL of 30% HBr/HOAc and heated to 100°C. The reaction mixture was stirred for 5 hours at this temperature and then the reaction was cooled and rotoevaporated to yield 1-amino-3-alkyl-1,3,4,5-tetrahydro-2H-3-benzazepin-2-one as the hydrobromide salt (100% yield).

**GENERAL PROCEDURE 1-B**Alkylation of 3-Amino-1,3,4,5-tetrahydro-2H-1-benzazepin-2-oneStep A

3-Amino-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one was prepared from  $\alpha$ -tetralone using the methods described in Armstrong et al. *Tetrahedron Letters*, 1994, 35, 3239, incorporated herein by reference. The following compounds were as prepared by this procedure for use in the following steps:

-- 89 --

5-methyl-3-amino-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one (from 4-methyl- $\alpha$ -tetralone (Aldrich)); and

5,5-dimethyl-3-amino-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one (from 4,4-dimethyl- $\alpha$ -tetralone (Aldrich)).

5 Step B

3-Amino-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one (4.43 g, 100M%) was suspended in t-butanol (30mL) and BOC-anhydride (7.5 mL, 130M%) was added dropwise. The reaction mixture was stirred for 2 hours and then it was rotoevaporated to a residue which was chromatographed with 60% ethyl acetate/hexanes to yield BOC-protected 3-amino-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one in 87% yield.

Step C

BOC-protected 3-amino-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one (1.5 g, 100M%) was dissolved in DMF (20mL) and NaH (95%, 0.13g, 100M%) was added in one portion. The reaction mixture was stirred for 1 hour and then the appropriate alkyl iodide (300M%) was added and stirring was continued for 12 hours. The reaction mixture was poured into water and extracted with ethyl acetate (3x). The ethyl acetate extracts were washed with water (3x) and then brine (1x). Treatment with  $\text{MgSO}_4$ , rotoevaporation, and chromatography (30% EtOAc/hexanes) yielded a BOC-protected 3-amino-1-alkyl-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one in 80% yield.

Step D

The BOC-protected 3-amino-1-alkyl-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one (1.0g, 100M%) was suspended in 30 mL of 1:1  $\text{CH}_2\text{Cl}_2$ /trifluoroacetic acid and the mixture was stirred for 4 hours. The reaction was then rotoevaporated to yield the 3-amino-1-alkyl-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one (100% yield).

-- 90 --

**EXAMPLE 1-A****Synthesis of 3-Amino-1,5-dimethyl-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one****Step A**

5           3-Amino-5-methyl-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one was prepared from 4-methyl- $\alpha$ -tetralone using the methods described in Armstrong et al. *Tetrahedron Letters*, 1994, 35, 3239, incorporated herein by reference.

**Step B**

          3-Amino-5-methyl-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one (9.3g 100M%)  
10       was dissolved in dioxane (300mL) and the solution was chilled to 0°C. BOC-anhydride (13.89g 130M%) was added and the ice bath was removed allowing the solution to come to room temperature and stirring was continued for 16 hours. The solution was rotary evaporated to remove dioxane to provide an off white solid. This solid was recrystallized from CHCl<sub>3</sub> to yield BOC-protected 3-amino-  
15       5-methyl-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one in 55% yield.

**Step C**

          BOC-protected 3-amino-5-methyl-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one (100 M%) was dissolved in DMF (20mL) and NaH (95%, 100 M%) was added in one portion and the reaction mixture was stirred for 1 hour. Methyl iodide (300  
20       M%) was added and this mixture was stirred for 12 hours. The reaction mixture was then poured into water and extracted with ethyl acetate (3x) then backwashed with water (3x) and then brine (1x). Treatment with MgSO<sub>4</sub>, rotoevaporation, and chromatography (5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) yielded BOC-protected 3-amino-1,5-dimethyl-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one in 75% yield.

**Step D**

          BOC-protected 3-amino-1,5-dimethyl-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one (100 M%) was suspended in 30 mL of 1:1 CH<sub>2</sub>Cl<sub>2</sub>/trifluoroacetic acid. The reaction mixture was stirred for 4 hours. The reaction was then rotoevaporated to yield 3-amino-1,5-dimethyl-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one (100%  
30       yield).



-- 91 --

**EXAMPLE 1-B****Synthesis of 5-(L-Alaninyl)-amino-3,3,7-trimethyl-5,7-dihydro-6H-benz[b]azepin-6-one Hydrochloride**

Following the General Procedure H and using N-t-Boc-L-alanine and 5-amino-3,3,7-trimethyl-5,7-dihydro-6H-benz[b]azepin-6-one hydrochloride (Example 1-C), 5-(N-t-Boc-L-alaninyl)amino-3,3,7-trimethyl-5,7-dihydro-6H-benz[b]azepine-6-one was prepared. Following the General Procedure (Example 1-C, Step B) and using 5-(N-t-Boc-L-alaninyl)amino-3,3,7-trimethyl-5,7-dihydro-6H-benz[b]azepin-6-one, 5-(L-alaninyl)amino-3,3,7-trimethyl-5,7-dihydro-6H-benz[b]azepin-6-one hydrochloride was prepared.

**EXAMPLE 1-C****Synthesis of 5-Amino-3,3,7-trimethyl-5,7-dihydro-6H-benz[b]azepin-6-one Hydrochloride**

**GENERAL PROCEDURE**  
**N-Alkylation of Lactams**

**Step A**

To a stirred solution of a BOC-protected  $\alpha$ -aminocaprolactam (6.87 g, 30 mmol) in DMF (150 mL) was added in portions 97% NaH (1.08g, 45 mmol). Bubbling occurred immediately and followed by heavy precipitation. After 10 minutes, benzyl bromide (3.93 mL, 33 mmol) was added. The precipitate dissolved quickly and in about 10 min. a clear solution was obtained. The reaction mixture was stirred overnight and then evaporated as completely as possible on a rotovap at 30°C. Ethyl acetate (100 mL) was added to the residue and this mixture was washed with water, brine, and dried over magnesium sulfate. After filtration and concentration, a thick liquid (10 g) was obtained which was then chromatographed over silica gel with 1:3 ethyl acetate/hexane as the eluant to provide 5.51 g (58%) of the N-benzylated product as an oil. Other lactams and alkylating agents may be used in this procedure to obtain a wide variety of N-alkylated lactams. Various bases, such as LiN(SiMe<sub>3</sub>), may also be employed.

-- 92 --

Following this General Procedure and using N-t-Boc-5-amino-3,3-dimethyl-5,7-dihydro-6H-benz[b]azepin-6-one (General Procedure 1-B, followed by Boc protection) and methyl iodide. N-t-Boc-5-amino-3,3,7-trimethyl-5,7-dihydro-6H-benz[b]azepin-6-one was prepared.

5

## GENERAL PROCEDURE

### BOC Removal Procedure

#### Step A

A stream of anhydrous HCl gas was passed through a stirred solution of the N-t-Boc protected amino acid in 1,4-dioxane (0.03-0.09 M), chilled in a ice bath to ~10°C under N<sub>2</sub>, for 10-15 minutes. The solution was capped, the cooling bath removed, and the solution was allowed to warm to room temperature with stirring for 2-8 hours, monitoring by TLC for the consumption of starting material. The solution was concentrated (and in some instances dissolved in CH<sub>2</sub>Cl<sub>2</sub> then re-concentrated and placed in vacuum oven at 60-70°C to remove most of the residual dioxane) and used without further purification.

Following this General Procedure and using N-t-Boc-5-amino-3,3,7-trimethyl-5,7-dihydro-6H-benz[b]azepin-6-one, the title compound was prepared.

20

## EXAMPLE 1-D

### Synthesis of 3-(S)-Amino-1-methyl-5-oxa-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one

#### Step A

3-(S)-Amino-5-oxa-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one was prepared from N-Boc-serine (Bachem) and 2-fluoro-1-nitrobenzene (Aldrich) using the method of R. J. DeVita et al., *Bioorganic and Medicinal Chemistry Lett.* **1995**, 5(12) 1281-1286, incorporated herein by reference.

#### Step B

Following the General Procedure of Step A of Example 1-C and using the product from Step A of this example, the title compound was prepared.

30

-- 93 --

**EXAMPLE 1-E**Synthesis of 3-(S)-Amino-1-ethyl-5-oxa-1,3,4,5-tetrahydro-2H-1-benzazepin-2-oneStep A

- 5        3-(S)-Amino-5-oxa-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one was prepared from N-Boc-serine (Bachem) and 2-fluoro-1-nitrobenzene (Aldrich) using the method of R. J. DeVita et al., *Bioorganic and Medicinal Chemistry Lett.* 1995, 5(12) 1281-1286, incorporated herein by reference.

Step B

- 10        Following the General Procedure of Step A of Example 1-C and using the product from Step A of this example, the title compound was prepared.

**EXAMPLE 1-F**Synthesis of 3-(S)-Amino-1-methyl-5-thia-1,3,4,5-tetrahydro-2H-1-benzazepin-2-one

15

The title compound was prepared from N-Boc-cystine (Novabio) and 2-fluoro-1-nitrobenzene (Aldrich) using the method of R. J. DeVita et al., *Bioorganic and Medicinal Chemistry Lett.* 1995, 5(12) 1281-1286, incorporated herein by reference, followed by the General Procedure of Step A of Example 1-C.

20

**EXAMPLE 1-G**Synthesis of 7-Amino-1,3,4,7,12,12a-hexahydropyrido[2,1-b][3]-benzazepin-6(2H)-one

25

Step A        Synthesis of N-Chloroacetyl-2-benzylpiperidine

Following General Procedure F and using 2-benzylpyridine, the title compound was prepared.

Physical data were as follows:

(MW = 251.8); mass spectroscopy (MH<sup>+</sup>) 252.0.

-- 94 --

**Step B**      Synthesis of 1,3,4,7,12,12a-hexahydropyrido[2,1-b][3]benzazepin-6(2H)-one

Following General Procedure G and using *N*-chloroacetyl-2-benzylpiperidine, the title compound was prepared.

5      Physical data were as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>): δ = 1.3-1.9 (6H); 2.42 (t, 1H); 3.08 (m, 2H); 3.47 (m, 1H); 3.96 (q, 2H); 4.66 (d, 1H); 7.2 (m, 4H).

(MW = 215.3); mass spectroscopy (MH<sup>+</sup>) 216.1.

10      **Step C**      Synthesis of 7-Oximo-1,3,4,7,12,12a-hexahydropyrido[2,1-b][3]benzazepin-6(2H)-one

Following General Procedure 3A (Step B) and using 1,3,4,7,12,12a-hexahydropyrido[2,1-b][3]benzazepin-6(2H)-one (from Step B above), the title compound was prepared.

Physical data were as follows:

15      (MW = 244.3); mass spectroscopy (MH<sup>+</sup>) 245.0.

**Step D**      Synthesis of 7-Amino-1,3,4,7,12,12a-hexahydropyrido[2,1-b][3]benzazepin-6(2H)-one

Following General Procedure 3A (Step C) and using 7-oximo-1,3,4,7,12,12a-hexahydropyrido[2,1-b][3]benzazepin-6(2H)-one (from Step C above), the title compound was prepared.

20      Physical data were as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>): δ = 1.3-1.9 (6H); 2.42 (t, 1H); 3.08 (m, 2H); 3.47 (m, 1H); 3.96 (q, 2H); 4.66 (d, 1H); 7.2 (m, 4H).

(MW = 230.3); mass spectroscopy (MH<sup>+</sup>) 231.1.

25

**EXAMPLE 1-H**

Synthesis of 1-(N'-L-Alaninyl)amino-4,5,6,7-tetrahydro-3,7-methano-3H-3-benzazonin-2(1H)-one

**Step A**      Synthesis of *N*-Chloroacetyl-3-phenylpiperidine

30

-- 95 --

Following General Procedure F and using 3-phenylpyridine hydrochloride (Aldrich), the title compound was prepared.

Step B      Synthesis of 4,5,6,7-Tetrahydro-3,7-methano-3H-3-benzazonin-2(1H)-one

5      Following General Procedure G and using *N*-chloroacetyl-3-phenylpiperidine, the title compound was prepared.

Physical data were as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>): δ = 1.32-1.57 (2H); 2.08 (m, 2H); 2.81 (t, 1H); 3.13 (bs, 1H); 3.37 (m, 2H); 4.36 (m, 2H); 4.50 (d, 1H).

10      (MW = 201.3); mass spectroscopy (MH<sup>+</sup>) 202.1.

Step C      Synthesis of 1-Oximo-4,5,6,7-tetrahydro-3,7-methano-3H-3-benzazonin-2(1H)-one

Following General Procedure 3A (Step B) and using the product from Step B above, the title compound was prepared.

15      Step D      Synthesis of 1-Amino-4,5,6,7-tetrahydro-3,7-methano-3H-3-benzazonin-2(1H)-one

Following General Procedure 3A (Step C) and using the product from Step C above, the title compound was prepared.

Physical data were as follows:

20      <sup>1</sup>H-nmr (CDCl<sub>3</sub>): δ = 2.86 (t, 1H); 3.17 (bs, 1H); 3.39 (dd, 1H); 4.40 (d, 1H); 4.50 (d, 1H); 5.39 (s, 1H).

(MW = 216.3); mass spectroscopy (MH<sup>+</sup>) 217.4.

Step E      Synthesis of 1-(N'-Boc-L-Alaninyl)amino-4,5,6,7-tetrahydro-3,7-methano-3H-3-benzazonin-2(1H)-one

25      Following General Procedure D and using *N*-tert-Boc-L-alanine (Aldrich) and the product from Step D above, the title compound was prepared.

Physical data were as follows:

(MW = 387.48); mass spectroscopy (MH<sup>+</sup>) 388.1.

-- 96 --

**Step F**      Synthesis of 1-(N'-L-Alaninyl)amino-4,5,6,7-tetrahydro-3,7-methano-3H-3-benzazonin-2(1H)-one

Following General Procedure E and using the product from Step E, the title compound was prepared.

5      Physical data were as follows:

$^1\text{H-nmr}$  ( $\text{CDCl}_3$ ):  $\delta$  = 2.85 (t, 1H); 3.16 (bs, 1H); 3.40 (dd, 1H); 3.67 (m, 1H); 4.35 (d, 1H); 4.56 (d, 1H); 6.40 (d, 1H).

(MW = 287.4); mass spectroscopy (MH+) 288.1.

10

**EXAMPLE 2-A**

Synthesis of 5-Amino-5,7-dihydro-6H-dibenzo[a,c]-cyclohepten-6-ol Hydrochloride

15

**Step A**      Synthesis of 5-Oximo-5,7-dihydro-6H-dibenzo[a,c]cyclohepten-6-one

20

A round bottom flask was charged with 5,7-dihydro-6H-dibenzo[a,c]cyclohepten-6-one (1.0 g, 4.81 mmol)(CAS# 1139-82-8, prepared as described in *Tetrahedron Letters*, Vol. 28, No. 23, (1987), pp 2633-2636, incorporated herein by reference) and butyl nitrite (0.673 mL, 5.77 mmol) (Aldrich) in  $\text{Et}_2\text{O}$ . The solution was cooled to  $0^\circ\text{C}$  and treated drop-wise with a saturated solution of  $\text{HCl(g)}/\text{Et}_2\text{O}$ . After 5 hours at  $0^\circ\text{C}$ , the resulting precipitate was filtered, rinsed with cold  $\text{Et}_2\text{O}$  and vacuum dried to give the title compound as a colorless solid.

25

NMR data was as follows:

$^1\text{H-nmr}$  ( $\text{CDCl}_3$ ):  $\delta$  = 7.26-7.74 (m, 8H), 3.84 (m, 2H).

$\text{C}_{15}\text{H}_{11}\text{NO}_2$  (MW = 237.26); mass spectroscopy (MH+) 238.

Anal. Calcd for  $\text{C}_{15}\text{H}_{11}\text{NO}_2$ ; C, 75.93 H, 4.67 N, 5.90. Found: C, 75.67 H, 4.83 N, 5.67.

30

**Step B**      Synthesis of 5-Amino-5,7-dihydro-6H-dibenzo[a,c]-cyclohepten-6-ol Hydrochloride

-- 97 --

The compound isolated above (0.489 g, 2.04 mmol) was dissolved in THF and added drop-wise to a well-stirred mixture of LAH (10.2 mL, 10.2 mmol)/THF. After heating to reflux for 25 hours under N<sub>2</sub> atmosphere the solution was quenched and worked-up according to Fieser's method. The  
5 resulting solid was rinsed with NH<sub>3</sub> sat/CHCl<sub>3</sub>, the filtrate evaporated and the title compound purified by chromatography (SiO<sub>2</sub>, CHCl<sub>3</sub>).

C<sub>15</sub>H<sub>15</sub>NO (MW = 225.290); mass spectroscopy (MH<sup>+</sup>) 226.

Anal. Calcd for C<sub>15</sub>H<sub>15</sub>NO; C, 79.97 H, 6.71 N, 6.22. Found: C, 80.19 H, 6.71 N, 5.91.

10

#### EXAMPLE 2-B

##### Synthesis of 5-[L-alaninyl]-amino-5,7-dihydro-6H-dibenzo[a,c]- cyclohepten-6-one

Following General Procedure D above using Boc-L-alanine (Aldrich) and  
15 5-amino-5,7-dihydro-6H-dibenzo[a,c]cyclohepten-6-ol hydrochloride (Example 2-A), the compound was prepared as a tan foam.

The resulting alcohol was oxidized as follows. To a stirred mixture of oxalyl chloride (0.15 mL, 1.2 mmol) in 10 mL of dichloromethane cooled to -78°C was added DMSO (0.106 mL, 1.5 mmol) and the mixture was stirred for  
20 10 minutes. A solution of the alcohol (0.1828 g, 0.60 mmol) in 20 mL of chloroform was added dropwise. The reaction mixture was stirred at -78°C for 2 hours, and then 0.5 mL (3.6 mmol) of triethylamine was added. Stirring was continued for 1 hour and then the mixture was allowed to warm to room temperature and stirring was continued at ambient temperature overnight. The  
25 mixture was then diluted with 50 mL of dichloromethane, washed with brine (3x), dried over magnesium sulfate, filtered and evaporated to dryness to give the crude product which was typically purified by column chromatography.

The Boc group was removed using 2.0 M HCl/dioxane. The title compound was isolated as an orange foam.

-- 98 --

$C_{18}H_{18}N_2O_2HCl$  (MW = 330.4); mass spectroscopy (MH<sup>+</sup> of freebase)  
295.

C. Dibenzazepinone Derivatives and Related Compounds

5

**GENERAL PROCEDURE 3-A**

Preparation of 5-Amino-7-alkyl-5,7-dihydro-  
6H-dibenz[b,d]azepin-6-one Derivatives

Step A

10        Following the General Procedure of Step A of Example 1-C and using 5,7-dihydro-6H-dibenz[b,d]azepin-6-one and an alkyl halide, the 7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one was prepared.

Step B

15        The 7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (1 eq.) was dissolved in THF and isoamylnitrite (1.2 eq.) was added. The mixture was cooled to 0°C in an ice bath. NaHMDS (1.1 eq., 1M in THF) was added dropwise. After stirring for 1 hour or until the reaction was complete, the mixture was concentrated then acidified with 1N HCl and extracted with EtOAc. The organic portion was dried and concentrated to yield a crude product which was purified by silica gel  
20        chromatography.

Step C

25        The resulting oxime was dissolved in EtOH/NH<sub>3</sub> (20:1) and hydrogenated in a bomb using Raney nickel and hydrogen (500 psi) at 100°C for 10 hours. The resulting mixture was filtered and concentrated to provide an oil which was purified by silica gel chromatography to yield the title compound.



-- 99 --

**GENERAL PROCEDURE 3-B****Preparation of Fluoro-substituted 5,7-dihydro-6H-dibenz[b,d]azepin-6-one Derivatives**

A modification of the procedure of Robin D. Clark and Jahangir, *Tetrahedron*, Vol. 49, No. 7, pp. 1351-1356, 1993<sup>15</sup> was used. Specifically, an appropriately substituted N-t-Boc-2-amino-2'-methylbiphenyl was dissolved in THF and cooled to -78°C. s-Butyl lithium (1.3M in cyclohexane, 2.2 eq.) was added slowly so that the temperature remained below -65°C. The resulting mixture was allowed to warm to -25°C and was stirred at that temperature for 1 hour. The mixture was cooled to -78°C. Dry CO<sub>2</sub> was bubbled through the mixture for 30 seconds. The mixture was allowed to warm to ambient temperature then was carefully quenched with water. The mixture was concentrated under reduced pressure then was adjusted to pH 3 with 1N HCl. The mixture was extracted with EtOAc and the organic portion was dried and concentrated to yield a crude material. The crude material was dissolved in methanol and the solution was saturated with HCl. The mixture was heated at reflux for 12 hours then was allowed to cool. The mixture was concentrated to provide crude lactam which was purified by chromatography or crystallization.

20

**GENERAL PROCEDURE 3-C****Resolution of 5-Amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one**

In a round bottom flask was added the racemic freebase amine (1.0 eq.) in methanol followed by di-*p*-toluoyl-D-tartaric acid monohydrate (1.0 eq.). The mixture was concentrated *in vacuo* to a residue and redissolved in a moderate volume of methanol and allowed to stir at room temperature open to the atmosphere (8-72 hours). The solid was removed by filtration. The enantiomeric excess was determined by chiral HPLC (Chiracel ODR) using 15% acetonitrile and 85% H<sub>2</sub>O with 0.1% trifluoroacetic acid and a flow rate of 1.0 mL/minute at 35°C. The resolved di-*p*-toluoyl-D-tartaric salt was then dissolved in EtOAc and saturated NaHCO<sub>3</sub> until pH 9-10 was reached. The layers were separated and the

30

-- 100 --

organic layer was washed again with saturated  $\text{NaHCO}_3$ ,  $\text{H}_2\text{O}$ , and brine. The organic layer was dried over  $\text{MgSO}_4$  and the drying agent was removed by filtration. The filtrate was concentrated *in vacuo*. The free amine was dissolved in MeOH and HCl (12M, 1.0 eq.) was added. The salt was concentrated *in vacuo* and the resulting film was triturated with EtOAc. The HCl salt was filtered and rinsed with EtOAc. The ee was determined by chiral HPLC.

### EXAMPLE 3-A

#### Synthesis of 5-Amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

##### Step A      Synthesis of 7-Methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

A round bottom flask was charged with sodium hydride (0.295 g, 7.46 mmol) in 9.0 mL of DMF and treated with 5,7-dihydro-6H-dibenz[b,d]azepin-6-one (1.3 g, 6.22 mmol) (CAS # 20011-90-9, prepared as described in Brown, et al., Tetrahedron Letters, No. 8, 667-670, (1971) and references cited therein, which are incorporated herein by reference). After stirring at 60°C for 1 hour, the solution was treated with methyl iodide (1.16 mL, 18.6 mmol) and stirring continued for 17 hours with the exclusion of light. After cooling, the reaction was diluted with  $\text{CH}_2\text{Cl}_2/\text{H}_2\text{O}$ , washed with  $\text{NaHSO}_4$  solution,  $\text{H}_2\text{O}$ , and dried over  $\text{Na}_2\text{SO}_4$ . Evaporation and flash chromatography ( $\text{SiO}_2$ ,  $\text{CHCl}_3$ ) gave 0.885 g (63%) of the title compound as a colorless solid.

NMR data was as follows:

$^1\text{H}$ -nmr ( $\text{CDCl}_3$ ):  $\delta$  = 7.62 (d, 2H), 7.26-7.47 (m, 6H), 3.51 (m, 2H), 3.32 (s, 3H).

$\text{C}_{15}\text{H}_{13}\text{NO}$  (MW = 223.27); mass spectroscopy ( $\text{MH}^+$ ) 223.

Anal. Calcd for  $\text{C}_{15}\text{H}_{13}\text{NO}$ ; C, 80.69 H, 5.87 N, 6.27. Found: C, 80.11 H, 5.95 N, 6.23.

-- 101 --

Step B      Synthesis of 7-Methyl-5-oximo-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

The compound isolated above (0.700 g, 3.14 mmol) was dissolved in 20 mL of toluene and treated with butyl nitrite (0.733 mL, 6.28 mmol). The reaction temperature was lowered to 0°C and the solution was treated with KHMDS (9.42 mL, 0.5 M) under N<sub>2</sub> atmosphere. After stirring for 1 hour the reaction was quenched with a saturated solution of NaHSO<sub>4</sub>, diluted with CH<sub>2</sub>Cl<sub>2</sub> and separated. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and the title compound purified by chromatography (SiO<sub>2</sub>, 98:2 CHCl<sub>3</sub>/MeOH) giving 0.59 g (80 %) as a colorless solid.

C<sub>15</sub>H<sub>12</sub>N<sub>2</sub>O<sub>2</sub> (MW = 252.275); mass spectroscopy (MH<sup>+</sup>) 252.

Anal. Calcd for C<sub>15</sub>H<sub>12</sub>N<sub>2</sub>O<sub>2</sub>; C, 71.42 H, 4.79 N, 11.10. Found: C, 71.24 H, 4.69 N, 10.87.

Step C      Synthesis of 5-Amino-7-Methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

The oxime isolated above (0.99 g, 3.92 mmol) was hydrogenated in a Parr apparatus at 35 psi over 10 % Pd/C (0.46 g) in 3A ethanol. After 32 hours, the reaction mixture was filtered through a plug of celite, the filtrate evaporated to a foam and treated with a saturated solution of HCl (g) in Et<sub>2</sub>O. The resulting colorless solid was filtered, rinsed with cold Et<sub>2</sub>O and vacuum dried to give 0.66 g (61 %) of the title compound.

NMR data was as follows:

<sup>1</sup>H-nmr (DMSOd<sub>6</sub>): δ = 9.11 (bs, 3H), 7.78-7.41(m, 8H), 4.83 (s, 1H), 3.25 (s, 3H).

C<sub>15</sub>H<sub>14</sub>N<sub>2</sub>O · HCl (MW = 274.753); mass spectroscopy (MH<sup>+</sup> free base) 238.  
Anal. Calcd for C<sub>15</sub>H<sub>14</sub>N<sub>2</sub>O · HCl; C, 65.57 H, 5.50 N, 10.19 Found: C, 65.27 H, 5.67 N, 10.13.

-- 102 --

**EXAMPLE 3-B**Synthesis of (S)- and (R)-5-(L-Alaninyl)-amino-7-methyl-  
5,7-dihydro-6H-dibenz[b,d]azepin-6-oneStep A      Synthesis of (S)- and (R)-5-(N-Boc-L-Alaninyl)-amino-7-methyl-  
5,7-dihydro-6H-dibenz[b,d]azepin-6-one

5                    Boc-L-Alanine (0.429 g, 2.26 mmol) (Aldrich) was dissolved in THF and  
treated with HOBt (0.305 g, 2.26 mmol), and 5-amino-7-methyl-5,7-dihydro-6H-  
dibenz[b,d]azepin-6-one (0.45 g, 1.89 mmol) (Example 3-A). The temperature  
was lowered to 0°C and the reaction mixture treated with EDC (0.449 g, 2.26  
10 mmol) (Aldrich) and stirred 17 hours under N<sub>2</sub>. The reaction mixture was  
evaporated, the residue diluted with EtOAc/H<sub>2</sub>O, washed 1.0 N HCl, sat.  
NaHCO<sub>3</sub>, brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The diastereomers were separated on a  
Chiralcel OD column using 10% IPA/heptane at 1.5 mL/minute.

Isomer 1: Retention time 3.37 minutes.

15                    NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>): δ = 7.62-7.33 (m, 9H), 5.26 (d, 1H), 5.08 (m, 1H),  
4.34 (m, 1H), 3.35 (s, 3H), 1.49 (s, 9H), 1.40 (d, 3H).

Optical Rotation: [α]<sub>20</sub> = - 96 @ 589 nm (c = 1, MeOH).

C<sub>23</sub>H<sub>27</sub>N<sub>3</sub>O<sub>4</sub> (MW = 409.489); mass spectroscopy (MH<sup>+</sup>) 409.

20                    Anal. Calcd for C<sub>23</sub>H<sub>27</sub>N<sub>3</sub>O<sub>4</sub>; C, 67.46 H, 6.64 N, 10.26. Found: C,  
68.42 H, 7.02 N, 9.81.

Isomer 2: Retention time 6.08 minutes.

NMR data was as follows:

25                    <sup>1</sup>H-nmr (CDCl<sub>3</sub>): δ = 7.74 (bd, 1H), 7.62-7.32 (m, 8H), 5.28 (d, 1H),  
4.99 (m, 1H), 4.36 (m, 1H), 3.35 (s, 3H), 1.49 (s, 9H), 1.46 (d, 3H).

Optical Rotation: [α]<sub>20</sub> = 69 @ 589 nm (c = 1, MeOH).

C<sub>23</sub>H<sub>27</sub>N<sub>3</sub>O<sub>4</sub> (MW = 409.489); mass spectroscopy (MH<sup>+</sup>) 409.

Anal. Calcd for C<sub>23</sub>H<sub>27</sub>N<sub>3</sub>O<sub>4</sub>; C, 67.46 H, 6.64 N, 10.26. Found: C,  
67.40 H, 6.62 N, 10.02

30

-- 103 --

**Step B**      Synthesis of (S)- and (R)-5-(L-Alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

The compounds isolated in Step A (each isomer separately) were dissolved in dioxane and treated with excess HCl (g). After stirring for 17 hours, the title  
5 compounds were isolated as colorless solids after evaporation and vacuum drying.

Isomer 1:

$C_{18}H_{19}N_3O_2 \cdot HCl$  (MW = 345.832); mass spectroscopy (MH<sup>+</sup> free base) 309.  
Optical Rotation:  $[\alpha]_{20} = -55$  @ 589 nm (c = 1, MeOH).

Isomer 2:

10  $C_{18}H_{19}N_3O_2 \cdot HCl$  (MW = 345.832); mass spectroscopy (MH<sup>+</sup> free base) 309.  
Optical Rotation:  $[\alpha]_{20} = 80$  @ 589 nm (c = 1, MeOH).

**EXAMPLE 3-C**

15      Synthesis of (S)- and (R)-5-(L-Valinyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

**Step A**      Synthesis of (S)- and (R)-5-(N-Boc-L-Valinyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

Boc-L-Valine (0.656 g, 3.02 mmol) (Aldrich) was dissolved in THF and treated with HOBt (0.408, 3.02 mmol), DIPEA (1.05 mL, 6.05 mmol) and 5-  
20 amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one hydrochloride (0.75 g, 2.75 mmol) (Example 3-A). The temperature was lowered to 0°C and the reaction mixture treated with EDC (0.601 g, 3.02 mmol) (Aldrich) and stirred 17 hours under N<sub>2</sub>. The reaction mixture was evaporated, the residue diluted with EtOAc/H<sub>2</sub>O, washed 1.0 N HCl, sat. NaHCO<sub>3</sub>, brine and dried over Na<sub>2</sub>SO<sub>4</sub>.  
25 The diastereomers were separated on a Chiralcel OD column using 10% IPA/heptane at 1.5 mL/minute.

Isomer 1: Retention time 3.23 minutes.

Optical Rotation:  $[\alpha]_{20} = -120$  @ 589 nm (c = 1, MeOH).

$C_{25}H_{31}N_3O_4$  (MW = 437.544); mass spectroscopy (MH<sup>+</sup>) 438

30 Isomer 2: Retention time 6.64 minutes.

-- 104 --

Optical Rotation:  $[\alpha]_{20} = 50 @ 589 \text{ nm}$  ( $c = 1$ , MeOH).

$\text{C}_{25}\text{H}_{31}\text{N}_3\text{O}_4$  (MW = 437.544); mass spectroscopy (MH+) 438

**Step B**      Synthesis of (S)- and (R)-5-(L-Valinyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

5            The compounds isolated in Step A (each isomer separately) were dissolved in dioxane and treated with excess HCl (g). After stirring for 17 hours, the title compounds were isolated as colorless solids after evaporation and vacuum drying.

Isomer 1:

$\text{C}_{20}\text{H}_{23}\text{N}_3\text{O}_2 \cdot \text{HCl}$  (MW = 373.88); mass spectroscopy (MH+ free base) 338.

10          Optical Rotation:  $[\alpha]_{20} = -38 @ 589 \text{ nm}$  ( $c = 1$ , MeOH).

Isomer 2:

$\text{C}_{20}\text{H}_{23}\text{N}_3\text{O}_2 \cdot \text{HCl}$  (MW = 373.88); mass spectroscopy (MH+ free base) 338.

Optical Rotation:  $[\alpha]_{20} = 97 @ 589 \text{ nm}$  ( $c = 1$ , MeOH).

15

**EXAMPLE 3-D**

Synthesis of (S)- and (R)-5-(L-tert-Leucine)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

**Step A**      Synthesis of (S)- and (R)-5-(N-Boc-L-tert-Leucinyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

20            Boc-L-tert-Leucine (0.698 g, 3.02 mmol) (Fluka) was dissolved in THF and treated with HOBt (0.408, 3.02 mmol), DIPEA (1.05 mL, 6.05 mmol) and 5-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one hydrochloride (0.75 g, 2.75 mmol) (Example 3-A). The temperature was lowered to 0°C and the reaction mixture treated with EDC (0.601 g, 3.02 mmol) (Alrich) and stirred 17  
25          hours under  $\text{N}_2$ . The reaction mixture was evaporated, the residue diluted with EtOAc/ $\text{H}_2\text{O}$ , washed 1.0 N HCl, sat.  $\text{NaHCO}_3$ , brine and dried over  $\text{Na}_2\text{SO}_4$ . The diastereomers were separated on a Chiralcel OD column using 10% IPA/heptane at 1.5 mL/minute.

Isomer 1: Retention time 3.28minutes.

30            Optical Rotation:  $[\alpha]_{20} = -128 @ 589 \text{ nm}$  ( $c = 1$ , MeOH).

-- 105 --

$C_{26}H_{33}N_3O_4$  (MW = 451.571); mass spectroscopy (MH+) 452

Isomer 2: Retention time 5.52 minutes.

Optical Rotation:  $[\alpha]_{20} = 26 @ 589 \text{ nm}$  (c = 1, MeOH).

$C_{26}H_{33}N_3O_4$  (MW = 451.571); mass spectroscopy (MH+) 452

5    Step B            Synthesis of (S)- and (R)-5-(L-tert-LeucinyI)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

The compounds isolated in Step A (each isomer separately) were dissolved in dioxane and treated with excess HCl (g). After stirring for 17 hours, the title compounds were isolated as colorless solids after evaporation and vacuum drying.

10   Isomer 1:

$C_{21}H_{25}N_3O_2 \cdot HCl$  (MW = 387.91); mass spectroscopy (MH+ free base) 352.

Optical Rotation:  $[\alpha]_{20} = -34 @ 589 \text{ nm}$  (c = 1, MeOH).

Isomer 2:

$C_{21}H_{25}N_3O_2 \cdot HCl$  (MW = 387.91); mass spectroscopy (MH+ free base) 352.

15   Optical Rotation:  $[\alpha]_{20} = 108 @ 589 \text{ nm}$  (c = 1, MeOH).

### EXAMPLE 3-E

Synthesis of 5-(N-Boc-Amino)-5,7-dihydro-6H,7H-dibenz[b,d]azepin-6-one

Step A            Synthesis of 5-Iodo-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

20        A solution of 5,7-dihydro-6H-dibenz[b,d]azepin-6-one (1.0 g, 4.77 mmol) (Example 3-A) and  $Et_3N$  (2.66 mL, 19.12 mmol) was stirred for 5.0 minutes at  $-15^\circ C$  in  $CH_2Cl_2$  and treated with TMSI (1.36 mL, 9.54 mmol). After stirring for 15 minutes  $I_2$  (1.81 g, 7.16 mmol) was added in a single portion and the reaction allowed to warm to  $5-10^\circ C$  over 3 h. The reaction was quenched with

25   sat.  $Na_2SO_3$ , diluted with  $CH_2Cl_2$  and separated. The organics were washed with  $Na_2SO_3$  and  $NaHSO_3$  and dried over  $MgSO_4$ . After filtration, the organics were concentrated to approximately 20 mL and diluted with an additional 20 mL of hexanes. The title compound was isolated as a tan precipitate by filtration.

Step B            Synthesis of 5-Azido-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

-- 106 --

The iodide isolated above was dissolved in DMF and treated with 1.2 equivalents of  $\text{NaN}_3$ . After stirring 17 hour at  $23^\circ\text{C}$ , the mixture was diluted with  $\text{EtOAc}/\text{H}_2\text{O}$ , separated, washed with brine and dried over  $\text{MgSO}_4$ . The title compound was triturated from hot  $\text{EtOAc}$  as a tan powder.

5    Step C            Synthesis of 5-(N-Boc-Amino)-5,7-dihydro-6H,7H-dibenz[b,d]azepin-6-one

The azide was dissolved in  $\text{THF}/\text{H}_2\text{O}$  and stirred at  $23^\circ\text{C}$  for 17 hours in the presence of 3.0 equivalents of  $\text{Ph}_3\text{P}$ . The reaction was diluted with 50 %  $\text{HOAc}/\text{toluene}$ , separated, the aqueous layer extracted with toluene and evaporated to an oily residue. This was taken to pH 7.0 by the addition of 1 N  $\text{NaOH}$ , the resulting  $\text{HOAc}$  salt was collected and vacuum dried. Finally, the compound was treated with Boc anhydride (1.05 equivalents) and  $\text{Et}_3\text{N}$  (2.1 equivalents) in  $\text{THF}$ . After stirring for 5 hours at  $23^\circ\text{C}$ , the reaction was filtered and the title compound isolated as a colorless powder.

15

**EXAMPLE 3-F**

Synthesis of 5-Amino-7-(2-methylpropyl)-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

20    Step A            Synthesis of 5-(N-Boc-Amino)-7-(2-methylpropyl)-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

A solution of 5-(N-Boc-amino)-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (0.2g, 0.617 mmol) (Example 3-E) in DMF was treated with  $\text{Cs}_2\text{CO}_3$  (0.22 g, 0.678 mmol) and warmed to  $60^\circ\text{C}$ . To the reaction mixture was added 1-iodo-2-methylpropane (0.078 mL, 0.678 mmol) and stirring continued for 17 hours. After cooling to  $23^\circ\text{C}$  the mixture was diluted with  $\text{CH}_2\text{Cl}_2$ , washed with several portions of brine and dried over  $\text{Na}_2\text{SO}_4$ . The title compound was purified by chromatography ( $\text{SiO}_2$ ,  $\text{CHCl}_3/\text{MeOH}$  9:1).

25                     $\text{C}_{23}\text{H}_{28}\text{N}_2\text{O}_3$  (MW = 380.41); mass spectroscopy ( $\text{MH}^+$ ) 381

                    Anal. Calcd for  $\text{C}_{23}\text{H}_{28}\text{N}_2\text{O}_3$ ; C, 72.61 H, 7.42 N, 7.36. Found: C, 72.31 H, 7.64 N, 7.17.

30



-- 107 --

**Step B**      Synthesis of 5-Amino-7-(2-methylpropyl)-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

The compound isolated in Step A was deprotected in dioxane saturated with gaseous HCl. The title compound was isolated as a slightly colored solid after evaporation and vacuum drying.

**EXAMPLE 3-G**

Synthesis of 5-Amino-7-(methoxyacetyl)-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

**Step A**      Synthesis of 5-(N-Boc-Amino)-7-(methoxyacetyl)-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

A solution of 5-(N-Boc-amino)-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (1.03, 3.08 mmol) (Example 3-E) in DMF was treated with Cs<sub>2</sub>CO<sub>3</sub> (1.10 g, 3.39 mmol) and warmed to 60°C. To the reaction mixture was added bromomethyl acetate (0.321 mL, 3.39 mmol) (Aldrich) and stirring continued for 17 hours. After cooling to 23°C, the mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed with several portions of brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The title compound was purified by chromatography (SiO<sub>2</sub>, CHCl<sub>3</sub>).

C<sub>22</sub>H<sub>24</sub>N<sub>2</sub>O<sub>5</sub> (MW = 396.44): mass spectroscopy (MH<sup>+</sup>) 397

Anal. Calcd for C<sub>22</sub>H<sub>24</sub>N<sub>2</sub>O<sub>5</sub>; C, 66.65 H, 6.10 N, 7.07. Found: C, 66.28 H, 5.72 N, 6.50.

**Step B**      Synthesis of 5-Amino-7-(methoxyacetyl)-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

The compound isolated in Step A was deprotected in dioxane saturated with gaseous HCl. The title compound was isolated as a colorless solid after evaporation and vacuum drying.

C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>3</sub> HCl (MW = 332.78); mass spectroscopy (MH<sup>+</sup> free base) 297.

-- 108 --

**EXAMPLE 3-H**Synthesis of 5-Amino-7-(3,3-dimethyl-2-butanonyl)-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

5      Step A      Synthesis of 5-(N-Boc-Amino)-7-(3,3-dimethyl-butanonyl)-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

A solution of 5-(N-Boc-amino)-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (0.2 g, 0.617 mmol) (Example 3-E) in DMF was treated with  $\text{Cs}_2\text{CO}_3$  (0.3 g, 0.925 mmol) and warmed to 60°C. To the reaction mixture was added 1-chloro-3,3-dimethyl-2-butanone (0.096 mL, 0.74 mmol) (Aldrich) and stirring continued for 17 hours. After cooling to 23 °C, the mixture was diluted with  $\text{CH}_2\text{Cl}_2$ , washed with several portions of brine and dried over  $\text{Na}_2\text{SO}_4$ . The title compound was isolated as a colorless solid.

$\text{C}_{25}\text{H}_{30}\text{N}_2\text{O}_4$  (MW = 422.522); mass spectroscopy (MH<sup>+</sup>) 423

15      Step B      Synthesis of 5-Amino-7-(3,3-dimethyl-2-butanonyl)-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

The compound isolated in Step A was deprotected in dioxane saturated with gaseous HCl. The title compound was isolated as a colorless solid after evaporation and vacuum drying.

20      **EXAMPLE 3-I**

Synthesis of L-Alaninyl-5-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one HydrochlorideStep A

Following General Procedure D and using N-t-Boc-L-alanine and 5-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one, N-t-Boc-L-alaninyl-5-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one was prepared.

Step B

Following the General Procedure of Step B of Example 1-C and using the N-t-Boc-L-alaninyl-5-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one, the title compound was prepared. Other substituted N-t-Boc-L-alaninyl-5-amino-

-- 109 --

7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-ones can also be prepared by this procedure.

### EXAMPLE 3-J

5                    Synthesis of L-Valinyl-5-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

Step A

Following General Procedure D and using N-t-Boc-L-valine and 5-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one, N-t-Boc-L-valinyl-5-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one was prepared.

Step B

Following the General Procedure of Step B of Example 1-C and using the N-t-Boc-L-valinyl-5-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one, the title compound was prepared. Other substituted N-t-Boc-L-valinyl-5-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-ones can also be prepared by this procedure.

### EXAMPLE 3-K

20                    Synthesis of 5-Amino-7-phenbutyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

Following General Procedure 3-A and using 5,7-dihydro-6H-dibenz[b,d]azepin-6-one (prepared as described in Brown, et. al., Tetrahedron Letters, No. 8, 667-670, (1971) and references cited therein, which are incorporated herein by reference) and 1-chloro-4-phenylbutane (Aldrich), the title compound was prepared.

25

### EXAMPLE 3-L

Synthesis of 5-Amino-7-cyclopropylmethyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

Following General Procedure 3-A and using 5,7-dihydro-6H-dibenz[b,d]azepin-6-one (prepared as described in Brown, et. al., Tetrahedron

30

-- 110 --

Letters, No. 8, 667-670, (1971) and references cited therein, which are incorporated herein by reference) and (bromomethyl)cyclopropane (Aldrich), the title compound was prepared.

5

**EXAMPLE 3-M**

Synthesis of 5-Amino-7-(2',2',2'-trifluoroethyl)-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

Following General Procedure 3-A and using 5,7-dihydro-6H-dibenz[b,d]azepin-6-one (prepared as described in Brown, et. al., Tetrahedron Letters, No. 8, 667-670, (1971) and references cited therein, which are incorporated herein by reference) and 1-bromo-2,2,2-trifluoroethane (Aldrich), the title compound was prepared.

15

**EXAMPLE 3-N**

Synthesis of 5-Amino-7-cyclohexyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

Following General Procedure 3-A and using 5,7-dihydro-6H-dibenz[b,d]azepin-6-one (prepared as described in Brown, et. al., Tetrahedron Letters, No. 8, pp. 667-670, (1971) and references cited therein, which are incorporated herein by reference) and bromocyclohexane (Aldrich), the title compound was prepared.

25

**EXAMPLE 3-O**

Synthesis of 5-(L-Alaninyl)amino-9-fluoro-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

**Step A**

2-Bromo-5-fluorotoluene was stirred in THF at -78C. s-BuLi (1.05 eq., 1.3 M in cyclohexane) was slowly added and the mixture was stirred for 45 minutes. Trimethylborate (1.5 eq.) was added and the mixture was allowed to warm to ambient temperature. After stirring for 1 hour, pinacol (2 eq.) was

-- 111 --

added. The mixture was stirred for 16 hours then was concentrated under reduced pressure. The resulting residue was slurried in  $\text{CH}_2\text{Cl}_2$  and filtered through Celite. The filtrate was concentrated to yield an oil which was purified by chromatography on deactivated silica gel ( $\text{Et}_3\text{N}$ ) to yield the arylboronate ester.

5    Step B

2-Bromoaniline (1 eq.) and di-t-butyl-dicarbonate (1.1 eq.) were stirred at  $80^\circ\text{C}$  for 20 hours. The resulting mixture was allowed to cool and was directly distilled using house vacuum to provide N-t-Boc-2-bromoaniline.

Step C

10       N-t-Boc-2-bromoaniline (Step 2, 1 eq.), the arylboronate ester (Step 1, 1.1 eq.),  $\text{K}_2\text{CO}_3$  (1.1 eq.) and tetrakis(triphenylphosphine)palladium(0) (0.02 eq.) were stirred in 20% water/dioxane under nitrogen. The solution was heated at reflux for 10 hours. The mixture was allowed to cool then was concentrated. The  
15       resulting residue was partitioned between water and chloroform. The organic portion was dried and concentrated to yield an oil which was purified by silica gel chromatography using 1:1  $\text{CH}_2\text{Cl}_2$ /hexanes.

Step D

Following General Procedure 3-B and using the substituted biphenyl from step 3, the 9-fluoro-5,7-dihydro-6H-dibenz[b,d]azepin-6-one was prepared.

20    Step E

9-Fluoro-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (1 eq., Step 4), cesium carbonate (1.1 eq., Aldrich) and methyl iodide (1.1 eq., Aldrich) were stirred in dry DMF at ambient temperature for 16 hours. The mixture was concentrated under reduced pressure to provide a residue which was partitioned between EtOAc  
25       and water. The organic portion was dried and concentrated to yield an oil which was purified by silica gel chromatography to 9-fluoro-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one.

Step F

-- 112 --

Following General Procedure 3-A, Step B and 9-fluoro-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one from Step 5, 5-amino-9-fluoro-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one was prepared.

**Step G**

- 5        Following the procedure of Example 3-I and using 5-amino-9-fluoro-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one from Step 6, the title compound was prepared.

**EXAMPLE 3-P**

- 10        Synthesis of 5-(L-Alaninyl)amino-13-fluoro-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

Following the procedure of Example 3-O and using 2-bromo-4-fluoroaniline (Step 2, Lancaster) and o-tolylboronic acid (Step 3, Aldrich), the title compound was prepared.

15

**EXAMPLE 3-Q**

Synthesis of 5-(L-Alaninyl)amino-10-fluoro-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

- Following the procedure of Example 3-O and using 2-bromo-4-fluorotoluene (Step 1), the title compound was prepared.

20

**EXAMPLE 3-R**

Synthesis of 5-(L-Alanyl)-amino-7-cyclopropylmethyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

- 25        Following the procedure of Example 3-I and using 5-amino-7-cyclopropylmethyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (Example 3-L), the title compound was prepared.

**EXAMPLE 3-S**

- 30        Synthesis of 5-(L-Alaninyl)amino-7-phenbutyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

-- 113 --

Following the procedure of Example 3-I and using 5-amino-7-phenbutyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (Example 3-K), the title compound was prepared.

5

**EXAMPLE 3-T**

Synthesis of 5-(L-Valinyl)amino-7-cyclopropylmethyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

Following the procedure of Example 3-J and using 5-amino-7-cyclopropylmethyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (Example 3-L), the  
10 title compound was prepared.

**EXAMPLE 3-U**

Synthesis of 5-(L-Valinyl)amino-7-phenbutyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

15 Following the procedure of Example 3-J and using 5-amino-7-phenbutyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (Example 3-K), the title compound was prepared.

**EXAMPLE 3-V**

20 Synthesis of 5-(L-Valinyl)amino-7-hexyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

Step A

Following General Procedure 3-A and using 5,7-dihydro-6H-dibenz[b,d]azepin-6-one (prepared as described in Brown, et. al., Tetrahedron  
25 Letters, No. 8, 667-670, (1971) and references cited therein, which are incorporated herein by reference) and 1-bromohexane (Aldrich), 5-amino-7-hexyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one was prepared.

Step B

Following the procedure of Example 3-J and using 5-amino-7-hexyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one, the title compound was prepared.  
30

-- 114 --

**EXAMPLE 3-W**Synthesis of 5-(L-Valinyl)amino-10-fluoro-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

Following the procedure of Example 3-J and using 5-amino-10-fluoro-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (as prepared in Example 3-Q), the title compound was prepared.

**EXAMPLE 3-X**

10 Synthesis of 5-(L-Valinyl)amino-13-fluoro-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

Following the procedure of Example 3-J and using the 5-amino-13-fluoro-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (as prepared in Example 3-P), the title compound was prepared.

15

**EXAMPLE 3-Y**Synthesis of 5-(L-Valinyl)amino-13-fluoro-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

Following the procedure of Example 3-J and using the 5-amino-9-fluoro-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (as prepared in Example 3-O), the title compound was prepared.

20

**EXAMPLE 3-Z**Synthesis of (5-Amino-7-methyl-1,2,3,4,5,7-hexahydro-6H-dicyclohexyl[b,d]azepin-6-one

25 The 5-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one hydrochloride (Example 3-A) was dissolved in a 1:1 mixture of EtOAc/HOAc. 5% Rh/C was added and the mixture was stirred at 60°C under 60 psi of hydrogen. After 3 days, the mixture was filtered and the filtrate was concentrated to provide an oil which was purified by SCX-cation exchange chromatography to yield the title compound.

30



-- 115 --

**EXAMPLE 3-AA**Synthesis of 5-(S)-Amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

Following General Procedure 3-C using racemic 5-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (1.0 eq.) and di-*p*-toluoyl-D-tartaric acid monohydrate (1.0 eq.) in methanol, the title compound was prepared as a solid. The product was collected by filtration. Enantiomeric excess was determined by chiral HPLC.

Desired enantiomer 1: retention time of 9.97 minutes.

10 Undesired enantiomer 2: retention time of 8.62 minutes.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>): δ = 9.39 (s, 2H), 7.75-7.42 (m, 8H), 4.80 (s, 1H), 3.30 (s, 3H).

C<sub>15</sub>H<sub>15</sub>ClN<sub>2</sub>O (MW = 274.75); mass spectroscopy (MH<sup>+</sup>) 239.1.

15 Anal Calcd for C<sub>15</sub>H<sub>15</sub>ClN<sub>2</sub>O<sub>3</sub>; C, 65.57; H, 5.50; N, 10.20; Found: C, 65.51, H, 5.61; N, 10.01.

**EXAMPLE 3-AB**Synthesis of 9-Amino-5,6-dihydro-4H-quinol[8,1-ab][3]benzazepin-8(9H)-one Hydrochloride

20

Step A      Synthesis of 8-Phenylquinoline

A degassed solution of 8-bromoquinoline (1.0 g, 4.81 mmol) (Aldrich) in dioxane (50 mL)/H<sub>2</sub>O (10 mL) was treated with phenylboronic acid (0.64 g, 5.29 mmol) (Aldrich), Pd(Ph<sub>3</sub>P)<sub>4</sub> (0.050 g, 0.04 mmol) and K<sub>2</sub>CO<sub>3</sub> (0.73 g, 5.29 mmol). After refluxing for 4 hours under a N<sub>2</sub> atmosphere the reaction was allowed to cool, diluted with EtOAc and separated. After drying over Na<sub>2</sub>SO<sub>4</sub> and SiO<sub>2</sub> chromatography (95:5 Hexanes/EtOAc) the titled compound was isolated as a colorless oil.

Physical data were as follows:

30 <sup>1</sup>H-nmr (CDCl<sub>3</sub>): δ = 8.97 (d, 1H), 8.22 (dd, 1H), 7.87-7.39 (m, 9H).

-- 116 --

$C_{15}H_{11}N$  (MW = 205); mass spectroscopy (MH+ ) 206.

**Step B**      Synthesis of 8-Phenyl-1,2,3,4-tetrahydroquinoline

The product from Step A (0.99 g, 4.82 mmol) was hydrogenated according to the procedure described by Honel, M., et. al., J.C.S. Perkin I, (1980), 1933-

5    1938, incorporated herein by reference.

Physical data were as follows:

$^1H$ -nmr ( $CDCl_3$ ):  $\delta$  = 7.46 (m, 3H), 7.38 (m, 2H), 6.98 (m, 2H), 6.70 (m, 1H), 3.27 (t, 2H), 2.86 (t, 2H), 1.96 (m, 2H).

$C_{15}H_{15}N$  (MW = 209); mass spectroscopy (MH+ ) 210.

10    **Step C**      Synthesis of 1-Chloromethylacetyl-8-phenyl-1,2,3,4-tetrahydroquinoline

The product from Step B (1.0 g, 4.78 mmol) was dissolved in  $CH_2Cl_2$  (20 mL)/  $H_2O$  (20 mL) and treated with  $NaHCO_3$  (0.602 g, 7.18 mmol) followed by chloroacetyl chloride (0.478 mL, 5.26 mmol). After stirring for 17 h at 23°C, the reaction was diluted with  $CH_2Cl_2$ , washed with saturated  $NaHCO_3$ , dried over  $Na_2SO_4$  and purified by  $SiO_2$  chromatography ( $CHCl_3$ /Hexanes 9:1). The product was isolated as a colorless solid.

Physical data were as follows:

$C_{17}H_{16}ClNO$  (MW = 286.77); mass spectroscopy (MH+ ) 287.

20    Anal. Calcd for  $C_{17}H_{16}ClNO$ ; C, 71.45 H, 5.64 N, 4.90. Found: C, 71.63 H, 5.60 N, 4.87.

**Step D**      Synthesis of 5,6-Dihydro-4H-quinolo[8,1-ab][3]benzazepin-8(9H)-one

The product from Step C (0.89 g, 3.11 mmol) was mixed thoroughly with  $AlCl_3$  (0.87 g, 6.54 mmol) at 23°C and the mixture heated neat at 100°C for 5-7 minutes. After vigorous gas evolution, the molten mixture was allowed to cool and extracted with several portions of  $CH_2Cl_2$ / $NaHCO_3$  (sat). The combined organic layers were dried over  $Na_2SO_4$  and the title compound was purified by chromatography ( $SiO_2$ ,  $CHCl_3$ /hexanes 9:1), yielding a colorless oil which solidified upon standing.

30      Physical data were as follows:

-- 117 --

$C_{17}H_{15}NO$  (MW = 249.312); mass spectroscopy (MH+ ) 250.

Anal. Calcd for  $C_{17}H_{15}NO$ ; C, 81.90 H, 6.06 N, 5.62. Found: C, 81.75 H, 6.11 N, 5.86.

**Step E**      Synthesis of 9-Oximo-5,6-Dihydro-4H-quinolizin-8(9H)-one

5

The product from Step D (0.490 g, 1.97 mmol) was dissolved in THF and butyl nitrite (0.46 mL, 3.93 mmol) and treated with KHMDS (0.5 M, 4.52 mL, 2.26 mmol) at 0°C. After stirring for 1 h, the reaction was quenched with cold 1 N HCl, extracted with EtOAc, the combined organic layers dried over  $Na_2SO_4$  and the product purified by  $SiO_2$  chromatography ( $CHCl_3/MeOH$ , 99:1). The title compound was isolated as a colorless solid.

Physical data were as follows:

$C_{17}H_{14}N_2O_2$  (MW = 278.3); mass spectroscopy (MH+ ) 279.

Anal. Calcd for  $C_{17}H_{14}N_2O_2$  .0.3317 mol  $H_2O$ ; C, 71.82 H, 5.19 N, 9.85. Found: C, 71.85 H, 5.09 N, 9.59.

15

**Step F**      Synthesis of 9-Amino-5,6-Dihydro-4H-quinolizin-8(9H)-one

The product from Step E (0.360 g, 1.29 mmol) was hydrogenated over Ra/Ni (0.05 g) in EtOH (50 mL)/  $NH_3$  (anhydrous) (5.0 mL) at 100°C and 500 psi for 10 h. The catalyst was removed by filtration and the resulting filtrate chromatographed over  $SiO_2$  ( $CHCl_3/MeOH$ , 98:2) yielding the titled compound as a colorless oil which solidified upon standing.

20

Physical data were as follows:

$C_{17}H_{16}N_2O$  (MW = 264.326); mass spectroscopy (MH+ ) 266.

Anal. Calcd for  $C_{17}H_{16}N_2O$ ; C, 77.25 H, 6.10 N, 10.60. Found: C, 77.23 H, 6.15 N, 10.49.

25

-- 118 --

**EXAMPLE 3-AC**Synthesis of 9-(N'-L-Alaninyl)amino-5,6-dihydro-4H-quinol[8,1-ab][3]benzazepin-8(9H)-one Hydrochloride5      Step A      Synthesis of 9-(N'-Boc-L-Alaninyl)amino-5,6-Dihydro-4H-quinol[8,1-ab][3]benzazepin-8(9H)-one

Following General Procedure D and using N-Boc-Alanine (Aldrich) and 9-amino-5,6-dihydro-4H-quinol[8,1-ab][3]benzazepin-8(9H)-one (from Example 3-AB), the title compound was prepared.

Physical data were as follows:

10       $C_{25}H_{29}N_3O_4$  (MW = 435.521); mass spectroscopy (MH<sup>+</sup>) 436.

Anal. Calcd for  $C_{25}H_{29}N_3O_4 \cdot 0.4102$  mol H<sub>2</sub>O; C, 67.79 H, 6.79 N, 9.49; Found: C, 67.83 H, 6.91 N, 9.29.

15      Step B      Synthesis of 9-(N'-L-Alaninyl)amino-5,6-dihydro-4H-quinol[8,1-ab][3]benzazepin-8(9H)-one Hydrochloride

Following General Procedure E and using the product from Step A, the title compound was prepared.

Physical data were as follows:

$C_{20}H_{21}N_3O_2 \cdot HCl$  (MW = 371.6); mass spectroscopy (MH<sup>+</sup> free base) 335.

20      **EXAMPLE 3-AD**Synthesis of 5-[L-alaninyl]-amino-7-(2-methylpropyl)-5,7-dihydro-6H-dibenz[b,d]azepin-6-one hydrochloride

Following General Procedure D above using Boc-L-alanine (Aldrich) and 5-amino-7-(2-methylpropyl)-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

25      hydrochloride (Example 3-F), the compound was prepared as a tan foam. The resulting Boc group was removed using 2.0 M HCl/dioxane. The title compound was isolated as a slightly colored solid after evaporation and vacuum drying.

$C_{21}H_{24}N_3O_2 \cdot HCl$  (MW = 386); mass spectroscopy (MH<sup>+</sup> of freebase) 351.

-- 119 --

**EXAMPLE 3-AE****Synthesis of 5-[L-alaninyl]-amino-5,7-dihydro-6H,7H-dibenz[b,d]azepin-6-one hydrochloride****5      Step A      Synthesis of 5-Amino-5,7-dihydro-6H,7H-dibenz[b,d]azepin-6-one hydrochloride**

5-(N-Boc-Amino)-5,7-dihydro-6H,7H-dibenz[b,d]azepin-6-one (Example 3-E) was treated with 2.0 M HCl/dioxane. After stirring for 17 h at 23°C, the title compound was isolated as a slightly colored solid after filtration and vacuum drying.

10       $C_{14}H_{12}N_2OHCl$  (MW = 260.72); mass spectroscopy (MH<sup>+</sup> of freebase) 225.

Anal. Calcd for  $C_{14}H_{12}N_2OHCl$ : C, 64.50 H, 5.03 N, 10.74. Found: C, 64.35 H, 4.99 N, 10.51.

**15      Step B      Synthesis of 5-[N-Boc-L-alaninyl]-amino-5,7-dihydro-6H,7H-dibenz[b,d]azepin-6-one**

The compound isolated above was coupled with Boc-L-alanine (Aldrich) following General Procedure D. The title compound was used without further purification.

20       $C_{22}H_{25}N_4O_4$  (MW = 395.45); mass spectroscopy (MH<sup>+</sup>) 396.  
Anal. Calcd for  $C_{22}H_{25}N_4O_4$ : C, 66.82 H, 6.37 N, 10.63. Found: C, 65.53 H, 6.16 N, 10.38.

**Step C      Synthesis of 5-[L-alaninyl]-amino-5,7-dihydro-6H,7H-dibenz[b,d]azepin-6-one hydrochloride**

25      The compound isolated above was deprotected using HCl/dioxane. The title compound was used without further purification after stirring for 17 h at 23 C and vacuum drying.

**D.      Benzodiazepine Derivatives and Related Compounds**

30

-- 120 --

**GENERAL PROCEDURE 4-A****N-1-Methylation of Benzodiazepines**

A solution of benzodiazepine (1 eq.) in DMF (0.1 M concentration) at 0°C was treated with potassium tert-butoxide (1.0 eq., 1.0 M solution in THF). After  
5 stirring for 30 minutes at 0°C, iodomethane (1.3 eq.) was added and stirring continued for 25 minutes. The mixture was diluted with methylene chloride and washed with water and brine. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The crude product was then either purified by trituration with  
1:1 ether/hexanes or chromatographed via HPLC using ethyl acetate/hexanes as  
10 the eluent.

**GENERAL PROCEDURE 4-B****Cbz Removal Procedure**

A flask was charged with the Cbz-protected 3-aminobenzodiazepine (1  
15 eq.). To this was added HBr (34 eq.; 30% solution in acetic acid). Within 20 minutes all of the starting material dissolves. The reaction was stirred for 5 hours at ambient temperature. Ether was added to the orange solution causing the HBr•amine salt to precipitate. The mixture was decanted. This process of adding ether and decanting was repeated thrice in an effort to remove acetic acid and  
20 benzyl bromide. Toluene was added and the mixture concentrated *in vacuo*. This step was also repeated. The HBr salt was partitioned between ethyl acetate and 1 M K<sub>2</sub>CO<sub>3</sub>. The aqueous layer was back-extracted with ethyl acetate. The combined organics were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated.

25

**GENERAL PROCEDURE 4-C****Boc Removal Procedure**

A solution of Boc-protected amine (1 eq.) in methylene chloride (0.15 M concentration) was cooled to 0°C and treated with trifluoroacetic acid (30 eq.).

-- 121 --

After 10 minutes at 0°C, the cooling bath was removed and stirring continued at ambient for 20 minutes to 1 hour. The mixture was concentrated *in vacuo* to remove excess trifluoroacetic acid. The residue was dissolved in methylene chloride and washed with saturated aqueous NaHCO<sub>3</sub> or 1 M K<sub>2</sub>CO<sub>3</sub> and brine.

- 5 The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated.

#### GENERAL PROCEDURE 4-D

##### Azide Transfer Reaction Using KHMDs

- 10 The azido derivative was prepared using the procedure described in John W. Butcher et al., *Tet. Lett.*, 37, 6685-6688 (1996), incorporated herein by reference.

#### GENERAL PROCEDURE 4-E

##### Azide Transfer Reaction Using LDA

- 15 To a solution of diisopropylamine (1.1 eq.) in 1 mL of dry THF cooled to -78°C was added n-butyl lithium (1.6M in hexane) (1.1 eq.) dropwise maintaining the reaction temperature at -78°C. The reaction mixture was stirred for 30 minutes at -78°C and then the lactam (0.471 mM) was added dropwise as a solution in 1 mL of dry THF. The reaction mixture was stirred at -78°C for 30  
20 minutes and then a pre-cooled solution of trisyl azide (1.2 eq.) was added as a solution in 1 mL of dry THF. The reaction mixture was stirred at -78°C for 20 minutes and then quenched with acetic acid (4.0 eq.). The reaction mixture was then stirred at 40°C for 2 hours. The reaction was then poured into EtOAc and washed with water, sodium bicarbonate and brine, and then dried over sodium  
25 sulfate, filtered and concentrated. The residue was purified by LC 2000 chromatography.

#### GENERAL PROCEDURE 4-F

##### Azido Group Reduction

-- 122 --

The azido group was reduced to the corresponding primary amine using the procedure described in John W. Butcher et al., *Tet. Lett.*, **37**, 6685-6688 (1996), incorporated herein by reference.

5

#### GENERAL PROCEDURE 4-G

##### N-Alkylation of Amides or Lactams Using Sodium Hydride or Potassium tert-Butoxide

To a slurry of sodium hydride or potassium tert-butoxide (1.1 eq) in 15 mL of dry DMF was added the appropriate amide (0.0042 moles) as a solution in 10 mL of DMF. The alkyl iodide was then added and a thick slurry resulted. The reaction became less thick as time elapsed and when complete by TLC the reaction had become homogeneous. The reaction mixture was poured over ice and extracted into ethyl acetate. The organic layer was washed with water, followed by brine. The organic layer was then dried over sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by HPLC (LC 2000), eluting with an ethyl acetate/hexane system.

15

#### GENERAL PROCEDURE 4-H

##### N-Alkylation of Amides or Lactams Using KHMDS

20

To the appropriate amide or lactam in THF cooled to -78°C was added KHMDS dropwise and the reaction mixture was stirred for 30 min. at -78°C. The alkyl iodide was then added dropwise while maintaining the temperature at -70°C. The cooling bath was then removed and reaction was allowed to warm to room temperature and stirring was continued for 2 hours. The reaction mixture was then poured over ice and extracted into ethyl acetate. The organic extracts were washed with water, followed by brine. The organic layer was then dried over sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by HPLC (LC 2000), eluting with an ethyl acetate/hexane system.

25

30



-- 123 --

**GENERAL PROCEDURE 4-I****N-Alkylation of Amides or Lactams Using Cesium Carbonate**

To a solution of the amide or lactam in DMF was added cesium carbonate (1.05 eq) and an alkyl iodide (1.1 eq). The mixture was allowed to stir overnight  
5 at room temperature and then the reaction mixture was diluted with ethyl acetate and washed with water, followed by brine. The organic layer was dried over sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by HPLC (LC 2000), eluting with an ethyl acetate/hexane system.

**10 GENERAL PROCEDURE 4-J****BOC Removal Procedure**

To an N-Boc protected compound was added  $\text{CH}_2\text{Cl}_2/\text{TFA}$  (4:1) at room temperature. The reaction mixture was stirred at room temperature for 3 hours and then concentrated. The residue was extracted into dichloromethane and  
15 washed with water, saturated sodium bicarbonate, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to give the free amine.

**GENERAL PROCEDURE 4-K****Azide Transfer Procedure**

20 This azide transfer procedure is a modification of the procedure described in Evans, D. A. et al. *J. Am. Chem. Soc.* **1990**, *112*, 4011-4030, incorporated herein by reference. To a solution of the lactam substrate (1.0 eq.) in THF (~0.1 M) under  $\text{N}_2$  at  $-78^\circ\text{C}$  was added a solution of  $\text{KN}(\text{TMS})_2$  (1.1 eq. of 0.5 M in Toluene, Aldrich) dropwise over a period of 2-10 minutes. A slight exotherm was  
25 often observed by an internal thermometer, and the resulting solution was stirred for 5-15 minutes, while re-cooling to  $-78^\circ\text{C}$ . Then, trisyl azide (1.1-1.5 eq., CAS No. 36982-84-0, prepared as described by references in the Evans reference above, which are incorporated herein by reference) in THF (~0.5 M), either precooled to  $-78^\circ\text{C}$  or at room temperature, was added via cannula over a period

-- 124 --

of 0.5-5 minutes. Again, a slight exotherm was generally noted. The resulting solution was stirred for from 5-10 minutes, while re-cooling to  $-78^{\circ}\text{C}$ . Then, AcOH (4.5-4.6 eq., glacial) was added, the cooling bath removed and the mixture allowed to warm to room temperature with stirring for 12-16 hours. The mixture  
5 was diluted with EtOAc, in a 2-5 volume multiple of the initial THF volume, and washed with dilute aq.  $\text{NaHCO}_3$  (1-2x), 0.1-1.0 M aq. HCl (0-2x), and brine (1x). The organic phase was then dried over  $\text{MgSO}_4$ , filtered, and concentrated to provide the crude product.

#### 10 GENERAL PROCEDURE 4-L

##### Azide Reduction to an Amine

A mixture of the azide in absolute EtOH (0.03-0.07 M) and 10% Pd/C (~ 1/3 by weight of the azide) was shaken in a Parr apparatus under  $\text{H}_2$  (35-45 psi) at room temperature for 3-6 hours. The catalyst was removed by filtration  
15 through a plug of Celite, rinsing with absolute EtOH, and the filtrate was concentrated to provide the crude amine product.

#### GENERAL PROCEDURE 4-M

##### Amide Alkylation Using Cesium Carbonate

20 This procedure is a modification of the procedure described in Claremon, D. A.; et al, PCT Application: WO 96/406555, incorporated herein by reference. To a mixture of 2,4-dioxo-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine (CAS No. 49799-48-6) in DMF (1.0 eq., 0.7 M) under  $\text{N}_2$  at room temperature was added  $\text{Cs}_2\text{CO}_3$  (2.2 eq.) and the appropriate alkyl halide (2.2 eq.). The mixture was  
25 stirred at room temperature for 5.5-16 hours. The mixture was partitioned between EtOAc and sat.  $\text{NaHCO}_3$ . The aqueous layer was extracted with EtOAc (1-2x) and the combined EtOAc extracts were dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated to provide the crude product.

-- 125 --

**GENERAL PROCEDURE 4-N**BOC Removal Procedure

A stream of anhydrous HCl gas was passed through a stirred solution of the N-t-Boc protected amino acid in 1,4-dioxane (0.03-0.09 M), chilled in a ice bath to ~10°C under N<sub>2</sub>, for 10-15 minutes. The solution was capped, the cooling bath removed, and the solution was allowed to warm to room temperature with stirring for 2-8 hours, monitoring by TLC for the consumption of starting material. The solution was concentrated (and in some instances dissolved in CH<sub>2</sub>Cl<sub>2</sub> then re-concentrated and placed in vacuum oven at 60-70°C to remove most of the residual dioxane) and used without further purification.

**EXAMPLE 4-A**Synthesis of 3-Amino-1,3-dihydro-5-(1-piperidiny)-2H-1,4-benzodiazepin-2-one

15 Step A      Preparation of 1,2-Dihydro-3H-1-methyl-5-(1-piperidiny)-1,4-benzodiazepin-2-one

A solution of phosphorous pentachloride (1.2 eq) in methylene chloride was added dropwise to a solution of 1-methyl-1,2,3,4-tetrahydro-3H-1,4-benzodiazepin-2,5-dione (Showell, G. A. et al. *J. Med. Chem.* **1994**, *37*, 719, incorporated herein by reference) in methylene chloride. The resultant yellowish-orange solution was stirred at ambient temperature for 2.5 hours; the solvent was removed *in vacuo*. The orange residue was redissolved in methylene chloride, cooled to 0 °C, and treated with a solution of piperidine (2 eq) and triethylamine (2 eq) in methylene chloride. The cooling bath was removed and the reaction stirred for 18 hours. The reaction mixture was washed with saturated aqueous NaHCO<sub>3</sub> (back-extracted with methylene chloride) and brine. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The residue was purified via HPLC eluting with a gradient of 4 to 10% methanol/methylene chloride affording the title intermediate as a yellow solid having a melting point of 103-105°C.

30      C<sub>15</sub>H<sub>19</sub>N<sub>3</sub>O (MW 257.37); mass spectroscopy 257.

-- 126 --

Anal. Calcd for  $C_{15}H_{19}N_3O$ : C, 70.01; H, 7.44; N, 16.33. Found: C, 69.94; H, 7.58; N, 16.23.

**Step B**      Preparation of 1,2-Dihydro-3H-1-methyl-3-oximido-5-(1-piperidinyl)-1,4-benzodiazepin-2-one

5            Potassium tert-butoxide (2.5 eq) was added in two portions to a  $-20^{\circ}\text{C}$  solution of 1,2-dihydro-3H-1-methyl-5-(1-piperidinyl)-1,4-benzodiazepin-2-one (1 eq.) in toluene. After stirring at  $-20^{\circ}\text{C}$  for 20 minutes, isoamyl nitrite (1.2 eq.; Aldrich) was added to the red reaction mixture. The reaction was stirred at  $-20^{\circ}\text{C}$  for 5 hours at which time the reaction was done by TLC. The cooling bath  
10 was removed and the reaction quenched with 0.5 M citric acid. After stirring for 10 minutes, diethyl ether was added. The suspension was stirred at ambient temperature overnight then filtered washing with ether. The resultant cream colored solid had a melting point of  $197-200^{\circ}\text{C}$ .

$^1\text{H}$  NMR data of the E/Z isomers was as follows:

15             $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  = 7.64 (1H, bs), 7.48 (2H, d,  $J=7.4$  Hz), 7.35-7.20 (6H, m), 6.75 (1H, bs), 3.8-3.2 (8H, m), 3.46 (3H, s), 3.42 (3H, s), 1.90-1.40 (12H, m).

$C_{15}H_{18}N_4O_2$  (MW = 286.37); mass spectroscopy 286.

20            **Step C**      Preparation of 1,2-dihydro-3H-1-methyl-3-[O-(ethylaminocarbonyl)-oximido]-5-(1-piperidinyl)-1,4-benzodiazepin-2-one

A mixture of 1,2-dihydro-3H-1-methyl-3-oximido-5-(1-piperidinyl)-1,4-benzodiazepin-2-one (1 eq.) in THF was treated with ethyl isocyanate (1.7 eq) and triethylamine (0.6 eq). The mixture was heated to  $64^{\circ}\text{C}$  for 4 hours. The mixture was concentrated and the residue purified by HPLC eluting with 5%  
25 methanol/methylene chloride.

$^1\text{H}$  NMR data of the E/Z isomers was as follows:

30             $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  = 7.50 (2H, dd,  $J=8.4, 1.5$  Hz), 7.35-7.22 (6H, m), 6.42 (1H, bt), 6.20 (1H, bt), 3.7-3.4 (8H, m), 3.46 (3H, s), 3.44 (3H, s), 3.25 (4H, m), 1.9-1.4 (12H, m), 1.12 (3H, t,  $J=6.3$  Hz), 1.10 (3H, t,  $J=6.3$  Hz).

-- 127 --

$C_{18}H_{23}N_5O_3$  (MW = 357.46); mass spectroscopy 357.

**Step D**      Preparation of 3-Amino-1,3-dihydro-2H-1-methyl-5-(1-piperidinyl)-1,4-benzodiazepin-2-one

5      The 1,2-dihydro-3H-1-methyl-3-[O-(ethylaminocarbonyl)oximido]-5-(1-piperidinyl)-1,4-benzodiazepin-2-one (1 eq.) was hydrogenated in methanol over 5% palladium on carbon (0.15 eq.) at 43 psi for 3.25 hours. The reaction was filtered through celite and concentrated *in vacuo*. The residue was taken up in methylene chloride and filtered a second time through celite. The filtrate was concentrated and the resultant foam was used immediately.

10

**EXAMPLE 4-B**

Synthesis of 3-(L-Alaninyl)-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one

15      **Step A**      Preparation of (S)-3-amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one, (1S)-7,7-dimethyl-2-oxobicyclo[2.2.1]heptane-1-methanesulfonate

20      The title intermediate was prepared according to Reider, P. J. et al. *J. Org. Chem.* **1987**, 52, 955, incorporated herein by reference, using 3-amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one (Bock M. G. et al.; *J. Org. Chem.* **1987**, 52, 3232, incorporated herein by reference) as the starting material.

**Step B**      Preparation of 3-[N'-(tert-Butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one

25      (S)-3-Amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one, (1S)-7,7-dimethyl-2-oxobicyclo[2.2.1]heptane-1-methanesulfonate was free based by partitioning between methylene chloride and 1M potassium carbonate. The free amine was then coupled with N-Boc-alanine following General Procedure D.

$C_{24}H_{28}N_4O_4$  (MW = 436.56); mass spectroscopy 436.

30      Anal. Calc. for  $C_{24}H_{28}N_4O_4$ : C, 66.03; H, 6.47; N, 12.84. Found: C, 65.79; H, 6.68; N, 12.80.

-- 128 --

**Step C**      Preparation of 3-(L-Alaninyl)-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one

Following General Procedure 4-C using 3-[N'-(tert-butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one, the  
5 title compound was prepared as a white foam.

Anal. Calc. for  $C_{19}H_{19}N_4O_2$ : C, 69.21; H, 6.64; N, 15.37. Found: C, 70.11; H, 6.85; N, 15.01.

**EXAMPLE 4-C**

10      Synthesis of 3-(L-Alaninyl)-amino-7-chloro-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one

**Step A**      Preparation of 3-(Benzyloxycarbonyl)-amino-7-chloro-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one

A solution of 3-(benzyloxycarbonyl)-amino-7-chloro-2,3-dihydro-5-phenyl-1H-1,4-Benzodiazepin-2-one (1 eq; Neosystem) in DMF was cooled to 0°C and  
15 treated with potassium *tert*-butoxide (1 eq; 1.0M solution in THF). The resultant yellow solution was stirred at 0°C for 30 minutes then quenched with methyl iodide (1.3 eq.). After stirring an additional 25 minutes the reaction was diluted with methylene chloride and washed with water and brine. The organic phase was  
20 dried over  $Na_2SO_4$ , filtered, and concentrated. The residue was purified via HPLC chromatography eluting with a gradient of 20-30% ethyl acetate/hexanes.

$C_{24}H_{20}ClN_3O_3$  (MW = 433.92); mass spectroscopy 433.

Anal. calcd for  $C_{24}H_{20}ClN_3O_3$ : C, 66.44; H, 4.65; N, 9.68. Found: C, 66.16; H, 4.50; N, 9.46.

25      **Step B**      Preparation of 3-Amino-7-chloro-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one

Following General Procedure 4-B using 3-(benzyloxycarbonyl)-amino-7-chloro-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a white foam which was used immediately in Step C.

-- 129 --

**Step C**      Preparation of 3-[N'-tert-Butylcarbamate)-L-alaninyl]-amino-7-chloro-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one

Following General Procedure D using N-Boc-L-alanine and 3-amino-7-chloro-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a white foam.

$C_{24}H_{28}ClN_4O_4$  (MW = 471.18); mass spectroscopy 471

Anal. calcd for  $C_{24}H_{28}ClN_4O_4$ : C, 61.21; H, 5.78; N, 11.90. Found: C, 61.24; H, 5.59; N, 11.67.

**Step D**      Preparation of 3-(L-Alaninyl)amino-7-chloro-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one

Following General Procedure 4-C using 3-[N'-tert-butylcarbamate)-L-alaninyl]-amino-7-chloro-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a white foam. The crude material was used immediately.

#### EXAMPLE 4-D

Synthesis of 3-(L-Alaninyl)amino-7-bromo-2,3-dihydro-1-methyl-5-(2-fluorophenyl)-1H-1,4-benzodiazepin-2-one

**Step A**      Preparation of 3-(Benzyloxycarbonyl)-amino-7-bromo-2,3-dihydro-1-methyl-5-(2-fluorophenyl)-1H-1,4-benzodiazepin-2-one

Following General Procedure 4-A using 3-(benzyloxycarbonyl)-amino-7-bromo-2,3-dihydro-5-(2-fluorophenyl)-1H-1,4-benzodiazepin-2-one (Neosystem), the title intermediate was prepared as a white foam.

$C_{24}H_{19}BrFN_3O_3$  (MW = 496.36); mass spectroscopy 497.

Anal. calcd for  $C_{24}H_{19}BrFN_3O_3$ : C, 58.08; H, 3.86; N, 8.47. Found: C, 57.90; H, 4.15; N, 8.20.

**Step B**      Preparation of 3-Amino-7-bromo-1,3-dihydro-1-methyl-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-2-one

Following General Procedure 4-B using 3-(benzyloxycarbonyl)-amino-7-bromo-2,3-dihydro-1-methyl-5-(2-fluorophenyl)-1H-1,4-benzodiazepin-2-one, the

-- 130 --

title intermediate was prepared as a white foam which was used immediately in Step C.

**Step C**      Preparation of 3-[N'-(*tert*-Butylcarbamate)-L-alaninyl]-amino-7-bromo-1,3-dihydro-1-methyl-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-2-one

Following General Procedure D using N-Boc-L-alanine (Novo) and 3-amino-7-bromo-1,3-dihydro-1-methyl-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a white foam.

$C_{24}H_{26}BrFN_4O_4$  (MW = 533.12); mass spectroscopy 533.2.

Anal. calcd for  $C_{24}H_{26}BrFN_4O_4$ : C, 54.04; H, 4.91; N, 10.50. Found: C, 53.75; H, 4.92; N, 10.41.

**Step D**      Preparation of 3-(L-Alaninyl)-amino-7-bromo-1,3-dihydro-1-methyl-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-2-one

Following General Procedure 4-C using 3-[N'-(*tert*-butylcarbamate)-L-alaninyl]-amino-7-bromo-1,3-dihydro-1-methyl-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a white foam. The crude material was used immediately.

#### EXAMPLE 4-E

Synthesis of 3-(N'-Methyl-L-alaninyl)-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one

**Step A**      Preparation of 3-[N'-(*tert*-Butylcarbamate)-N'-methyl-L-alaninyl]-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one

Following General Procedure D and using (S)-3-amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one (Example 4-B) and N-*tert*-Boc-N-methyl-alanine (Sigma), the title intermediate was obtained as a white solid.

$C_{25}H_{30}N_4O_4$  (MW = 450.2); mass spectroscopy (M+1) 451.2.

Anal. calcd for  $C_{25}H_{30}N_4O_4$ : C, 66.65; H, 6.71; N, 12.44. Found: C, 66.66; H, 6.89; N, 12.21.

**Step B**      Preparation of 3-(N'-Methyl-L-alaninyl)-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one



-- 131 --

Following General Procedure 4-C and using 3-[N'-(*tert*-butylcarbamate)-N'-methyl-L-alaninyl]-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a white foam.

$C_{20}H_{22}N_4O_2$  (MW = 350.46); mass spectroscopy (M+1) 351.4.

5        Anal. calcd for  $C_{20}H_{22}N_4O_2$ : C, 68.55; H, 6.33; N, 15.99. Found, C, 68.36; H, 6.20; N, 15.79.

#### EXAMPLE 4-F

10        Synthesis of 3-(L-Alaninyl)amino-7-chloro-2,3-dihydro-1-methyl-5-(2-chlorophenyl)-1H-1,4-benzodiazepin-2-one

Step A        Preparation of 3-(Benzyloxycarbonyl)-amino-7-chloro-2,3-dihydro-1-methyl-5-(2-chlorophenyl)-1H-1,4-benzodiazepin-2-one

15        Following General Procedure 4-A using 3-(benzyloxycarbonyl)-amino-7-chloro-2,3-dihydro-5-(2-chlorophenyl)-1H-1,4-benzodiazepin-2-one (Neosystem), the title intermediate was prepared as a white solid having a melting point of 232-233°C.

$C_{24}H_{19}Cl_2N_3O_3$  (MW = 468.36); mass spectroscopy 468.

20         $^1H$  NMR (300 MHz,  $CDCl_3$ ):  $\delta$  = 7.67 (1H, m), 7.52 (1H, dd, J=2.4, 8.7 Hz), 7.42-7.26 (9H, m), 7.07 (1H, d, J=2.4 Hz), 6.70 (1H, d, J=8.3 Hz), 5.35 (1H, d, J=8.4 Hz), 5.14 (2H, ABq, J=19.6 Hz), 3.47 (3H, s).

$^{13}C$  NMR (75 MHz,  $CDCl_3$ ):  $\delta$  = 166.66, 165.65, 155.72, 140.52, 136.99, 136.0, 132.87, 131.99, 131.47, 131.40, 131.38, 131.16, 130.54, 130.06, 128.45, 128.08, 128.03, 127.72, 127.22, 123.28, 122.01, 68.95, 67.02, 35.32.

25        Step B        Preparation of 3-Amino-7-chloro-1,3-dihydro-1-methyl-5-(2-chlorophenyl)-2H-1,4-benzodiazepin-2-one

Following General Procedure 4-B using 3-(benzyloxycarbonyl)-amino-7-chloro-2,3-dihydro-1-methyl-5-(2-chlorophenyl)-1H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a white foam which was used immediately in Step C.

-- 132 --

Step C      Preparation of 3-[N'-(*tert*-Butylcarbamate)-L-alaninyl]-amino-7-chloro-1,3-dihydro-1-methyl-5-(2-chlorophenyl)-2H-1,4-benzodiazepin-2-one

Following General Procedure D using N-Boc-L-alanine and 3-amino-7-chloro-1,3-dihydro-1-methyl-5-(2-chlorophenyl)-2H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a white foam.

$C_{24}H_{26}Cl_2N_4O_4$  (MW = 505.44); mass spectroscopy 505.2.

Step D      Preparation of 3-(L-Alaninyl)-amino-7-chloro-1,3-dihydro-1-methyl-5-(2-chlorophenyl)-2H-1,4-benzodiazepin-2-one

10

Following General Procedure 4-C using 3-[N'-(*tert*-butylcarbamate)-L-alaninyl]-amino-7-chloro-1,3-dihydro-1-methyl-5-(2-chlorophenyl)-2H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a white foam. The crude material was used immediately.

15

#### EXAMPLE 4-G

Synthesis of 3-(L-Alaninyl)amino-5-cyclohexyl-2,3-dihydro-1-methyl-1H-1,4-Benzodiazepin-2-one

Step A      Preparation of 3-(Benzyloxycarbonyl)-amino-5-cyclohexyl-2,3-dihydro-1-methyl-1H-1,4-benzodiazepin-2-one

20

Following General Procedure 4-A using 3-(benzyloxycarbonyl)-amino-5-cyclohexyl-2,3-dihydro-1H-1,4-benzodiazepin-2-one (Neosystem), the title intermediate was prepared as a white solid having a melting point of 205-206°C.

$C_{24}H_{27}N_3O_3$  (MW = 405.54); mass spectroscopy 405.

$^1H$  NMR (300 MHz,  $CDCl_3$ ):  $\delta$  = 7.54 (1H, d, J=7.9 Hz), 7.48 (1H, d, J=7.7 Hz), 7.36-7.26 (7H, m), 6.54 (1H, d, J= 8.3 Hz), 5.15 (1H, d, J=8.0 Hz), 5.09 (2H, ABq, J=17.1 Hz), 3.39 (3H, s), 2.77 (1H, m), 2.01 (1H, bd, J=13.6 Hz), 1.85 (1H, bd, J=12.4 Hz), 1.68-1.49 (4H, m), 1.34-1.02 (4H, m).

25

-- 133 --

**Step B**      Preparation of 3-Amino-5-cyclohexyl-1,3-dihydro-1-methyl-2H-1,4-benzodiazepin-2-one

Following General Procedure 4-B using 3-(benzyloxycarbonyl)-amino-5-cyclohexyl-2,3-dihydro-1-methyl-1H-1,4-benzodiazepin-2-one, the title

5 intermediate was prepared as a white foam which was used immediately in Step C.

$C_{16}H_{21}N_3O$  (MW + H = 272.1763); mass spectroscopy 272.1766

**Step C**      Preparation of 3-[N'-(*tert*-Butylcarbamate)-L-alaninyl]-amino-5-cyclohexyl-1,3-dihydro-1-methyl-2H-1,4-benzodiazepin-2-one

Following General Procedure D using N-Boc-L-alanine and 3-amino-5-cyclohexyl-1,3-dihydro-1-methyl-2H-1,4-benzodiazepin-2-one, the title  
10 intermediate was prepared as a white foam.

$C_{24}H_{34}N_4O_4$  (MW = 442.62); mass spectroscopy (M+H) 443.2.

**Step D**      Preparation of 3-(L-Alaninyl)amino-5-cyclohexyl-1,3-dihydro-1-methyl-2H-1,4-benzodiazepin-2-one

Following General Procedure 4-C using 3-[N'-(*tert*-butylcarbamate)-L-alaninyl]-amino-5-cyclohexyl-1,3-dihydro-1-methyl-2H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a white foam. The crude material was used  
15 immediately.

$C_{19}H_{26}N_4O_2$  (M+H = 343.2136); mass spectroscopy found 343.2139.

20

#### EXAMPLE 4-H

Synthesis of 3-(L-Alaninyl)amino-2,3-dihydro-1-methyl-7-nitro-5-phenyl-1H-1,4-benzodiazepin-2-one

**Step A**      Preparation of 2-[N-( $\alpha$ -Isopropylthio)-N'-(benzyloxycarbonyl)-glycinyl]-amino-5-nitrobenzophenone

25

A solution of  $\alpha$ -(isopropylthio)-N-(benzyloxycarbonyl)glycine (1 eq; prepared according to Zoller, V. and D. Ben-Ishai *Tetrahedron* **1975**, *31*, 863, incorporated herein by reference) in dry THF was cooled to 0 °C and treated with oxalyl chloride (1 eq.) and 3 drops of DMF. After stirring for 15 minutes at 0°C,  
30 the cooling bath was removed and stirring continued at ambient temperature for 40 minutes. The solution was recooled to 0°C. A solution of 2-amino-5-

-- 134 --

nitrobenzophenone (0.9 eq.; Acros) and 4-methylmorpholine (2.0 eq.) in dry THF was added via cannulation to the acid chloride. The cooling bath was removed and the reaction stirred at ambient temperature for 5 hours. The reaction was diluted with methylene chloride and washed with 0.5 M citric acid, saturated aqueous NaHCO<sub>3</sub>, and brine. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The residue was purified via preparative LC2000 eluting with a gradient of 15-20% ethyl acetate/hexanes giving an off-white foam.

C<sub>26</sub>H<sub>25</sub>N<sub>3</sub>O<sub>6</sub>S (MW = 507.61); mass spectroscopy found 507.9.

Anal. calcd for C<sub>26</sub>H<sub>25</sub>N<sub>3</sub>O<sub>6</sub>S: C, 61.53; H, 4.96; N, 8.28. Found: C, 61.70; H, 4.99; N, 8.22.

Step B            Preparation of 2-[N-( $\alpha$ -Amino)-N'-(benzyloxycarbonyl)-glyciny]-amino-5-nitrobenzophenone

Ammonia gas was bubbled into a solution 2-[N-( $\alpha$ -isopropylthio)-N'-(benzyloxycarbonyl)-glyciny]-amino-5-nitrobenzophenone (1 eq) in THF at 0°C. After 35 minutes mercury(II) chloride (1.1 eq) was added. The ice bath was removed and ammonia gas was continued to bubble through the suspension for 4 hours. The bubbler was removed and the reaction continued to stir for 16 hours. The mixture was filtered through celite washing with THF. The filtrate was concentrated *in vacuo*. The crude solid was used in step C without further purification.

Step C            Preparation of 3-(Benzyloxycarbonyl)-amino-2,3-dihydro-7-nitro-5-phenyl-1H-1,4-benzodiazepin-2-one

2-[N-( $\alpha$ -Amino)-N'-(benzyloxycarbonyl)-glyciny]-amino-5-nitrobenzophenone (1 eq) was treated with glacial acetic acid and ammonium acetate (4.7 eq). The suspension was stirred at ambient temperature for 21 hours. After concentrating the reaction in vacuo, the residue was partitioned between ethyl acetate and 1 N NaOH. The aqueous layer was back-extracted with ethyl acetate. The combined organics were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The residue was purified via flash chromatography eluting with a gradient of 2-3% isopropyl alcohol/methylene chloride.

-- 135 --

$C_{23}H_{18}N_4O_5$  (MW = 430.45); mass spectroscopy found (M+H) 431.2.

Anal. calcd for  $C_{23}H_{18}N_4O_5$ : C, 64.18; H, 4.22; N, 13.02. Found: C, 64.39; H, 4.30; N, 13.07.

5      Step D      Preparation of 3-(Benzyloxycarbonyl)-amino-2,3-dihydro-1-methyl-7-nitro-5-phenyl-1H-1,4-benzodiazepin-2-one

Following General Procedure 4-A and using 3-(benzyloxycarbonyl)-amino-2,3-dihydro-7-nitro-5-phenyl-1H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a yellow foam.

$C_{24}H_{20}N_4O_5$  (MW = 444.48); mass spectroscopy found (M+H) 445.2.

10      Anal. calcd for  $C_{24}H_{20}N_4O_5$ : C, 64.86; H, 4.54; N, 12.60. Found: C, 65.07; H, 4.55; N, 12.46.

Step E      Preparation of 3-Amino-1,3-dihydro-1-methyl-7-nitro-5-phenyl-2H-1,4-benzodiazepin-2-one

15      Following General Procedure 4-B and using 3-(benzyloxycarbonyl)-amino-2,3-dihydro-1-methyl-7-nitro-5-phenyl-1H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a yellow foam which was used immediately in Step F.

Step F      Preparation of 3-[N'-(tert-Butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-1-methyl-7-nitro-5-phenyl-1H-1,4-benzodiazepin-2-one

20      Following General Procedure D using N-Boc-L-alanine and 3-amino-1,3-dihydro-1-methyl-7-nitro-5-phenyl-2H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a yellow solid.

$C_{24}H_{27}N_5O_6$  (MW = 481.56); mass spectroscopy found (M+H) 482.3.

25      Anal. calcd for  $C_{24}H_{27}N_5O_6$ : C, 59.88; H, 5.61; N, 14.55. Found: C, 60.22; H, 5.75; N, 13.91.

Step G      Preparation of 3-(L-Alaninyl)-amino-2,3-dihydro-1-methyl-7-nitro-5-phenyl-1H-1,4-benzodiazepin-2-one

30      Following General Procedure 4-C using 3-[N'-(tert-butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-1-methyl-7-nitro-5-phenyl-1H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a yellow foam. The crude material was used immediately.

-- 136 --

## EXAMPLE 4-I

Synthesis of 3-(L-Alaninyl)amino-2,3-dihydro-1-methyl-5-(2-fluorophenyl)-1H-1,4-benzodiazepin-2-one5      Step A      Preparation of 3-Amino-1,3-dihydro-1-methyl-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-2-one

A flask was charged with 3-(benzyloxycarbonyl)-amino-7-bromo-2,3-dihydro-1-methyl-5-(2-fluorophenyl)-1H-1,4-benzodiazepin-2-one (1 eq.; Example 4-D, Step A) and 10% palladium on carbon. Methanol was added, and the flask was placed under a balloon of H<sub>2</sub>. The reaction was stirred for 21 hours. The mixture was filtered through celite washing with methanol. The filtrate was concentrated to a white solid.

C<sub>16</sub>H<sub>14</sub>FN<sub>3</sub>O (MW = 283.33); mass spectroscopy found (M+H) 284.1.

15      Step B      Preparation of 3-[N'-(tert-Butylcarbamate)-L-alaninyl]-amino-1,3-dihydro-1-methyl-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-2-one

Following General Procedure D using N-Boc-L-alanine and 3-amino-1,3-dihydro-1-methyl-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a white solid.

C<sub>24</sub>H<sub>27</sub>FN<sub>4</sub>O<sub>4</sub> (MW = 454.50); mass spectroscopy found (M+H) 455.4.

20      Anal. calcd for C<sub>24</sub>H<sub>27</sub>FN<sub>4</sub>O<sub>4</sub>: C, 63.44; H, 5.95; N, 12.33. Found: C, 63.64; H, 6.08; N, 12.16.

Step C      Preparation of 3-(L-Alaninyl)-amino-7-bromo-1,3-dihydro-1-methyl-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-2-one

25      Following General Procedure 4-C using 3-[N'-(tert-butylcarbamate)-L-alaninyl]-amino-1,3-dihydro-1-methyl-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a white foam. The crude material was used immediately.

## EXAMPLE 4-J

30      Synthesis of 3-(L-Alaninyl)-amino-2,3-dihydro-1-methyl-5-(3-fluorophenyl)-1H-1,4-benzodiazepin-2-oneStep A      Preparation of 2-Amino-3'-fluorobenzophenone

-- 137 --

A solution of 3-bromofluorobenzene (1 eq.) in THF was cooled to  $-78^{\circ}\text{C}$  under nitrogen and treated with *tert*-butyllithium (2.05 eq., 1.6 M solution in pentane) at a rate of 40 mL/h. The internal temperature did not rise above  $-74^{\circ}\text{C}$ . The orange solution was stirred at  $-78^{\circ}\text{C}$  for 30 minutes prior to the addition of anthranilonitrile (0.6 eq.) as a solution in THF. The reaction was warmed to  $0^{\circ}\text{C}$  and stirred for 2 hours. 3N HCl was added to the mixture and stirring continued for 30 minutes. The reaction was diluted with ethyl acetate and the layers were separated. The aqueous layer was back-extracted thrice with ethyl acetate. The combined extracts were washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. The residue was purified via HPLC eluting with 93:7 hexanes/ethyl acetate.

$\text{C}_{13}\text{H}_{10}\text{FNO}$  (MW = 215.24); mass spectroscopy found (M+H) 216.3.

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.44-7.19 (6H, m), 6.74 (1H, d,  $J=8.0$  Hz), 6.61 (1H, dd,  $J=0.94, 7.9$  Hz), 6.10 (2H, bs).

15 **Step B**      Preparation of 2-[N-( $\alpha$ -Isopropylthio)-N'-(benzyloxycarbonyl)-glycinyll]-amino-3'-fluorobenzophenone

A solution of  $\alpha$ -(isopropylthio)-N-(benzyloxycarbonyl)glycine (1 eq; prepared according to Zoller, V. and D. Ben-Ishai *Tetrahedron* **1975**, *31*, 863, incorporated herein by reference) in dry THF was cooled to  $0^{\circ}\text{C}$  and treated with oxalyl chloride (1 eq.) and 3 drops of DMF. After stirring for 15 minutes at  $0^{\circ}\text{C}$ , the cooling bath was removed and stirring continued at ambient temperature for 40 minutes. The solution was recooled to  $0^{\circ}\text{C}$ . A solution of 2-amino-3'-fluorobenzophenone (0.9 eq.) and 4-methylmorpholine (2.0 eq.) in dry THF was added via cannulation to the acid chloride. The cooling bath was removed and the reaction stirred at ambient temperature for 5 hours. The reaction was diluted with methylene chloride and washed with 0.5 M citric acid, saturated aqueous  $\text{NaHCO}_3$ , and brine. The organic phase was dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. The residue was purified via preparative LC2000 eluting with a gradient of 15-20% ethyl acetate/hexanes giving an off-white foam.

30  $\text{C}_{26}\text{H}_{25}\text{N}_2\text{O}_4\text{S}$  (MW = 480.60); mass spectroscopy found (M+ $\text{NH}_4^+$ ) 498.3.

-- 138 --

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  11.39 (1H, s), 8.59 (1H, d,  $J=6.0$  Hz), 7.63-7.55 (2H, m), 7.48-7.27 (9H, m), 7.14 (1H, dt,  $J=1.2, 8.4$  Hz), 5.94 (1H, d,  $J=7.2$  Hz), 5.58 (1H, d,  $J=8.7$  Hz), 5.17 (2H, ABq,  $J=14.7$  Hz), 3.25 (1H, sep,  $J=6.6$  Hz), 1.44 (3H, d,  $J=6.0$  Hz), 1.28 (3H, d,  $J=6.6$  Hz).

5    Step C            Preparation of 2-[N-( $\alpha$ -Amino)-N'-(benzyloxycarbonyl)-glycinyll]-amino-3'-fluorobenzophenone

Ammonia gas was bubbled into a solution of 2-[N-( $\alpha$ -isopropylthio)-N'-(benzyloxycarbonyl)-glycinyll]-amino-3'-fluorobenzophenone (1 eq) in THF at  $0^\circ\text{C}$ . After 35 minutes mercury(II) chloride (1.1 eq) was added. The ice bath  
10    was removed and ammonia gas was continued to bubble through the suspension for 4 hours. The bubbler was removed and the reaction continued to stir for 16 hours. The mixture was filtered through celite washing with THF. The filtrate was concentrated *in vacuo*. The crude solid was used in step D without further purification.

15    Step D            Preparation of 3-(Benzyloxycarbonyl)-amino-2,3-dihydro-5-(3-fluorophenyl)-1H-1,4-benzodiazepin-2-one

2-[N-( $\alpha$ -Amino)-N'-(benzyloxycarbonyl)-glycinyll]-amino-3'-fluorobenzophenone (1 eq) was treated with glacial acetic acid and ammonium acetate (4.7 eq). The suspension was stirred at ambient temperature for 21 hours.  
20    After concentrating the reaction in *vacuo*, the residue was partitioned between ethyl acetate and 1 N NaOH. The aqueous layer was back-extracted with ethyl acetate. The combined organics were washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. The residue was purified via flash chromatography eluting with a gradient of 2-3% isopropyl alcohol/methylene chloride.

25             $\text{C}_{23}\text{H}_{18}\text{FN}_3\text{O}_3$  (MW = 403.44); mass spectroscopy found (M+H) 404.4.

Anal. calcd for  $\text{C}_{23}\text{H}_{18}\text{FN}_3\text{O}_3 \cdot 0.5\text{H}_2\text{O}$ : C, 66.98; H, 4.64; N, 10.18.

Found: C, 67.20; H, 4.64; N, 9.77.

Step E            Preparation of 3-(Benzyloxycarbonyl)-amino-2,3-dihydro-1-methyl-5-(3-fluorophenyl)-1H-1,4-benzodiazepin-2-one



-- 139 --

Following General Procedure 4-A and using 3-(benzyloxycarbonyl)-amino-2,3-dihydro-5-(3-fluorophenyl)-1H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a yellow foam.

$C_{24}H_{20}FN_3O_3$  (MW = 417.47); mass spectroscopy found (M+H) 418.3.

5      Anal. calcd for  $C_{24}H_{20}FN_3O_3$ : C, 69.06; H, 4.83; N, 10.07. Found: C, 69.33; H, 4.95; N, 9.82.

Step F      Preparation of 3-Amino-1,3-dihydro-1-methyl-5-(3-fluorophenyl)-2H-1,4-benzodiazepin-2-one

10      Following General Procedure 4-B and using 3-(benzyloxycarbonyl)-amino-2,3-dihydro-1-methyl-5-(3-fluorophenyl)-1H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a yellow foam which was used immediately in Step G.

Step G      Preparation of 3-[N'-(tert-Butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-1-methyl-5-(3-fluorophenyl)-1H-1,4-benzodiazepin-2-one

15      Following General Procedure D using N-Boc-L-alanine and 3-amino-1,3-dihydro-1-methyl-5-(3-fluorophenyl)-2H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a yellow solid.

$C_{24}H_{27}FN_4O_4$  (MW = 454.50); mass spectroscopy found (M+H) 455.3.

20      Anal. calcd for  $C_{24}H_{27}FN_4O_4$ : C, 63.42; H, 5.99; N, 12.33. Found: C, 63.34; H, 6.01; N, 12.08.

Step H      Preparation of 3-(L-Alaninyl)-amino-2,3-dihydro-1-methyl-5-(3-fluorophenyl)-1H-1,4-benzodiazepin-2-one

25      Following General Procedure 4-C using 3-[N'-(tert-butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-1-methyl-5-(3-fluorophenyl)-1H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a yellow foam. The crude material was used immediately.

#### EXAMPLE 4-K

30      Synthesis of 3-(L-Alaninyl)amino-2,3-dihydro-1-methyl-5-(4-fluorophenyl)-1H-1,4-benzodiazepin-2-one

Step A      Preparation of 2-Amino-4'-fluorobenzophenone

-- 140 --

A solution of 4-bromofluorobenzene (1 eq.) in THF was cooled to -78°C under nitrogen and treated with *tert*-butyllithium (2.05 eq., 1.6 M solution in pentane) at a rate of 40 mL/h. The internal temperature did not rise above -74°C. The orange solution was stirred at -78°C for 30 minutes prior to the addition of anthranilonitrile (0.6 eq.) as a solution in THF. The reaction was warmed to 0°C and stirred for 2 hours. 3N HCl was added to the mixture and stirring continued for 30 minutes. The reaction was diluted with ethyl acetate and the layers were separated. The aqueous layer was back-extracted thrice with ethyl acetate. The combined extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The residue was purified via HPLC eluting with 93:7 hexanes/ethyl acetate.

C<sub>13</sub>H<sub>10</sub>FNO (MW = 215.24); mass spectroscopy found (M+H) 216.3.

Anal. calcd for C<sub>13</sub>H<sub>10</sub>FNO: C, 72.55; H, 4.68; N, 6.51. Found: C, 72.80; H, 4.51; N, 6.74.

15 **Step B**      Preparation of 2-[N-( $\alpha$ -Isopropylthio)-N'-(benzyloxycarbonyl)-glycinyll-amino-4'-fluorobenzophenone

A solution of  $\alpha$ -(isopropylthio)-N-(benzyloxycarbonyl)glycine (1 eq; prepared according to Zoller, V. and D. Ben-Ishai *Tetrahedron* **1975**, *31*, 863, incorporated herein by reference) in dry THF was cooled to 0°C and treated with oxalyl chloride (1 eq.) and 3 drops of DMF. After stirring for 15 minutes at 0°C, the cooling bath was removed and stirring continued at ambient temperature for 40 minutes. The solution was recooled to 0°C. A solution of 2-amino-4'-fluorobenzophenone (0.9 eq.) and 4-methylmorpholine (2.0 eq.) in dry THF was added via cannulation to the acid chloride. The cooling bath was removed and the reaction stirred at ambient temperature for 5 hours. The reaction was diluted with methylene chloride and washed with 0.5 M citric acid, saturated aqueous NaHCO<sub>3</sub>, and brine. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The residue was purified via preparative LC2000 eluting with a gradient of 15–20% ethyl acetate/hexanes giving an off-white foam.

-- 141 --

$C_{26}H_{25}N_2O_4S$  (MW = 480.60); mass spectroscopy found  $(M+NH_4^+)$   
498.2.

$^1H$  NMR (300 MHz,  $CDCl_3$ )  $\delta$  11.28 (1H, s), 8.56 (1H, d,  $J=8.4$  Hz),  
7.78-7.73 (2H, m), 7.61-7.53 (2H, m), 7.36-7.32 (5H, m), 7.20-7.14 (3H, m),  
5 5.98 (1H, d,  $J=7.5$  Hz), 5.57 (1H, d,  $J=7.8$  Hz), 5.16 (2H, ABq,  $J=14.7$  Hz),  
3.25 (1H, sep,  $J=6.0$  Hz), 1.43 (3H, d,  $J=6.3$  Hz), 1.27 (3H, d,  $J=6.6$  Hz).

Step C      Preparation of 2-[N-( $\alpha$ -Amino)-N'-(benzyloxycarbonyl)-glycinyll]-  
amino-4'-fluorobenzophenone

Ammonia gas was bubbled into a solution of 2-[N-( $\alpha$ -isopropylthio)-N'-  
10 (benzyloxycarbonyl)-glycinyll]-amino-3'-fluorobenzophenone (1 eq) in THF at  
0°C. After 35 minutes mercury(II) chloride (1.1 eq) was added. The ice bath  
was removed and ammonia gas was continued to bubble through the suspension  
for 4 hours. The bubbler was removed and the reaction continued to stir for 16  
hours. The mixture was filtered through celite washing with THF. The filtrate  
15 was concentrated *in vacuo*. The crude solid was used in step D without further  
purification.

Step D      Preparation of 3-(Benzyloxycarbonyl)amino-2,3-dihydro-5-(4-  
fluorophenyl)-1H-1,4-benzodiazepin-2-one

2-[N-( $\alpha$ -Amino)-N'-(benzyloxycarbonyl)-glycinyll]-amino-4'-  
20 fluorobenzophenone (1 eq) was treated with glacial acetic acid and ammonium  
acetate (4.7 eq). The suspension was stirred at ambient temperature for 21 hours.  
After concentrating the reaction in *vacuo*, the residue was partitioned between  
ethyl acetate and 1 N NaOH. The aqueous layer was back-extracted with ethyl  
acetate. The combined organics were washed with brine, dried over  $Na_2SO_4$ ,  
25 filtered, and concentrated. The residue was purified via flash chromatography  
eluting with a gradient of 2-3% isopropyl alcohol/methylene chloride.

$C_{23}H_{18}FN_3O_3$  (MW = 403.44); mass spectroscopy found  $(M+H)$  404.4.

Anal. calcd for  $C_{23}H_{18}FN_3O_3 \cdot 1.25H_2O$ : C, 64.85; H, 4.85. Found: C,  
64.80; H, 4.55.

-- 142 --

**Step E**      Preparation of 3-(Benzyloxycarbonyl)-amino-2,3-dihydro-1-methyl-5-(4-fluorophenyl)-1H-1,4-benzodiazepin-2-one

Following General Procedure 4-A and using 3-(benzyloxycarbonyl)-amino-2,3-dihydro-5-(4-fluorophenyl)-1H-1,4-benzodiazepin-2-one, the title intermediate  
5 was prepared as a yellow foam.

$C_{24}H_{20}FN_3O_3$  (MW = 417.47); mass spectroscopy found (M+H) 418.2.

Anal. calcd for  $C_{24}H_{20}FN_3O_3$ : C, 69.06; H, 4.83; N, 10.07. Found: C, 69.35; H, 4.93; N, 9.97.

10      **Step F**      Preparation of 3-Amino-1,3-dihydro-1-methyl-5-(4-fluorophenyl)-2H-1,4-benzodiazepin-2-one

Following General Procedure 4-B and using 3-(benzyloxycarbonyl)-amino-2,3-dihydro-1-methyl-5-(4-fluorophenyl)-1H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a yellow foam which was used immediately in Step G.

15      **Step G**      Preparation of 3-[N'-(*tert*-Butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-1-methyl-5-(3-fluorophenyl)-1H-1,4-benzodiazepin-2-one

Following General Procedure D using N-Boc-L-alanine and 3-amino-1,3-dihydro-1-methyl-5-(3-fluorophenyl)-2H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a yellow solid.

20       $C_{24}H_{27}FN_4O_4$  (MW = 454.50); mass spectroscopy found (M+H) 455.4.

Anal. calcd for  $C_{24}H_{27}FN_4O_4 \cdot 1.5H_2O$ : C, 59.86; H, 6.28; N, 11.64.

Found: C, 60.04; H, 5.62; N, 11.27.

**Step H**      Preparation of 3-(L-Alaninyl)-amino-2,3-dihydro-1-methyl-5-(4-fluorophenyl)-1H-1,4-benzodiazepin-2-one

25      Following General Procedure 4-C using 3-[N'-(*tert*-butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-1-methyl-5-(4-fluorophenyl)-1H-1,4-benzodiazepin-2-one, the title intermediate was prepared as a yellow foam. The crude material was used immediately.

30

**EXAMPLE 4-L**

Synthesis of 3-(N'-L-Alaninyl)amino-2,3-dihydro-1-isobutyl-

-- 143 --

5-phenyl-1H-1,4-benzodiazepin-2-oneStep A

1,3-Dihydro-5-phenyl-2H-1,4-benzodiazepin-2-one (prepared according to the procedure of M. G. Bock et al., *J. Org. Chem.* **1987**, **52**, 3232-3239, incorporated herein by reference) was alkylated with isobutyl iodide using General Procedure 8-G to afford 1,3-dihydro-1-isobutyl-5-phenyl-2H-1,4-benzodiazepin-2-one.

Step B

Following General Procedures 4-D and 4-F and using the product from Step A, 3-amino-1,3-dihydro-1-isobutyl-5-phenyl-2H-1,4-benzodiazepin-2-one was prepared.

Step C

The product from Step B and N-Boc-L-alanine (Sigma) were coupled using General Procedure D, followed by removal of the Boc group using General Procedure 8-J, to afford 3-(N'-L-alaninyl)amino-1,3-dihydro-1-isobutyl-5-phenyl-2H-1,4-benzodiazepin-2-one.

By substituting isopropyl iodide, *n*-propyl iodide, cyclopropylmethyl iodide and ethyl iodide for isobutyl iodide in Step A above, the following additional intermediates were prepared:

20        3-(N'-L-alaninyl)amino-1,3-dihydro-1-isopropyl-5-phenyl-2H-1,4-benzodiazepin-2-one

          3-(N'-L-alaninyl)amino-1,3-dihydro-1-propyl-5-phenyl-2H-1,4-benzodiazepin-2-one

25        3-(N'-L-alaninyl)amino-1,3-dihydro-1-cyclopropylmethyl-5-phenyl-2H-1,4-benzodiazepin-2-one

30        3-(N'-L-alaninyl)amino-1,3-dihydro-1-ethyl-5-phenyl-2H-1,4-benzodiazepin-2-one.

-- 144 --

**EXAMPLE 4-M**Synthesis of 3-(N'-L-Alaninyl)amino-1-methyl-5-phenyl-1,3,4,5-tetrahydro-2H-1,5-benzodiazepin-2-oneStep A

5           1,3,4,5-Tetrahydro-5-phenyl-2H-1,5-benzodiazepin-2-one (CAS No. 32900-17-7) was methylated using General Procedure 4-I to afford 1-methyl-5-phenyl-1,3,4,5-tetrahydro-2H-1,5-benzodiazepin-2-one.

Step B

10           Following General Procedures 4-E and 4-F and using the product from Step A, 3-amino-1-methyl-5-phenyl-1,3,4,5-tetrahydro-2H-1,5-benzodiazepin-2-one was prepared.

Step C

15           The product from Step B and N-Boc-L-alanine (Sigma) were coupled using General Procedure D, followed by removal of the Boc group using General Procedure 8-N, to afford 3-(N'-L-alaninyl)amino-1-methyl-5-phenyl-1,3,4,5-tetrahydro-2H-1,5-benzodiazepin-2-one.

**EXAMPLE 4-N**Synthesis of 3-(N'-L-Alaninyl)amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

20           3-Amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine (CAS No. 131604-75-6) was coupled with N-Boc-L-alanine (Sigma) using General Procedure D, followed by removal of the Boc group using General Procedure 4-N, to afford the title compound.

25

**EXAMPLE 4-O**Synthesis of 3-((R)-Hydrazinopropionyl)amino-2,3-dihydro-1-methyl-5-phenyl)-1H-1,4-benzodiazepin-2-one

30    Part 1       Synthesis of (R)-N,N'-Di-BOC-2-Hydrazinopropionic Acid  
                  Step A

-- 145 --

To (S)-(-)-4-benzyl-2-oxazolidanone (Aldrich) in THF cooled to -50°C was added n-butyl lithium 1.1 eq. (1.6 M in hexane) dropwise. The reaction mixture was allowed to warm to -20°C and then was re-cooled to -78°C and propionyl chloride (1.1 eq) was added in one portion. The reaction mixture  
5 was allowed to stir an additional 15 min. at -78°C and then was allowed to warm to room temperature. The reaction was then quenched with a saturated solution of sodium bicarbonate and extracted with ethyl acetate. The organic extracts were washed with water, followed by brine and then dried over sodium sulfate, filtered and concentrated to give (S)-(-)-3-propionyl-4-benzyl-2-oxazolidanone.

10 Step B

To a solution of (S)-(-)-3-propionyl-4-benzyl-2-oxazolidanone in THF at -78°C was added KHMDS (1.05 eq.) (Aldrich) dropwise. The reaction mixture was allowed to stir at -78°C for 30 min. and then a precooled solution of di-tert-butyl-azodicarboxylate (Aldrich) was added via a cannula. After 5 min. 2.6 eq. of  
15 acetic acid was added. The reaction mixture was then extracted with dichloromethane and the organic layer was washed with 1M potassium phosphate. The organic layer was then dried over sodium sulfate, filtered and concentrated to give (S)-(-)-3-[(R)-N,N'-di-BOC-2-hydrazinopropionyl]-4-benzyl-2-oxazolidanone.

Step C

20 To (S)-(-)-3-[(R)-N,N'-di-BOC-2-hydrazinopropionyl]-4-benzyl-2-oxazolidanone (0.49 moles) at 0°C in 8 mL of THF and 3 mL of water was added LiOH (1.7 eq.) and H<sub>2</sub>O<sub>2</sub> (3.0 eq.) and the reaction mixture was stirred at room temperature for 3 hours. The reaction mixture was then poured into a separatory funnel and diluted with water. The aqueous mixture was extracted with ethyl  
25 acetate and then acidified to pH 2.0 with 1N HCl and extracted with ethyl acetate. The organic layer was then dried over sodium sulfate, filtered and solvent removed to give (R)-N,N'-di-BOC-2-hydrazinopropionic acid which was used without further purification.

-- 146 --

**Part 2**

3-Amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one was coupled to (R)-N,N'-di-BOC-2-hydrazinopropionic acid using General Procedure D. The Boc group was removed by dissolving the Boc-protected compound in a 1:1-2:1 mixture of CH<sub>2</sub>Cl<sub>2</sub> and trifluoroacetic acid. The resulting solution was stirred until tlc indicated complete conversion, typically 2 hours. The solution was then stripped to dryness and the residue was taken up in ethyl acetate or CH<sub>2</sub>Cl<sub>2</sub>. The solution was washed with saturated aqueous NaHCO<sub>3</sub> and the aqueous phase was adjusted to a basic pH, then extracted with ethyl acetate or CH<sub>2</sub>Cl<sub>2</sub>. The organic phase was washed with saturated aqueous NaCl and dried over MgSO<sub>4</sub>. The solution was stripped free of solvent on a rotary evaporator to afford the title compound.

**EXAMPLE 4-P**

Synthesis of 3-Amino-2,4-dioxo-1,5-bis-(1-methylethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

Step A      Synthesis of 2,4-Dioxo-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

2,4-Dioxo-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine (CAS No. 49799-48-6) was prepared from 1,2-phenylenediamine (Aldrich) and malonic acid (Aldrich) using the procedure of Claremon, D. A. et al., PCT Application: WO 96/40655, incorporated herein by reference.

Step B      Synthesis of 2,4-Dioxo-1,5-bis-(1-methylethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

2,4-Dioxo-1,5-bis-(1-methylethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine (CAS No. 113021-84-4) was prepared following General Procedure 4-M using the product from Step A and 2-iodopropane (Aldrich). Purification was by flash chromatography eluting with EtOAc/hexanes (3:7 gradient to 1:1), then recrystallization from EtOAc/hexanes.

Step C      Synthesis of 3-Azido-2,4-dioxo-1,5-bis-(1-methylethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine



-- 147 --

Following General Procedure 4-K using the product from Step B, 3-azido-2,4-dioxo-1,5-bis-(1-methylethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine (CAS No. 186490-50-6) was prepared as a white solid. The product was purified by flash chromatography eluting with hexanes/EtOAc (4:1) to provide a separable  
5 23:1 mixture of pseudo-axial/pseudo-equatorial azides. The pure pseudo-axial azide was used in the next step.

**Step D**      Synthesis of 3-Amino-2,4-dioxo-1,5-bis-(1-methylethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

Following General Procedure 4-L using the product from Step C, 3-amino-  
10 2,4-dioxo-1,5-bis-(1-methylethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine (CAS No. 186490-51-7) was prepared as a white solid. Purification was by flash chromatography eluting with CH<sub>2</sub>Cl<sub>2</sub>/MeOH (98:2 gradient to 95:5). The isolated pseudo-axial amine atropisomer was completely converted to the pseudo-equatorial amine atropisomer by heating in toluene to 100-105 °C for 15 minutes, and the  
15 pseudo-equatorial amine atropisomer was used in the next step. The isomers were distinguished by <sup>1</sup>H-NMR in CDCl<sub>3</sub>. Selected <sup>1</sup>H-NMR (CDCl<sub>3</sub>): Pseudo-axial amine 4.40 (s, 1H); Pseudo-equatorial amine 3.96 (s, 1H).

**EXAMPLE 4-Q**

20      Synthesis of 3-(R-2-Thienylglyciny)l-amino-2,4-dioxo-1,5-bis-(1-methylethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine Hydrochloride

**Step A**      Synthesis of N-(t-Butoxycarbonyl)-R-2-thienylglycine

N-(t-Butoxycarbonyl)-R-2-thienylglycine (CAS No. 74462-03-1) was  
25 prepared from L-α-(2-thienyl)glycine (Sigma) by the procedure described in Bodansky, M. et al; *The Practice of Peptide Synthesis*; Springer Verlag; 1994, p. 17, incorporated herein by reference.

**Step B**      Synthesis of 3-[N'-(t-Butoxycarbonyl)-R-2-thienylglyciny]l-amino-2,4-dioxo-1,5-bis-(1-methylethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine  
30

-- 148 --

Following General Procedure J above using the product from Example 4-P and the product from Step A above, 3-[N'-(*t*-butoxycarbonyl)-R-2-thienylglyciny]-amino-2,4-dioxo-1,5-bis-(1-methylethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was prepared as a white foam. Purification was by flash chromatography eluting with CH<sub>2</sub>Cl<sub>2</sub>/EtOAc (9:1 gradient to 5:1).

Step C      Synthesis of 3-(R-2-Thienylglyciny)amino-2,4-dioxo-1,5-bis-(1-methylethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine Hydrochloride

Following General Procedure 4-N above using the product from Step B, the title compound was prepared as a white solid.

#### EXAMPLE 4-R

Synthesis of 3-(L-Alaniny)-amino-2,4-dioxo-1,5-bis-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine Hydrochloride

Step A      Synthesis of 2,4-Dioxo-1,5-bis-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

2,4-Dioxo-1,5-bis-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine (CAS No. 23954-54-3) was prepared following General Procedure 4-M using the product from Example 4-P, Step A and iodomethane (Aldrich). The white solid product precipitated during partial concentration of the reaction after work-up, and was isolated by filtration.

Step B      Synthesis of 3-Azido-2,4-dioxo-1,5-bis-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

For this substrate, General Procedure 4-K was modified in the following manner. Initially the product from Step A was suspended (not a solution) in THF at -78°C, and following addition of the KN(TMS)<sub>2</sub> solution, this suspension was allowed to warm to -35°C over a period of 12 minutes, during which the suspension became a solution, and was re-cooled to -78°C; then treated as described in the General Procedure. 3-Azido-2,4-dioxo-1,5-bis-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was purified by flash chromatography eluting

-- 149 --

with  $\text{CHCl}_3/\text{EtOAc}$  (7:1), then trituration from hot  $\text{CHCl}_3$  with hexanes and cooled to  $-23^\circ\text{C}$ . The product was isolated as a white solid.

Step C      Synthesis of 3-Amino-2,4-dioxo-1,5-bis-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

5            Following General Procedure 4-L using the product from Step B, 3-amino-2,4-dioxo-1,5-bis-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was prepared as a white solid. The crude product was used without further purification.

Step D      Synthesis of 3-[N'-(t-Butoxycarbonyl)-L-alaninyl]-amino-2,4-dioxo-1,5-bis-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

10           Following General Procedure I above using N-Boc-L-alanine (Novabiochem) and the product from Step C, 3-[N'-(t-butoxycarbonyl)-L-alaninyl]-amino-2,4-dioxo-1,5-bis-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was prepared as a white foam. Purification was by flash chromatography eluting with  $\text{CH}_2\text{Cl}_2/\text{EtOAc}$  (2:1 gradient to 1:1).

15      Step E      Synthesis of 3-(L-alaninyl)-amino-2,4-dioxo-1,5-bis-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine Hydrochloride

Following General Procedure 4-N above using the product from Step D, the title compound was prepared as an off-white amorphous solid.

20

**EXAMPLE 4-S**

Synthesis of 3-(L-Alaninyl)amino-2,4-dioxo-1,5-bis-(2-methylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine Hydrochloride

Step A      Synthesis of 2,4-Dioxo-1,5-bis-(2-methylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

25           2,4-Dioxo-1,5-bis-(2-methylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was prepared following General Procedure 4-M using the product from Example 4-P, Step A and 1-iodo-2-methylpropane (Aldrich). Purification was by flash chromatography eluting with  $\text{EtOAc}/\text{hexanes}$  (3:7 gradient to 1:1), then recrystallization from  $\text{EtOAc}/\text{hexanes}$ .

30      Step B      Synthesis of 3-Azido-2,4-dioxo-1,5-bis-(2-methylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

-- 150 --

Following General Procedure 4-K (a precipitate formed during the addition of the  $\text{KN}(\text{TMS})_2$ , but dissolved upon addition of the trisyl azide) using the product from Step A, 3-azido-2,4-dioxo-1,5-bis-(2-methylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was prepared as a white solid. The product  
5 was purified by flash chromatography eluting with hexanes/EtOAc (4:1) and a second flash chromatography eluting with  $\text{CH}_2\text{Cl}_2$ /hexanes/EtOAc (10:10:1 gradient to 8:6:1).

Step C      Synthesis of 3-Amino-2,4-dioxo-1,5-bis-(2-methylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

10      Following General Procedure 4-L using the product from Step B, 3-amino-2,4-dioxo-1,5-bis-(2-methylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was prepared as a white solid. Purification was by flash chromatography eluting with  $\text{CH}_2\text{Cl}_2$ /MeOH (98:2 gradient to 95:5, with 5%  $\text{NH}_3$  in the MeOH).

Step D      Synthesis of 3-[N'-(t-Butoxycarbonyl)-L-alaninyl]-amino-2,4-dioxo-1,5-bis-(2-methylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

15      Following General Procedure I above using N-Boc-L-alanine (Novabiochem) and the product from Step C, 3-[N'-(t-butoxycarbonyl)-L-alaninyl]-amino-2,4-dioxo-1,5-bis-(2-methylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was prepared as a white foam. Purification was by flash  
20 chromatography eluting with  $\text{CH}_2\text{Cl}_2$ /EtOAc (3:1 gradient to 3:2).

Step E      Synthesis of 3-(L-Alaninyl)-amino-2,4-dioxo-1,5-bis-(2-methylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine Hydrochloride

25      Following General Procedure 4-N above using the product from Step D, the title compound was prepared as an amorphous white solid.

### EXAMPLE 4-T

**Step A**      **Synthesis of 3-[N'-(*t*-Butoxycarbonyl)-S-phenylglycinyll-amino-2,4-dioxo-1,5-bis-(2-methylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine**

**Step B**      Synthesis of 3-(S-PhenylglycinyI)-amino-2,4-dioxo-1,5-bis-(2-methylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine Hydrochloride

### EXAMPLE 4-U

**Step A**      **Synthesis of 2,4-Dioxo-1,5-bis-(cyclopropylmethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine**

**Step B**      **Synthesis of 3-Azido-2,4-dioxo-1,5-bis-(cyclopropylmethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine**

-- 152 --

For this substrate General Procedure 4-K was modified in the following manner. Initially the product from Step A was suspended (not a solution) in THF at -78°C, and following addition of the KN(TMS)<sub>2</sub> solution, this suspension was allowed to warm to -30°C, during which the suspension became a solution, and  
5 was re-cooled to -78°C. Upon re-cooling to -78°C a precipitate began to form, therefore the reaction flask containing the mixture was partially raised above the cooling bath until the internal temperature rose to -50°C; then the trisyl azide solution was added. The cooling bath was removed and the mixture allowed to warm to -20°C whereupon the mixture had become a nearly homogenous solution,  
10 and the AcOH was added. Then the mixture was treated as described in the general procedure. 3-Azido-2,4-dioxo-1,5-bis-(cyclopropylmethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was purified by trituration with hot to room temperature EtOAc, followed by recrystallization from hot to -23°C CHCl<sub>3</sub>/EtOAc/EtOH (5:5:1) and isolated as a white solid.

15 Step C      Synthesis of 3-Amino-2,4-dioxo-1,5-bis-(cyclopropylmethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

Following General Procedure 4-L using the product from Step B, 3-amino-2,4-dioxo-1,5-bis-(cyclopropylmethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was prepared as a white solid. Purification was by flash chromatography eluting  
20 with CH<sub>2</sub>Cl<sub>2</sub>/MeOH (98:2 gradient to 95:5, with 5% NH<sub>3</sub> in the MeOH) followed by recrystallization from warm CH<sub>2</sub>Cl<sub>2</sub>/hexanes (1:1) to -23°C.

Step D      Synthesis of 3-[N'-(t-Butoxycarbonyl)-L-alaninyl]-amino-2,4-dioxo-1,5-bis-(cyclopropylmethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

25      Following General Procedure I above using N-Boc-L-alanine (Novabiochem) and the product from Step C, 3-[N'-(t-butoxycarbonyl)-L-alaninyl]-amino-2,4-dioxo-1,5-bis-(cyclopropylmethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was prepared as a white foam. Purification was by flash chromatography eluting with CH<sub>2</sub>Cl<sub>2</sub>/EtOAc (3:1 gradient to 2:1).

-- 153 --

**Step E**      Synthesis of 3-(L-Alaninyl)-amino-2,4-dioxo-1,5-bis-(cyclopropylmethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine Hydrochloride

Following General Procedure 4-N above using the product from Step D,  
5      the title compound was prepared as an off-white solid.

**EXAMPLE 4-V**

Synthesis of 3-(L-Alaninyl)-amino-2,4-dioxo-1,5-bis-(2,2-dimethylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine Hydrochloride

10      **Step A**      Synthesis of 2,4-Dioxo-1,5-bis-(2,2-dimethylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

To a stirred suspension of the product from Example 4-P, Step A (1.0 eq., 17.08 g) in DMSO (500 mL) at room temperature was added neopentyl iodide (43.01 g, 2.24 eq., Aldrich) and Cs<sub>2</sub>CO<sub>3</sub> (72.65 g, 2.3 eq., Aldrich). The  
15      resulting mixture was heated to 75°C for 30 minutes, then additional Cs<sub>2</sub>CO<sub>3</sub> (31.59 g, 1.0 eq.) was added and the mixture rapidly stirred at 75°C for 6 hours. The mixture was allowed to cool and H<sub>2</sub>O (500 mL) and EtOAc (1000 mL) were added. The phases were partitioned and the organic phase washed with H<sub>2</sub>O (1x500 mL), 1 M aq. HCl (2x500 mL), and brine (1x500 mL). Then, the organic  
20      phase was dried over MgSO<sub>4</sub>, filtered, concentrated, and purified by flash chromatography eluting with hexanes/EtOAc (3:2 gradient to 2:3) to provide 2,4-dioxo-1,5-bis-(2,2-dimethylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine as a white solid.

25      **Step B**      Synthesis of 3-Azido-2,4-dioxo-1,5-bis-(2,2-dimethylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

Following General Procedure 4-K using the product from Step A, 3-azido-2,4-dioxo-1,5-bis-(2,2-dimethylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was prepared as a white solid. The product was purified by flash chromatography eluting with hexanes/CH<sub>2</sub>Cl<sub>2</sub>/EtOAc (10:5:1 gradient to 5:5:1) to provide a  
30      separable 13:1 mixture of pseudo-axial/pseudo-equatorial azides. The pure

-- 154 --

pseudo-axial azide was used in the next step. Selected  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):

Pseudo-axial azide 5.12 (s, 1H); Pseudo-equatorial azide 4.03 (s, 1H).

**Step C**      Synthesis of 3-Amino-2,4-dioxo-1,5-bis-(2,2-dimethylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

- 5      Following General Procedure 4-L using the product from Step B, 3-amino-2,4-dioxo-1,5-bis-(2,2-dimethylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was prepared as a white solid. Purification was by flash chromatography eluting with  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  (98:2 gradient to 95:5, with 5%  $\text{NH}_3$  in the MeOH). The isolated white solid product was identified as a ~4:1 mixture of pseudo-axial and
- 10      pseudo-equatorial amines atropisomers by  $^1\text{H-NMR}$ . The mixture was heated in toluene to 100 °C for 20 minutes, then re-concentrated to provide the pure pseudo-equatorial amine atropisomer, as a white solid, and this was for the next step. Selected  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): Pseudo-axial amine 4.59 (s, 1H); Pseudo-equatorial amine 4.03 (s, 1H).

- 15      **Step D**      Synthesis of 3-[N'-(*t*-Butoxycarbonyl)-L-alaninyl]-amino-2,4-dioxo-1,5-bis-(2,2-dimethylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

- Following General Procedure I above using N-Boc-L-alanine (Novabiochem) and the product from Step C, 3-[N'-(*t*-butoxycarbonyl)-L-alaninyl]-amino-2,4-dioxo-1,5-bis-(2,2-dimethylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was prepared as a white foam. Purification was by flash
- 20      chromatography eluting with  $\text{CH}_2\text{Cl}_2/\text{EtOAc}$  (4:1 gradient to 5:2).

- Step E**      Synthesis of 3-(L-Alaninyl)-amino-2,4-dioxo-1,5-bis-(2,2-dimethylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine Hydrochloride
- 25

Following General Procedure 4-N above using the product from Step D, the title compound was prepared as an off-white solid.



-- 155 --

## EXAMPLE 4-W

Synthesis of 3-(L-Alaninyl)amino-2,4-dioxo-1,5-bis-phenyl-  
2,3,4,5-tetrahydro-1H-1,5-benzodiazepine hydrochlorideStep A      Synthesis of 2,4-Dioxo-1,5-bis-phenyl-2,3,4,5-tetrahydro-1H-1,5-  
benzodiazepine

5                    This procedure is a modification of the procedure described in Chan, D. M. T. *Tetrahedron Lett.* **1996**, 37, 9013-9016, incorporated herein by reference. A mixture of the product from Example 4-P, Step A (1.0 eq., 7.50 g), Ph<sub>3</sub>Bi (2.2 eq., 41.26 g, Aldrich), Cu(OAc)<sub>2</sub> (2.0 eq., 15.48 g, Aldrich), Et<sub>3</sub>N (2.0 eq., 8.62 g) in CH<sub>2</sub>Cl<sub>2</sub> (100 mL) was stirred under N<sub>2</sub> at room temperature for 6 days (monitoring by TLC). The solids were removed by filtration through a plug of Celite rinsing with CH<sub>2</sub>Cl<sub>2</sub>/MeOH (3x75 mL). The filtrate was concentrated, dissolved in hot CH<sub>2</sub>Cl<sub>2</sub>/MeOH (9:1) and filtered through a large plug of silica gel eluting with CH<sub>2</sub>Cl<sub>2</sub>/MeOH (9:1, 2L). The filtrate was concentrated and the residue purified by flash chromatography eluting with straight CH<sub>2</sub>Cl<sub>2</sub> gradient to CH<sub>2</sub>Cl<sub>2</sub>/MeOH (9:1). 2,4-Dioxo-1,5-bis-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine crystallized during concentration of the fractions containing the product, and was isolated by filtration as a white solid.

20                    Step B      Synthesis of 3-Azido-2,4-dioxo-1,5-bis-phenyl-2,3,4,5-tetrahydro-  
1H-1,5-benzodiazepine

                    For this substrate, General Procedure 4-K was modified in the following manner. Initially the product from Step A was suspended (not a solution) in THF at -70°C, and following addition of the KN(TMS)<sub>2</sub> solution, this suspension was allowed to warm to -20°C over a period of 10 minutes, during which the suspension became a solution, and was re-cooled to -70°C; then treated as described in the general procedure. The title compound was purified by trituration with hot CHCl<sub>3</sub>/hexanes (1:1) to yield 3-azido-2,4-dioxo-1,5-bis-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine as a white solid.

-- 156 --

**Step C**      Synthesis of 3-Amino-2,4-dioxo-1,5-bis-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

Following General Procedure 4-L using the product from Step B, 3-amino-2,4-dioxo-1,5-bis-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was prepared as a white solid. Purification was by flash chromatography eluting with CH<sub>2</sub>Cl<sub>2</sub>/MeOH (98:2 gradient to 95:5, with 5% NH<sub>3</sub> in the MeOH).

**Step D**      Synthesis of 3-[N'-(*t*-Butoxycarbonyl)-L-alaninyl]-amino-2,4-dioxo-1,5-bis-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

Following General Procedure I above using N-Boc-L-alanine (Novabiochem) and the product from Step C, 3-[N'-(*t*-butoxycarbonyl)-L-alaninyl]-amino-2,4-dioxo-1,5-bis-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was prepared as a white foam. Purification was by flash chromatography eluting with CH<sub>2</sub>Cl<sub>2</sub>/EtOAc (4:1 gradient to 3:1).

**Step E**      Synthesis of 3-(L-Alaninyl)-amino-2,4-dioxo-1,5-bis-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine Hydrochloride

Following General Procedure 4-N above using the product from Step D, the title compound was prepared as a white amorphous solid.

**EXAMPLE 4-X**

**20**      Synthesis of 3-Amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one

Following the method of R. G. Sherrill et al., *J. Org. Chem.*, **1995**, 60, 730-734, incorporated herein by reference, and using glacial acetic acid and HBr gas, the title compound was prepared.

**25**

-- 157 --

## EXAMPLE 4-Y

Synthesis of 3-(L-Valinyl)-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-oneStep A      Synthesis of 3-[N'-(*tert*-Butylcarbamate)-L-valinyl]-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one

5

(S)-3-Amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one, (1S)-7,7-dimethyl-2-oxobicyclo[2.2.1]heptane-1-methanesulfonate (Example 4-B, Step A) was free based by partitioning between methylene chloride and 1M potassium carbonate. The free amine was then coupled with N-Boc-valine following General Procedure D to give the title compound.

10

$C_{26}H_{32}N_4O_4$  (MW 464.62); mass spectroscopy 464.3.

Anal. Calcd for  $C_{26}H_{32}N_4O_4$ : C, 67.22; H, 6.94; N, 12.06. Found: C, 67.29; H, 6.79; N, 11.20.

Step B      Synthesis of 3-(L-valinyl)-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one

15

Following General Procedure 4-C and using 3-[N'-(*tert*-butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepine-2-one, the title compound was prepared as a white foam.

$C_{21}H_{23}N_4O_2$  (MW 363.48); mass spectroscopy (M+H) 364.2.

20

## EXAMPLE 4-Z

Synthesis of 3-(L-*tert*-Leucinyl)-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-oneStep A      Synthesis of 3-[N'-(*tert*-Butylcarbamate)-L-*tert*-leucinyl]-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one

25

(S)-3-amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one, (1S)-7,7-dimethyl-2-oxobicyclo[2.2.1]heptane-1-methanesulfonate (Example 4-B, Step A) was free based by partitioning between methylene chloride and 1M potassium carbonate. The free amine was then coupled with N-Boc-*tert*-leucine following General Procedure D to give the title compound.

30

$C_{27}H_{35}N_4O_4$  (MW 479.66); mass spectroscopy 479.

-- 158 --

**Step B**      Synthesis of 3-(L-*tert*-Leucinyl)-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one

Following General Procedure 4-C and using 3-[N'-(*tert*-butylcarbamate)-L-*tert*-leucinyl]-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepine-2-one,

5      the title compound was prepared as a white foam.

Anal. Calcd for C<sub>22</sub>H<sub>25</sub>N<sub>4</sub>O<sub>2</sub>•0.5H<sub>2</sub>O: C, 68.19; H, 7.02; N, 14.40.

Found: C, 68.24; H, 7.00; N, 14.00.

**EXAMPLE 4-AA**

10      Synthesis of 3-(L-Alaninyl)-amino-2,3-dihydro-1,5-dimethyl-1H-1,4-benzodiazepine

2,3-Dihydro-1,5-dimethyl-1H-1,4-benzodiazepine was prepared following General Procedures 8-I (using methyl iodide), 4-D and 4-F. Coupling of this intermediate with Boc-L-alanine (Novo) using General Procedure D.

15      The Boc group was removed by dissolving the Boc-protected compound in a 1:1-2:1 mixture of CH<sub>2</sub>Cl<sub>2</sub> and trifluoroacetic acid. The resulting solution was stirred until tlc indicated complete conversion, typically 2 hours. The solution was then stripped to dryness and the residue was taken up in ethyl acetate or CH<sub>2</sub>Cl<sub>2</sub>. The solution was washed with saturated aqueous NaHCO<sub>3</sub> and the  
20      aqueous phase was adjusted to a basic pH, then extracted with ethyl acetate or CH<sub>2</sub>Cl<sub>2</sub>. The organic phase was washed with saturated aqueous NaCl and dried over MgSO<sub>4</sub>. The solution was stripped free of solvent on a rotary evaporator to afford the title compound which was used without further purification.

25      **EXAMPLE 4-AB**

Synthesis of 3-(L-3-Thienylglycinyl)amino-2,4-dioxo-1,5-bis-(2,2-dimethylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

**Step A**      Synthesis of N-(*t*-Butoxycarbonyl)-L-3-thienylglycine

30      N-(*t*-Butoxycarbonyl)-L-3-thienylglycine was prepared from L-α-(3-thienyl)glycine (Sigma) by the procedure described in Bodansky, M. et al; *The*

-- 159 --

*Practice of Peptide Synthesis*; Springer Verlag; 1994, p. 17, incorporated herein by reference.

5      Step B      Synthesis of 3-[N'-(*t*-Butoxycarbonyl)-L-3-thienylglyciny]-amino-2,4-dioxo-1,5-bis-(2,2-dimethylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

Following General Procedure D above using the product from Example 4-V, Step C and the product from Step A above, 3-[N'-(*t*-butoxycarbonyl)-L-3-thienylglyciny]-amino-2,4-dioxo-1,5-bis-(2,2-dimethylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine was prepared.

10      Step C      Synthesis of 3-(L-3-Thienylglyciny)amino-2,4-dioxo-1,5-bis-(2,2-dimethylpropyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

Following General Procedure 4-N above using the product from Step B, the title compound was prepared.

15

#### EXAMPLE 4-AC

Synthesis of 2-(L-Alaniny)-amino-3H-fluoreno[1,9-ef]-2,4-dihydro-1-methyl-1H-1,4-diazepin-3-one

20      Step A      Preparation of 1-[N-( $\alpha$ -isopropylthio)-N'-(benzyloxycarbonyl)-glyciny]-amino-9-fluorenon

A solution of  $\alpha$ -(isopropylthio)-N-(benzyloxycarbonyl)glycine (1 eq; prepared according to Zoller, V. and D. Ben-Ishai *Tetrahedron* 1975, 31, 863, incorporated herein by reference) in dry THF was cooled to 0°C and treated with oxalyl chloride (1 eq.) and 3 drops of DMF. After stirring for 15 minutes at 0°C, the cooling bath was removed and stirring continued at ambient temperature for 40 minutes. The solution was recooled to 0°C. A solution of 1-amino-9-fluorenone (0.9 eq.; Aldrich) and 4-methylmorpholine (2.0 eq., Aldrich) in dry THF was added via cannulation to the acid chloride. The cooling bath was removed and the reaction stirred at ambient for 5 hours. The reaction was diluted with methylene chloride and washed with 0.5 M citric acid, saturated aqueous NaHCO<sub>3</sub>, and brine. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The

25

30

-- 160 --

residue was triturated with 1:1 diethyl ether/hexanes giving the title compound and a yellow solid.

MS Calcd for  $C_{26}H_{25}N_2O_4S$ : 461.15 ( $MH^+$ ), found 461.3

Anal. Calcd for  $C_{26}H_{24}N_2O_4S$ : C, 67.81; H, 5.25; N, 6.08. Found: C, 67.97; H, 5.26; N, 6.14.

**Step B**      Preparation of 1-[N-( $\alpha$ -amino)-N'-(benzyloxycarbonyl)-glyciny]-amino-9-fluorenone

Ammonia gas was bubbled into a solution 1-[N-( $\alpha$ -isopropylthio)-N'-(benzyloxycarbonyl)-glyciny]-amino-9-fluorenone (1 eq) in THF at 0°C. After 35 minutes mercury(II) chloride (1.1 eq) was added. The ice bath was removed and ammonia gas was continued to bubble through the suspension for 4 hours. The bubbler was removed and the reaction continued to stir for 16 hours. The mixture was filtered through celite washing with THF. The filtrate was concentrated *in vacuo*. The crude solid was used in step C without further purification.

**Step C**      Preparation of 2-(benzyloxycarbonyl)-amino-3H-Fluoreno[1,9-ef]-2,4-dihydro-1H-1,4-diazepin-3-one

1-[N-( $\alpha$ -Amino)-N'-(benzyloxycarbonyl)-glyciny]-amino-9-fluorenone (1 eq.) was treated with glacial acetic acid and ammonium acetate (4.7 eq.). The suspension was stirred at ambient temperature for 21 hours. After concentrating the reaction in vacuo, the residue was partitioned between ethyl acetate and 1 N NaOH. The aqueous layer was back-extracted with ethyl acetate. The combined organics were washed with brine, dried over  $Na_2SO_4$ , filtered, and concentrated. The residue was triturated with 3:1 diethyl ether/methylene chloride.

Anal. Calcd for  $C_{23}H_{17}N_3O_3 \cdot 0.25 H_2O$ : C, 71.22; H, 4.55; N, 10.83. Found: C, 71.50; H, 4.44; N, 10.84.

**Step D**      Preparation of 2-(benzyloxycarbonyl)-amino-3H-fluoreno[1,9-ef]-2,4-dihydro-1-methyl-1H-1,4-diazepin-3-one

Following General Procedure 4-A above using 2-(benzyloxycarbonyl)-amino-3H-fluoreno[1,9-ef]-2,4-dihydro-1H-1,4-diazepin-3-one, the title

-- 161 --

intermediate was prepared as a yellow solid.

Anal. Calcd for  $C_{24}H_{19}N_3O_3$ : C, 72.53; H, 4.82; N, 10.57. Found: C, 72.37; H, 5.01; N, 10.36.

5      Step E      Preparation of 2-amino-3H-fluoreno[1,9-ef]-2,4-dihydro-1-methyl-1H-1,4-diazepin-3-one

Following General Procedure 4-B above using 2-(benzyloxycarbonyl)-amino-3H-fluoreno[1,9-ef]-2,4-dihydro-1-methyl-1H-1,4-diazepin-3-one, the title intermediate was prepared as a yellow foam which was used immediately in Step F.

10      Step F      Preparation of 2-[N'-(tert-butylcarbamate)-L-alaninyl]-amino-3H-Fluoreno[1,9-ef]-2,4-dihydro-1-methyl-1H-1,4-diazepin-3-one

Following General Procedure D using N-Boc Alanine (Novabiochem) and 2-amino-3H-fluoreno[1,9-ef]-2,4-dihydro-1-methyl-1H-1,4-diazepin-3-one, the title intermediate was prepared as a yellow solid.

15      MS Calcd for  $C_{24}H_{27}N_4O_4$ : 435.21 ( $MH^+$ ); found 435.29.

Step G      Preparation of 2-(L-alaninyl)-amino-3H-fluoreno[1,9-ef]-2,4-dihydro-1-methyl-1H-1,4-diazepin-3-one

Following General Procedure 4-C using 2-[N'-(tert-butylcarbamate)-L-alaninyl]-amino-3H-fluoreno[1,9-ef]-2,4-dihydro-1-methyl-1H-1,4-diazepin-3-one, the title intermediate was prepared as a yellow foam.

#### EXAMPLE 4-AD

Synthesis of 5-(L-Alaninyl)-amino-7-methyl-1,2,4-triazolo[4,3-d][1,4]benzodiazepin-6(7H)-one

25      Step A      Preparation of 1,3-dihydro-5-(ethylthio)-1-methyl-2H-1,4-benzodiazepin-2-one

A rapidly stirred solution of 1-methyl-1,2,3,4-tetrahydro-3H-1,4-benzodiazepin-2,5-dione (Showell, G. A. et al. *J. Med. Chem.* **1994**, 37, 719, incorporated herein by reference) (1 eq.) and pyridine (1.1 eq., Aldrich) in  
30       $CH_2Cl_2$  (0.16 M in dione), cooled to -40 to -45°C under  $N_2$  in an oven-dried flask, was treated dropwise with trifluoromethanesulfonic anhydride (1.1 eq.,

-- 162 --

Aldrich). The resulting mixture (light yellow color; precipitate) was stirred at -40 to -35°C for 20 minutes and then at 0°C (ice bath) to 10°C for 14.5 hours (note: ice in Dewar melted slowly overnight). The resulting orange solution (some precipitate) was recooled to 0°C and treated dropwise with ethanethiol (1.2 eq., Aldrich). The resulting mixture was stirred at 0 to 4°C under N<sub>2</sub> for 8 hours then partitioned between CH<sub>2</sub>Cl<sub>2</sub> and saturated aqueous NaHCO<sub>3</sub>. The aqueous phase was extracted thrice with CH<sub>2</sub>Cl<sub>2</sub>. The organic extracts were combined, dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated *in vacuo*. The mixture was purified via flash chromatography using a gradient from CH<sub>2</sub>Cl<sub>2</sub> to 30:70 CH<sub>2</sub>Cl<sub>2</sub>/Ethyl acetate as the eluent.

MS Calcd for C<sub>12</sub>H<sub>15</sub>NOS: 235.09 (MH<sup>+</sup>), found 235.0.

Anal. Calcd for C<sub>12</sub>H<sub>14</sub>NOS: C, 61.51; H, 6.02; N, 11.96. Found: C, 61.55; H, 5.99; N, 11.74.

**Step B**      Preparation of 7-methyl-1,2,4-triazolo[4,3-d][1,4]benzodiazepin-6(7H)-one

A mixture of 1,3-dihydro-5-(ethylthio)-1-methyl-2H-1,4-benzodiazepin-2-one (1 eq.) and formic hydrazide (5.8 eq. Aldrich) in n-butanol (0.1 M in benzodiazepine) was stirred at reflux under N<sub>2</sub> for 24 hours. An additional 1.67 eq. of formic hydrazide was added and refluxing continued an additional 16 hours. The yellow solution was evaporated in vacuo and the residue was purified via flash chromatography eluting with a gradient from 98:2 to 96:4 CH<sub>2</sub>Cl<sub>2</sub>/MeOH. The product was obtained as a white solid.

MS Calcd for C<sub>11</sub>H<sub>11</sub>N<sub>4</sub>O: 215.09, found 215.3.

Anal Calcd for C<sub>11</sub>H<sub>10</sub>N<sub>4</sub>O: C, 61.67; H, 4.71; N, 26.15. Found: C, 61.56; H, 4.71; N, 26.08.

**Step C**      Preparation of 5(5H)-azido-7-methyl-1,2,4-triazolo[4,3-d][1,4]benzodiazepin-6(7H)-one



-- 163 --

Following General Procedure 4-K using 5H-7-methyl-1,2,4-triazolo[4,3-d][1,4]benzodiazepin-6(7H)-one, the title compound was prepared as a pale yellow solid.

MS Calcd for C<sub>11</sub>H<sub>9</sub>N<sub>7</sub>O: 255.09, FDMS found 255.0.

5 IR (solution in CHCl<sub>3</sub>) 2138, 2115 cm<sup>-1</sup>.

Step D      Preparation of 5(5H)-amino-7-methyl-1,2,4-triazolo[4,3-d][1,4]benzodiazepin-6(7H)-one

A mixture of 5(5H)-azido-7-methyl-1,2,4-triazolo[4,3-d][1,4]benzodiazepin-6(7H)-one in ethyl acetate (0.1 M) under N<sub>2</sub> was treated  
10 with 10% Pd on carbon (0.4 equiv., Engelhard). The reaction vessel was flushed with H<sub>2</sub> and stirring continued for 3 hours under a balloon of H<sub>2</sub>. The vessel was flushed with N<sub>2</sub> and the contents filtered through celite 545 washing with ethyl acetate. The filtrate was concentrated *in vacuo* to give a white powder.

MS Calcd for C<sub>11</sub>H<sub>12</sub>N<sub>5</sub>O: 230.10 (MH<sup>+</sup>), found 230.1.

15 Step E      Preparation of 5(5H)-[N'-(*tert*-butylcarbamate)-L-alaninyl]-amino-7-methyl-1,2,4-triazolo[4,3-d][1,4]benzodiazepin-6(7H)-one

Following General Procedure D using 5(5H)-amino-7-methyl-1,2,4-triazolo[4,3-d][1,4]benzodiazepin-6(7H)-one and N-Boc Alanine (Novabiochem), the title intermediate was prepared as a white foam.

20 MS Calcd for C<sub>19</sub>H<sub>25</sub>N<sub>6</sub>O<sub>4</sub>: 401.19 (MH<sup>+</sup>), found 401.1.

Anal. Calcd for C<sub>19</sub>H<sub>24</sub>N<sub>6</sub>O<sub>4</sub>: C, 55.74; H, 6.15; N, 20.53. Found: C, 56.06; H, 6.42; N, 20.20.

Step F      Preparation of 5(5H)-(L-alaninyl)-amino-7-methyl-1,2,4-triazolo[4,3-d][1,4]benzodiazepin-6(7H)-one

25 Following General Procedure 4-C using 5(5H)-[N'-(*tert*-butylcarbamate)-L-alaninyl]-amino-7-methyl-1,2,4-triazolo[4,3-d][1,4]benzodiazepin-6(7H)-one, the title compound was prepared as a white foam.

MS Calcd for C<sub>14</sub>H<sub>17</sub>N<sub>6</sub>O<sub>2</sub>: 301.14 (MH<sup>+</sup>), found 301.1.

-- 164 --

**EXAMPLE 4-AE****Synthesis of 3-(L-Alaninyl)-amino-2,3-dihydro-1-methyl-5-piperidinyl-1H-1,4-benzodiazepin-2-one****Step A Preparation of 3-[N'-tert-butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-1-methyl-5-(1-piperidinyl)-1H-1,4-benzodiazepin-2-one**

Following General Procedure D above using 3-amino-1,3-dihydro-1-methyl-5-(1-piperidinyl)-2H-1,4-benzodiazepin-2-one (Example 4-A) and N-Boc Alanine (Novabiochem), the title compound was prepared as a white foam.

MS Calcd for  $C_{23}H_{33}N_5O_4$  444.26 ( $MH^+$ ), found 444.4.

Anal. Calcd for  $C_{23}H_{33}N_5O_4 \cdot 0.5 H_2O$ : C, 61.04; H, 7.57; N, 15.47.

Found: C, 61.09; H, 7.29; N, 15.21.

**Step B Preparation of 3-(L-alaninyl)-amino-2,3-dihydro-1-methyl-5-(1-piperidinyl)-1H-1,4-benzodiazepin-2-one**

Following General Procedure 4-C using 3-[N'-tert-butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-1-methyl-5-(1-piperidinyl)-1H-1,4-benzodiazepin-2-one, the title compound was prepared.

**EXAMPLE 4-AF****Synthesis of 3-(L-Alaninyl)-amino-2,3-dihydro-5-isopropyl-1-methyl-1H-1,4-benzodiazepin-2-one****Step A Preparation of 3-(benzyloxycarbonyl)-amino-2,3-dihydro-5-isopropyl-1H-1,4-benzodiazepin-2-one**

A slurry of 2-(benzotriazol-1-yl)-N-(benzyloxycarbonyl)glycine (1.1 equiv.; Katritzky, A. R. et al. *J. Org. Chem.* **1990**, 55, 2206, incorporated herein by reference) in THF (0.3 M) was cooled to 0°C and treated with oxalyl chloride (1.1 equiv.) in a dropwise manner. To the slurry was added dropwise DMF (0.1 equiv.); stirring was continued at 0 °C for 1 hour. A solution of 1-(2-aminophenyl)-2-methyl-1-propanone (1.0 equiv.; Robl, J. A. *Synthesis* **1991**, 56, incorporated herein by reference) and N-methylmorpholine (2.2 equiv.) in THF (1 M in propanone), pre-cooled to 0 °C, was added via cannula. Upon completion of the addition, the reaction was warmed to ambient temperature. The mixture was filtered, washing the filter cake with THF. The filtrate was transferred to a

-- 165 --

three-neck flask and treated with ammonia gas through a dispersion tube for 15 minutes. Methanol (0.3 M in propanone) was added and the ammonia continued to be bubbled through the solution for 1 hour. The reaction was concentrated, diluted with ethyl acetate, and re-concentrated; this was repeated again. The  
5 residue was diluted with ethyl acetate and washed twice with 1 N NaOH. The aqueous washes were back-extracted with ethyl acetate. The combined organics were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Acetic acid was added and concentrated *in vacuo*. The syrup was dissolved in acetic acid (0.6 M in propanone) and treated with ammonium acetate (4.0 equiv.). The  
10 reaction was stirred at ambient temperature for 18 hours. The resultant solid was filtered, washing with H<sub>2</sub>O.

MS Calcd for C<sub>20</sub>H<sub>22</sub>N<sub>3</sub>O<sub>3</sub>: 352.17, found 352.5.

**Step B**      Preparation of 3-(benzyloxycarbonyl)-amino-2,3-dihydro-5-isopropyl-1-methyl-1H-1,4-benzodiazepin-2-one

15      Following General Procedure 4-A using 3-(benzyloxycarbonyl)-amino-2,3-dihydro-5-isopropyl-1H-1,4-benzodiazepin-2-one, the title compound was prepared as a white solid.

MS Calcd for C<sub>21</sub>H<sub>24</sub>N<sub>3</sub>O<sub>3</sub>: 366.18 (MH<sup>+</sup>), found 366.2.

Anal. Calcd for C<sub>21</sub>H<sub>23</sub>N<sub>3</sub>O<sub>3</sub>•0.25H<sub>2</sub>O: C, 68.18; H, 6.40; N, 11.36.

20      Found: C, 68.36; H, 6.28; N, 11.48.

**Step C**      Preparation of 3-amino-2,3-dihydro-5-isopropyl-1-methyl-1H-1,4-benzodiazepin-2-one

Following General Procedure 4-B using 3-(benzyloxycarbonyl)-amino-2,3-dihydro-5-isopropyl-1-methyl-1H-1,4-benzodiazepin-2-one, the title compound  
25 was prepared as a white foam.

MS Calcd for C<sub>13</sub>H<sub>18</sub>N<sub>3</sub>O: 232.14 (MH<sup>+</sup>), found 232.19.

-- 166 --

**Step D**      Preparation of 3-[N'-(*tert*-butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-5-isopropyl-1-methyl-1H-1,4-benzodiazepin-2-one

Following General Procedure D using 3-amino-2,3-dihydro-5-isopropyl-1-methyl-1H-1,4-benzodiazepin-2-one and N-Boc alanine (Novabiochem), the title  
5 compound was prepared as a white solid.

MS Calcd for  $C_{21}H_{31}N_4O_4$ : 403.23 (MH<sup>+</sup>), found 403.46.

**Step E**      Preparation of 3-(L-alaninyl)-amino-2,3-dihydro-5-isopropyl-1-methyl-1H-1,4-benzodiazepin-2-one

Following General Procedure 4-C using 3-[N'-(*tert*-butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-5-isopropyl-1-methyl-1H-1,4-benzodiazepin-2-one,  
10 the title compound was prepared as a white foam.

MS Calcd for  $C_{16}H_{23}N_4O_2$ : 303.18 (MH<sup>+</sup>), found 303.21.

Anal Calcd for  $C_{16}H_{22}N_4O_2 \cdot 0.3 H_2O$ : C, 62.44; H, 7.40; N, 18.20.

Found: C, 62.58; H, 7.10; N, 17.79.

15

**EXAMPLE 4-AG**

Synthesis of 3-(L-Alaninyl)-amino-2,3-dihydro-5-*n*-propyl-1-methyl-1H-1,4-benzodiazepin-2-one

**Step A**      Preparation of 1-(2-aminophenyl)-1-butanone

20 A solution of anthranilonitrile (1 equiv., Aldrich) in diethyl ether (2.4 M) was cooled to 0°C and treated with propylmagnesium chloride (2.5 equiv., Aldrich; 2.0 M in Et<sub>2</sub>O) in a dropwise manner over the course of an hour. After addition of 25% of the Grignard reagent, an additional 1/10 volume of Et<sub>2</sub>O was added. The cooling bath was removed and stirring of the suspension continued for  
25 5 hours. The reaction was returned to 0 °C and cautiously quenched with 3 N HCl. The cooling bath was removed and stirring continued for 30 minutes. The mixture was made basic by the addition of solid NaOH. The contents were extracted thrice with ethyl acetate; brine was added to help break up the suspension. The combined extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>,

-- 167 --

filtered, and concentrated. The crude residue was chromatographed eluting with 95:5 hexanes/ethyl acetate.

MS Calcd for  $C_{10}H_{13}NO$ : 163.10, found 163.18.

Anal. Calcd for  $C_{10}H_{13}NO \cdot 0.2 H_2O$ : C, 72.00; H, 8.10; N, 8.40.

5 Found: C, 72.36; H, 8.25; N, 8.76

**Step B**      Preparation of 3-(benzyloxycarbonyl)-amino-2,3-dihydro-5-isopropyl-1H-1,4-benzodiazepin-2-one

A slurry of 2-(benzotriazol-1-yl)-N-(benzyloxycarbonyl)glycine (1.1 equiv.; Katritzky, A. R. et al. *J. Org. Chem.* 1990, 55, 2206, incorporated herein  
10 by reference) in THF (0.3 M) was cooled to 0 °C and treated with oxalyl chloride (1.1 equiv.) in a dropwise manner. To the slurry was added dropwise DMF (0.1 equiv.); stirring was continued at 0 °C for 1 hour. A solution of 1-(2-aminophenyl)-1-butanone and N-methylmorpholine (2.2 equiv.) in THF (1 M in butanone), pre-cooled to 0 °C, was added via cannula. Upon completion of the  
15 addition, the reaction was warmed to ambient temperature. The mixture was filtered, washing the filter cake with THF. The filtrate was transferred to a three-neck flask and treated with ammonia gas through a dispersion tube for 15 minutes. Methanol (0.3 M in butanone) was added and the ammonia continued to be bubbled through the solution for 1 hour. The reaction was concentrated, diluted  
20 with ethyl acetate, and re-concentrated; this was repeated again. The residue was diluted with ethyl acetate and washed twice with 1 N NaOH. The aqueous washes were back-extracted with ethyl acetate. The combined organics were washed with brine, dried over  $Na_2SO_4$ , filtered, and concentrated. Acetic acid was added and concentrated *in vacuo*. The syrup was dissolved in acetic acid (0.6 M in  
25 butanone) and treated with ammonium acetate (4.0 equiv.). The reaction was stirred at ambient temperature for 18 hours. The resultant solid was filtered, washing with  $H_2O$ .

MS Calcd for  $C_{20}H_{22}N_3O_3$ : 352.17, found 352.4.

-- 168 --

Step C      Preparation of 3-(benzyloxycarbonyl)-amino-2,3-dihydro-5-n-propyl-1-methyl-1H-1,4-benzodiazepin-2-one

Following General Procedure 4-A using 3-(benzyloxycarbonyl)-amino-2,3-dihydro-5-n-propyl-1H-1,4-benzodiazepin-2-one, the title compound was prepared  
5 as a white solid.

MS Calcd for  $C_{21}H_{24}N_3O_3$ : 366.18 (MH<sup>+</sup>), found 366.2.

Step D      Preparation of 3-amino-2,3-dihydro-5-n-propyl-1-methyl-1H-1,4-benzodiazepin-2-one

Following General Procedure 4-B using 3-(benzyloxycarbonyl)-amino-2,3-dihydro-5-n-propyl-1-methyl-1H-1,4-benzodiazepin-2-one, the title compound was  
10 prepared as a white foam. This compound was used immediately in Step E.

Step E      Preparation of 3-[N'-(tert-butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-5-n-propyl-1-methyl-1H-1,4-benzodiazepin-2-one

Following General Procedure D using 3-amino-2,3-dihydro-5-n-propyl-1-methyl-1H-1,4-benzodiazepin-2-one and N-Boc alanine (Novabiochem), the title  
15 compound was prepared as a white solid.

MS Calcd for  $C_{21}H_{31}N_4O_4$ : 403.23 (MH<sup>+</sup>), found 403.4.

Step F      Preparation of 3-(L-alaninyl)-amino-2,3-dihydro-5-n-propyl-1-methyl-1H-1,4-benzodiazepin-2-one

Following General Procedure 4-C using 3-[N'-(tert-butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-5-n-propyl-1-methyl-1H-1,4-benzodiazepin-2-one the  
20 title compound was prepared as a white foam.

**EXAMPLE 4-AH**

25      Synthesis of 3-(L-Alaninyl)amino-4-n-butyl-3,4-dihydro-1-methyl-1H-1,4-benzodiazepin-2,5-dione

Step A      Preparation of 4-n-butyl-3,4-dihydro-1-methyl-1H-1,4-benzodiazepin-2,5-dione

A solution of 13 mmol of 3,4-dihydro-1-methyl-1H-1,4-benzodiazepin-2,5-dione (Tett. Lett. 1994, 50(30), 9051, incorporated herein by reference) in 30mL  
30 dry dimethylformamide is treated, dropwise at 0°C under nitrogen cover, with one equivalent of potassium-t-butoxide (Aldrich; 1.0 M in THF). After forty-five

-- 169 --

minutes at 0°C, iodobutane is introduced via syringe over several minutes. The reaction mixture is stirred at ambient temperature seventy-five minutes, diluted with methylene chloride, and then washed with water and saturated sodium chloride. The organic solution is dried over anhydrous sodium sulfate, filtered  
5 and concentrated in vacuo to give a yellow oil. Flash column chromatography (silica gel; ethyl acetate/hexane (7/1) eluent) provides pure material as a colorless oil (90% yield).

$C_{14}H_{18}N_2O_2$  (MW = 246.3)

Anal. Calcd for  $C_{14}H_{18}N_2O_2$ : C, 68.27; H, 7.37; N, 11.37. Found: C,  
10 68.53; H, 7.11; N, 11.41.

**Step B**      Preparation of 3-azido-4-n-butyl-3,4-dihydro-1-methyl-1H-1,4-benzodiazepine-2,5-dione

Following General Procedure 4-K using 4-n-butyl-3,4-dihydro-1-methyl-1H-1,4-benzodiazepin-2,5-dione the title intermediate was prepared as a waxy  
15 solid.

$C_{14}H_{17}N_5O_2$  (MW=287.3); Exact Mass FAB+ Theory 288.1461  
Found 288.1459

**Step C**      Preparation of 3-amino-4-n-butyl-3,4-dihydro-1-methyl-1H-1,4-benzodiazepine-2,5-dione

20 A solution of 0.4 mmol of azide (see Step B above) in ethyl acetate is treated with 170mg of 10% Pd/C (Englehard) and hydrogenated overnight via the static pressure of a hydrogen-filled balloon attached to the reaction flask via a syringe and septum. The catalyst is removed by filtration and the filtrate

-- 170 --

concentrated in vacuo to give a yellow oil. This was purified via chromatography (silica gel; 1 mil Chromatotron plate; 95/5 methylene chloride/methanol[7N ammonia])

Anal. Calcd for  $C_{14}H_{19}N_3O_2$ : C, 64.35; H, 7.33; N, 16.08. Found: C, 64.58; H, 7.19; N, 15.94.

**Step D**      Preparation of 3-(N-tert-butylcarbamate-L-alaninyl)amino-4-n-butyl-3,4-dihydro-1-methyl-1H-1,4-benzodiazepin-2,5-dione

Following General Procedure D using N-Boc-alanine and the aminobenzodiazepinedione from Step C above, the title intermediate was obtained as an amorphous white solid.

Anal. Calcd. for  $C_{22}H_{32}N_4O_5$ : C, 61.09; H, 7.46; N, 12.95. Found: C, 60.83; H, 7.51; N, 12.69

**Step E**      Preparation of 3-(L-alaninyl)amino-4-n-butyl-3,4-dihydro-1-methyl-1,4-benzodiazepin-2,5-dione

Following General Procedure 4-C using the Boc-protected intermediate from Step D above, the title intermediate is obtained as a white solid.

$C_{17}H_{24}N_4O_3$  (MW=332.4) Exact Mass FAB+ Theory MW= 333.1927  
Found MW=333.1924

**EXAMPLE 4-AI**

Synthesis of 3-(L-alaninyl)amino-1,3-dihydro-5-ethylthio-1-methyl-2H-1,4-benzodiazepin-2-one

**Step A**      Preparation of 1,3-dihydro-5-ethylthio-1-methyl-2H-1,4-benzodiazepin-2-one

A solution of 1.0 mmol of 3,4-dihydro-1-methyl-1H-1,4-benzodiazepin-2,5-dione (Tett. Lett. 1994, 50(30), 9051) in 8mL dry dichloromethane (Aldrich Sure Seal), in an oven-dried round bottom flask under nitrogen cover, is treated with 1.1 equivalents of anhydrous pyridine (Mallinkrodt). The reaction is cooled to  $-51^{\circ}\text{C}$  in a dry ice/acetone bath and treated, dropwise via syringe over three minutes, with 1.1 equivalents of trifluoromethanesulfonic anhydride (Aldrich; sealed ampules). The reaction mixture is allowed to stir twenty minutes while



-- 171 --

maintaining a reaction temperature between -47 and -35°C. The temperature is brought to 0°C over 1-2 minutes and then maintained at that temperature for one hour. Ethanethiol (3.4 equivalents) is introduced via syringe and the mixture allowed to stir overnight, still immersed in an ice water-filled Dewar. The temperature had risen to 16°C by morning. The reaction mixture is partitioned between methylene chloride and aqueous sodium bicarbonate. The aqueous portion is further extracted three times with methylene chloride and the combined extracts dried over anhydrous sodium sulfate, filtered and concentrated in vacuo to give a yellow oil. Purification is effected by flash chromatography (silica gel; gradient elution using methylene chloride/ethyl acetate in 100/1, 95/5, 90/10 and 85/15. The title compound is obtained as a colorless oil which crystallizes on standing.

Anal. Calcd for  $C_{12}H_{14}N_2OS$ : C, 61.51; H, 6.02; N, 11.96. Found: C, 61.55; H, 5.99; N, 11.74.

**Step B**      Preparation of 3-azido-1,3-dihydro-5-ethylthio-1-methyl-2H-1,4-benzodiazepin-2-one

Following General Procedure 4-K using the ethylthiobenzodiazepine intermediate from Step A above, the title intermediate was prepared as a white solid.

Anal. Calcd. for  $C_{12}H_{13}N_5OS$ : C, 52.35; H, 4.76; N, 25.44. Found: C, 52.63; H, 4.67; N, 25.39.

**Step C**      Preparation of 3-amino-1,3-dihydro-5-ethylthio-1-methyl-2H-1,4-benzodiazepin-2-one

A solution of 1.90 mmol of azide (see Step B above) in 13mL of tetrahydrofuran to which had been added 1 mL of water is treated with an excess of triphenylphosphine (2.8 equivalents) added in one portion as a solid. The reaction is stirred at room temperature under nitrogen cover for twenty hours and then diluted with ethyl acetate. The solution is extracted with 1N HCl three times, and the combined extracts are rendered basic by the addition of 5N NaOH. This is extracted three times with ethyl acetate and the combined extracts washed once

-- 172 --

with saturated aqueous sodium chloride, dried over anhydrous sodium sulfate, filtered and concentrated in vacuo on a rotary evaporator at 20°C. The resulting oil is used immediately.

$C_{12}H_{15}N_3OS$  (MW=249.3) FAB+ Exact Mass: Theory 250.1014

5 Found 250.1011

Step D Preparation of 3-(N-tert-butylcarbamate-L-alaninyl)amino-1,3-dihydro-5-ethylthio-1-methyl-2H-1,4-benzodiazepin-2-one

Following General Procedure D using N-Boc-alanine and the aminobenzodiazepin-2-one from Step C above, the title intermediate was obtained  
10 as an amorphous white solid.

Anal. Calcd. for  $C_{20}H_{28}N_4O_4S$ : C, 57.12; H, 6.71; N, 13.32. Found: C, 56.85; H, 6.77; N, 13.12

Step E Preparation of 3-(L-alaninyl)amino-4-n-butyl-3,4-dihydro-1-methyl-1,4-benzodiazepin-2,5-dione

15 Following General Procedure 4-C using the Boc-protected intermediate from Step D above, the title intermediate is obtained as a colorless oil.

$C_{15}H_{20}N_4O_2S$  (MW=320.4) FAB+ Exact Mass Theory 321.1385  
Found 321.1388.

20 **EXAMPLE 4-AJ**

Synthesis of 3-L-Alaninyl-amino-5-phenyl-1-methyl-2H-1,5-diazepin-2-one

Step A Preparation of 1-phenyl-4-piperidinone

The title intermediate was prepared from aniline (Aldrich) and methyl acrylate (Aldrich) by the literature procedure of Hermant, R. M., et al. *J. Am. Chem. Soc.*, 1990, 112, 1214-1221, incorporated herein by reference.  
25

Step B Preparation of 5-phenyl-2H-1,5-diazepin-2-one

To the product from Step A (12.54 g, 71.56 mmols) in glacial acetic acid (60 mL) and concentrated sulfuric acid (30 mL) at 0 °C was added sodium azide (5.12 g, 78.7 mmols) in five portions of 1.024 g over a period of 4 hours. The  
30 resultant pale yellow mixture was allowed to warm to room temperature with

-- 173 --

stirring under nitrogen for 17 hours. The mixture was poured onto ice and neutralized to pH 7 with 5 M aq. NaOH. The product was extracted into methylene chloride (2x400 mL) and the combined organic extracts were dried over sodium sulfate, filtered, and concentrated to a pale yellow solid. Flash chromatography purification eluting with EtOAc gradient to EtOAc/MeOH (95:5) yielded 12.37g (91%) of the title intermediate as a pale yellow solid.

$C_{11}H_{14}N_2O$  (MW. 190.25), mass spectroscopy ( $MH^+$ ), 191.4.

Anal. Calcd. For  $C_{11}H_{14}N_2O$ : C, 69.45; H, 7.42; N, 14.72, Found: C, 69.74; H, 7.23; N, 15.00.

10 Step C      Preparation of 5-phenyl-1-methyl-2H-1,5-diazepin-2-one

Following General Procedure 4-G using the product from Step B, methyl iodide, and potassium *tert*-butoxide, the title intermediate was prepared. HPLC purification eluting with EtOAc yielded the product as a white solid.

$C_{12}H_{16}N_2O$  (MW. 204.27), mass spectroscopy ( $MH^+$ ), 205.2.

15      Anal. Calcd. For  $C_{12}H_{16}N_2O$ : C, 70.56; H, 7.90; N, 13.71, Found: C, 70.65; H, 7.70; N, 13.95.

Step D      Preparation of 3-azido-5-phenyl-1-methyl-2H-1,5-diazepin-2-one

Following the General Procedure 4-E using the product from Step C, the title intermediate was prepared. HPLC purification eluting with EtOAc/hexanes (60:40), and a second HPLC purification eluting with methylene chloride/methanol (98:2), yielded the product as a light yellow solid.

$C_{12}H_{15}N_3O$  (MW. 245), mass spectroscopy ( $MH^+$ ), 246.4.

Step E      Preparation of 3-amino-5-phenyl-1-methyl-2H-1,5-diazepin-2-one

Following the General Procedure 4-F using the product from Step D, the title intermediate was prepared as yellow oil which upon standing solidified.

$C_{12}H_{17}N_3O$  (MW. 219), mass spectroscopy ( $MH^+$ ), 220.3.

25      Anal. Calcd. For  $C_{12}H_{17}N_3O$ : C, 65.73; H, 7.81; N, 19.16, Found: C, 65.94; H, 7.37; N, 18.85.

-- 174 --

**Step F**      Preparation of 3-[N'-(*t*-butoxycarbonyl)-L-alaninyl]-amino-5-phenyl-1-methyl-2H-1,5-diazepin-2-one

Following General Procedure D using the product from Step E and Boc-L-Alanine (Nova Biochem) the title intermediate was prepared. HPLC purification eluting with EtOAc/hexanes (1:1) afforded the title compound as a white solid.

Exact Mass, anal. calcd. for  $C_{20}H_{31}N_4O_4$ : Theory, 391.2345, Found, 391.2342.

**Step G**      Preparation of 3-(L-alaninyl)-amino-5-phenyl-1-methyl-2H-1,5-diazepin-2-one

Following the General Procedure 4-N using the product from Step F the title intermediate was prepared. HPLC purification eluting with 95:5 methylene chloride/methanol gave the title intermediate.

Exact mass anal. calcd. Form.  $C_{15}H_{23}N_4O_2$ , Theory, 291.1821; Found, 291.1816.

**EXAMPLE 4-AK**

Synthesis of 3-(S)-Phenylglycinyl]-amino-5-phenyl-1-methyl-2H-2,3,4,5-tetrahydro-1,5-benzodiazepine-2-one

**Step A**      Preparation of 3-[N'-(*t*-Butoxycarbonyl)-(S)-phenylglycinyl]-amino-1-methyl-5-phenyl-1,3,4,5-tetrahydro-2H-1,5-benzodiazepine 2-one

Following modified General Procedure J using the product from Example 4-M, Step B, and Boc-L-Phenylglycine (Nova Biochem), the title intermediate was prepared. The modification was that the reaction was only stirred for 6 hours. HPLC purification eluting with 80/20 hexanes/EtOAc afforded the separated diastereomers; isomer 1 (first eluting) and isomer 2 (second eluting).

$C_{29}H_{32}N_4O_4$  (MW 500.60); mass spectroscopy for isomer 1:  $MH^+$  501.2;  $MH^-$ , 499.3; mass spectroscopy for isomer 2:  $MH^+$  501.2;  $MH^-$ , 499.3.

**Step B**      Synthesis of 3-((S)-phenylglycinyl)-amino-1-methyl-5-phenyl-1,3,4,5-tetrahydro-2H-1,5-benzodiazepine 2-one

Following the General Procedure 4-N using the products (Isomers 1 and 2 brought through reaction sequence separately) from Step A, the title intermediates

-- 175 --

were prepared. HPLC purification eluting with 95/5 methylene chloride/methanol afforded the title intermediates as light yellow oils.

$C_{24}H_{24}N_4O_2$  (MW 400.48); mass spectroscopy for isomer 1:  $MH^+$ , 401.3;  $MH^-$ , 399.2; mass spectroscopy for isomer 2:  $MH^+$ , 401.2;  $MH^-$ , 399.3.

5

#### EXAMPLE 4-AL

##### Synthesis of 3-(L-Norvalinyl)-amino-2,4-dioxo-1,5-bis-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

##### 10 Step A      Preparation of 3-[N'-(*t*-Butoxycarbonyl)-L-norvalinyl]-amino-2,4-dioxo-1,5-bis-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

Following General Procedure D using the product from Example 4-R, Step C and Boc-L-norvaline (BACHEM) the title intermediate was prepared. HPLC purification eluting with EtOAc/Hexanes (60:40) afforded the title intermediate as a white solid.

15  $C_{21}H_{30}N_4O_5$  (MW 418.49).

Anal. Calcd. for  $C_{21}H_{30}N_4O_5$  hemihydrate: C, 59.00; H, 7.39; N, 13.10.

Found: C, 59.35; H, 7.58; N, 12.86.

Exact Mass calcd. for  $C_{21}H_{31}N_4O_5$ : Theory 419.2294, Found 419.2289.

##### 20 Step B      Preparation of 3-(L-norvalinyl)-amino-2,4-dioxo-1,5-bis-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

Following General Procedure 4-N using the product from Step A, the title intermediate was prepared and used without further purification.

$C_{16}H_{22}N_4O_3$  (MW 318.38)

Exact Mass calcd. for  $C_{16}H_{23}N_4O_3$ : Theory 319.1770, Found 319.1774.

25

-- 176 --

**EXAMPLE 4-AM****Synthesis of 3-(L-norvalinyl)-amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine****5      Step A      Preparation of 3-[N'-(*t*-Butoxycarbonyl)-L-norvalinyl]-amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine**

Following General Procedure D using 3-amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine (CAS No. 131604-75-6) and Boc-L-norvaline (BACHEM) the title intermediate was prepared. HPLC purification with EtOAc/hexanes (1:1) afforded the title intermediate as a white  
10      solid.

$C_{26}H_{32}N_4O_5$  (MW 480.57); mass spectroscopy,  $MH^+$  481.2.

Anal. Calcd. for  $C_{26}H_{32}N_4O_5$ : C, 64.98; H, 6.71; N, 11.66. Found: C, 64.87; H, 6.83; N, 11.53.

**15      Step B      Preparation of 3-(L-norvalinyl)-amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine**

Following General Procedure 4-N using the product from Step A, the title intermediate was prepared. HPLC purification eluting with methylene chloride/methanol (9:1) afforded the title intermediate as an off white solid.

$C_{21}H_{24}N_4O_3$  (MW 380.45); mass spectroscopy,  $MH^+$  381.1.

20      Exact Mass calcd. for  $C_{21}H_{25}N_4O_3$ : Theory 381.1889, Found 381.1928.

**EXAMPLE 4-AN****Synthesis of 3-(L-alaninyl)-amino-1,5-bis-methyl-2H-2,3,4,5-tetrahydro-1,5-benzodiazepine-2-one****25      Step A      Preparation of N,N'-Dimethyl-1,2-phenylenediamine**

Following the literature procedures of Stetter, H., *Chem. Ber.*, 1953, 86, 161 and Cheeseman, G. W. H., *J. Chem. Soc.*, 1955, 3308, incorporated herein

-- 177 --

by reference, the title intermediate was prepared from 1,2-phenylenediamine (Aldrich) as a low melting solid.

$C_8H_{12}N_2$  (MW 136); mass spectroscopy, 136.1.

Exact mass anal. Calcd. for  $C_8H_{13}N_2$ : Theory, 137.1079, Found, 137.081

5    **Step B**        Preparation of 1,5-Bis-methyl-2,3,4,5-tetrahydro-2H-1,5-benzadiazepine-2-one

To a solution of the product from Step A (400 mg, 2.94 mmols) in 5M Aq. HCl (30 mL) was added acrylic acid (0.202 mL, 3.23 mmols, Aldrich) and the mixture heated to reflux for 18 hours. The black mixture was allowed to cool,  
10 then poured onto ice and the pH adjusted to 10 with 5M Aq. NaOH. The product was extracted into  $CH_2Cl_2$  (200 mL) and washed with water (100 mL) and brine (100 mL). The organic phase was dried over  $Na_2SO_4$ , filtered, and concentrated to give a black oil. HPLC purification eluting with hexanes/EtOAc (1:1) afforded 364 mg of the title intermediate as a brown oil.

15         $C_{11}H_{14}N_2O$  (MW 190.25); mass spectroscopy,  $MH^+$  191.4

Anal. Calcd. for  $C_{11}H_{14}N_2O$ : Theory, C, 69.45; H, 7.42; N, 14.72;  
Found, C, 69.26; H, 7.40; N, 14.64.

**Step C**        Preparation of 3-Azido-1,5-bis-methyl-2,3,4,5-tetrahydro-2H-1,5-benzadiazepine-2-one

20        Following General Procedure 4-E using the product from Step B, the title intermediate was prepared. HPLC purification eluting with hexanes/EtOAc (7:3) afforded the title intermediate as a light brown oil.

$C_{11}H_{13}N_5O$  (MW 231.26), mass spectroscopy,  $MH^+$  232.2

Exact mass Anal. Calcd. for  $C_{11}H_{14}N_5O$ : Theory, 232.1198; Found,  
25    232.1196.

**Step D**        Preparation of 3-Amino-1,5-bis-methyl-2,3,4,5-tetrahydro-2H-1,5-benzadiazepine-2-one

Following procedure 4-F using the product from Step C, the title intermediate was prepared and used without chromatographic purification.

30         $C_{11}H_{15}N_3O$  (MW 205.26), mass spectroscopy  $MH^+$  206.2.

-- 178 --

Exact mass Anal. Calcd. for  $C_{11}H_{16}N_3O$ : Theory, 206.1293; Found, 206.1295.

**Step E**      Preparation of 3-[N'-(*t*-Butoxycarbonyl)-L-alaninyl]-amino-1,5-bis-methyl-2,3,4,5-tetrahydro-2H-1,5-benzodiazepine-2-one

5            Following General Procedure D using the product from Step D and Boc-L-Alanine (Nova Biochem), the title intermediate was prepared. HPLC purification eluting with EtOAc/hexanes (6:4) afforded the title intermediate as a white foamy solid.

$C_{19}H_{28}N_4O_4$  (MW 376.45), mass spectroscopy  $MH^+$ , 377.4,  $MH^-$ ,  
10    375.3.

Anal. Calcd. for  $C_{19}H_{28}N_4O_4$ : Theory, C, 60.62; H, 7.50; N, 14.88.  
Found, C, 60.68; H, 7.42; N, 14.38.

**Step F**      Preparation of 3-(L-Alaninyl)-amino-1,5-bis-methyl-2,3,4,5-tetrahydro-2H-1,5-benzodiazepine-2-one

15            Following the procedure 4-N using the product from Step E, the title intermediate was prepared. HPLC purification eluting with EtOAc/hexanes (6:4) afforded the title intermediate as a thick yellow oil.

$C_{14}H_{20}N_4O_2$ : (MW 276.34), mass spectroscopy,  $MH^+$ , 277.2.

Using the following procedures, the following additional intermediates can  
20    be prepared for use in this invention.

### GENERAL PROCEDURE C-H

The intermediates shown in Table C-A were synthesized in parallel in using the following procedure:

25    **Step A**

To a solution of 3-(tert-butoxycarbonyl)amino-2,3-dihydro-5-phenyl-1H-1,4-benzodiazepin-2-one (CA No. 125:33692: 100 mg, 0.28 mmol) in 1 mL of anhydrous DMF was added 600  $\mu$ L of a solution of 0.5 M potassium bis(trimethylsilyl)amide (0.30 mmol) in toluene. Neat alkyl halide (0.56 mmol; as  
30    indicated in Table C-A) was added immediately in one portion and the reaction



-- 179 --

mixture was left undisturbed overnight. When an alkyl chloride was used, 1 equivalent of sodium iodide was added to the reaction mixture. After concentration under reduced pressure, the crude reaction residue was partitioned between methylene chloride (2 mL) and aqueous saturated bicarbonate (2 mL) and then passed through a 5 g Extralut QE cartridge (EM Science; Gibbstown, NJ) using 10 mL of methylene chloride. The resulting filtrate was concentrated under reduced pressure and the crude product was further purified using automated semi-preparative HPLC (YMC 20 X 50 mm Silica column; gradient elution; 0-5 % (5.5 min.), 5-20 % (3.5 min.), 20-100 % (2 min.), 100% (4 min.) ethyl acetate/methylene chloride, flow rate of 25 mL/min.). Product provided the expected M + 1 peak by IEX MS and was carried on without further purification and characterization.

#### Step B

The product obtained from Step A was dissolved in 5 mL of a 15 % TFA/methylene chloride solution and allowed to stand undisturbed for 16 h. After concentration under reduced pressure, the TFA salt was dissolved in methanol and loaded directly onto a 1 g SCX column. The column was washed 3 X with 2 mL portions of methanol and the product was eluted from the column using 6 mL of 2.0 M solution of ammonia/methanol. After concentration under reduced pressure, the product were characterized by IEX MS and carried on without further purification.

#### Step C

To the crude product obtained from Step B (1.05 equiv.) was added sequentially a 0.3 mM stock solution of HOBt•H<sub>2</sub>O (1.05 equiv.) in DMF, a 0.3 mM stock solution of N-t-BOC-L-alanine (1.0 equiv.) in THF and 0.3 mM stock solution of 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide (1.05 equiv.) in THF. After standing undisturbed for 24 h, the reaction mixture was concentrated and the residue redissolved in 2 mL of a 10% methanol/methylene chloride solution. This solution was then filtered through a pre-washed (methanol) 1 g SCX (Varian

-- 180 --

Sample Preparation) column using an additional 8 mL of the same solvent. For Example C-V a 1 g Si column (Varian Sample Preparation) was used. The filtrate was concentrated under a stream of nitrogen to approximately 1/3 its original volume and then passed over a plug (500 mg) of AG 1-8x anion exchange resin (BioRad; Hercules, California; columns were pre-washed with 1N NaOH, water and methanol) using an additional 10 mL of methanol. The resulting filtrate was concentrated under reduced pressure and the crude product was carried on without further purification after characterization by IEX MS.

#### Step D

The crude product obtained from Step C was dissolved in 5 mL of a 15 % TFA/methylene chloride solution and allowed to stand undisturbed for 16 h. After concentration under reduced pressure, the TFA salt was dissolved in methanol and loaded directly onto a 1 g SCX column. The column was washed 3 X with 2 mL portions of methanol and the product was eluted from the column using 6 mL of 2.0 M solution of ammonia/methanol. After concentration under reduced pressure, the product was characterized by IEX MS and carried on without further purification. The intermediates prepared by this method are shown in Table C-A.

TABLE C-A  
Intermediates

Ex.	Alkyl Halide	Intermediate	MS
C-A	3-Fluorobenzyl bromide (Aldrich)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(3-fluorobenzyl)-1H-1,4-benzodiazepin-2-one	431.1
C-B	Benzyl bromide (Aldrich)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(benzyl)-1H-1,4-benzodiazepin-2-one	513.2
C-C	<i>tert</i> -Butylbenzyl bromide (Aldrich)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(4- <i>tert</i> -butylbenzyl)-1H-1,4-benzodiazepin-2-one	469.2
C-D	2-Bromoethylcyclohexane (Fairfield)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(2-cyclohexylethyl)-1H-1,4-benzodiazepin-2-one	433.2

-- 181 --

Ex.	Alkyl Halide	Intermediate	MS
C-E	1-Bromo-3,3-dimethylbutane (Wiley)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(3,3-dimethylbutyl)-1H-1,4-benzodiazepin-2-one	407.2
C-F	Methyl alpha-bromophenylacetate (Aldrich)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(1-methoxycarbonyl-1-phenylmethyl)-1H-1,4-benzodiazepin-2-one	471.2
C-G	1-Bromo-2-ethylbutane (Aldrich)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(2-ethylbutyl)-1H-1,4-benzodiazepin-2-one	407.2
C-H	Bromomethylcyclohexane (Aldrich)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(cyclohexylmethyl)-1H-1,4-benzodiazepin-2-one	419.2
5 C-I	2-(Bromoethyl)benzene (Aldrich)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(2-phenylethyl)-1H-1,4-benzodiazepin-2-one	427.2
C-J	3-(Bromopropyl)benzene (K and K Laboratories)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(3-phenylpropyl)-1H-1,4-benzodiazepin-2-one	441.2
C-K	N-(2-Bromoethyl)phthalimide (Aldrich)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(2-(N-phthalimidyl)ethyl)-1H-1,4-benzodiazepin-2-one	496.2
C-L	2-Phenylbenzyl bromide (Aldrich)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(2-biphenylmethyl)-1H-1,4-benzodiazepin-2-one	489.2
C-M	Tetrahydrofurfuryl bromide (Lancaster)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-((2-tetrahydrofuran-2-yl)methyl)-1H-1,4-benzodiazepin-2-one	407.2
10 C-N	2-Bromomethyl-1,4-benzodioxane (Acros)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(2-(1,4-benzodioxan-2-yl)methyl)-1H-1,4-benzodiazepin-2-one	471.2
C-O	3-Bromomethyl-5-chlorobenzo[b]thiophene (Maybridge)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-((3-(5-chlorobenzo[b]thien-2-yl)methyl)-1H-1,4-benzodiazepin-2-one	503.1

-- 182 --

Ex.	Alkyl Halide	Intermediate	MS
C-P	1-Bromopinacolone (Lancaster)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(3,3-dimethyl-2-oxopropyl)-1H-1,4-benzodiazepin-2-one	421.1
C-Q	5-(Bromomethyl)benzofurazan (Maybridge)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(5-benzofurazanylmethyl)-1H-1,4-benzodiazepin-2-one	455.2
C-R	3-Phenoxypropyl bromide (Aldrich)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(3-phenoxypropyl)-1H-1,4-benzodiazepin-2-one	457.2
C-S	6-(Bromomethyl)-2-(trifluoromethyl)quinoline (Maybridge)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(6-(2-trifluoromethylquinolinyl)methyl)-1H-1,4-benzodiazepin-2-one	533.2
5 C-T	1-Bromo-2-methylbutane (Aldrich)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(2-methylbutyl)-1H-1,4-benzodiazepin-2-one	393.2
C-U	Ethyl bromide (Aldrich)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(ethyl)-1H-1,4-benzodiazepin-2-one	351.2
C-V	3-Picolyl chloride hydrochloride (Aldrich)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(3-pyridylmethyl)-1H-1,4-benzodiazepin-2-one	414.1
C-W	1-(2-Chloroacetyl)indoline (Maybridge)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(2-oxo-2-(N-indolinyl)ethyl)-1H-1,4-benzodiazepin-2-one	482.2
C-Y	4-(Chloromethyl)-3,5-dimethylisoxazole (Aldrich)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-((4-(3,5-dimethyl)isoxazolyl)methyl)-1H-1,4-benzodiazepin-2-one.	432.2
10 C-Z	2-Bromoethyl methyl ether (Aldrich)	3-(L-alaninyl)amino-5-phenyl-2,3-dihydro-1-(2-methoxyethyl)-1H-1,4-benzodiazepin-2-one	381.2

-- 183 --

## EXAMPLE C-AA

Synthesis of (S)-3-(L-phenylglyciny)amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one

5      Step A      Synthesis of (S)-3-(N'-(tert-Butoxycarbonyl)-L-phenylglyciny)amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one

To a solution of triethyl amine (519 uL, 3.8 mmol) and (S)-3-amino-5-phenyl-2-oxo-1,4-benzodiazepine (1.0 g, 3.8 mmol) (prepared according to the procedure of M. G. Bock et al., *J. Org. Chem.* **1987**, **52**, 3232-3239, incorporated herein by reference) in 100 mL of anhydrous methylene chloride at -20°C was added N-Boc-L-phenylglycine fluoride (Carpino et al, *J. Org. Chem.* **1991**, **56**, 2611-2614, incorporated herein by reference) in one portion. The reaction mixture was stirred for 15 min. and quenched with saturated aqueous bicarbonate (10 mL). The layers were separated, the organic layer washed sequentially with saturated aqueous bicarbonate, water and brine and then dried over sodium sulfate. Purification of the crude product using silica gel chromatography (10-50% ethyl acetate / hexane) gave 1.3 g (69%) of a hydroscopic white foam.

NMR data was as follows:

20      <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ = 1.35 (br s, 9H), 3.41 (s, 3H), 5.30-5.45 (m, 2H), 5.75-5.95 (m, 1H), 7.15-7.75 (m, 15H).

IR (CDCl<sub>3</sub>): 1709.7, 1676.6, 1489, 1166.3 cm<sup>-1</sup>.

IEX MS (M+1): 498.0.

25      Step B      Synthesis of (S)-3-(L-phenylglyciny)amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one

(S)-3-(N'-(tert-Butoxycarbonyl)-L-phenylglyciny)amino-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-2-one (1.27 g, 2.55 mmol) was added to 50 mL of a stirring solution of 15 % TFA in methylene chloride in one portion. After stirring 1 h, the reaction mixture was concentrated under reduced pressure and the residue dissolved in 100 mL of methylene chloride. This solution was washed twice with saturated sodium bicarbonate, once with brine and then dried

30

-- 184 --

over sodium sulfate. Purification of the crude product using silica gel column chromatography (5-10% methanol/methylene chloride) gave 743 mg (73%) of a very light green foam.

NMR data was as follows:

5  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 2.05 (br s, 1 H), 3.45 (s, 3 H), 5.51 (d, J = 8.39 Hz, 1H), 7.15-7.70 (m, 14 H), 8.60 (d, J = 830 Hz, 1 H).

IR ( $\text{CDCl}_3$ ): 1673.3, 1601.1, 1506.1  $\text{cm}^{-1}$ .

IEX MS (M+1): 399.2.

10

#### EXAMPLE C-AB

##### Synthesis of 3-(L-Alaninyl)amino-2,3-dihydro-1-(2-oxo-2-phenylethyl)-5-phenyl-1H-1,4-benzodiazepin-2-one

##### Step A Synthesis of 3-(Benzoxycarbonyl)amino-2,3-dihydro-1-(2-oxo-2-phenylethyl)-5-phenyl-1H-1,4-benzodiazepin-2-one

15 To a solution of 3-(Benzoxycarbonyl)amino-2,3-dihydro-5-phenyl-1H-1,4-benzodiazepin-2-one (Bock, M. G. et al, *Tetrahedron Lett.* 1987, 28, 939, incorporated herein by reference; 4.0 g, 10.4 mmol) in 40 mL of anhydrous DMF at 0°C was added potassium *tert*-butoxide (1.51 g, 13.5 mmol) in one portion. The reaction mixture was stirred 20 min. and  $\alpha$ -bromoacetophenone (Lancaster; 20 Windham, NH; 2.9 g, 14.6 mmol) was added. The reaction mixture was warmed to room temperature over 30 min. and then diluted with 100 mL of water and 200 mL of methylene chloride. The layers were separated. The organic layer was extracted with water and dried over sodium sulfate. Purification of the crude product by silica gel column chromatography (0-5% ethyl acetate/methylene 25 chloride) gave 4.2 g (81%) of an off white foam.

NMR data was as follows:

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  = 5.16 (s, 2 H), 5.34 (s, 2 H), 5.50 (d, J = 8.33 Hz, 1 H), 6.70 (d, J = 8.28 Hz, 1 H), 7.20-7.70 (m, 12 H), 7.91 (d, J = 7.54 Hz, 2 H).

30 IR ( $\text{CHCl}_3$ ): 1706.04, 1685.3, 1505.9, 1489.1, 1450.3, 1244.7  $\text{cm}^{-1}$ .

-- 185 --

IEX MS (M+1): 504.3.

**Step B**      Synthesis of 3-Amino-2,3-dihydro-1-(2-oxo-2-phenylethyl)-5-phenyl-1H-1,4-benzodiazepin-2-one

A solution of 3-(Benzyloxycarbonyl)amino-2,3-dihydro-1-(2-oxo-2-phenylethyl)-5-phenyl-1H-1,4-benzodiazepin-2-one (3.7 g, 7.36 mmol) in 100 mL of anhydrous methylene chloride was cooled to 0°C under nitrogen. A stream of anhydrous HBr gas was then bubbled through this solution for 1 h. The bubbler was removed and the reaction was warmed to room temperature under nitrogen. After stirring 1 h the reaction was concentrated under vacuum and the residue was redissolved in 20 mL of methylene chloride. The crude HBr salt of the product was precipitated from solution using 300 mL of anhydrous ether and collected by filtration as a light yellow solid. After washing with ether, the solid was dissolved in methylene chloride and saturated sodium bicarbonate. The layers were separated and the organic layer was extracted with saturated sodium bicarbonate. The combined aqueous layers were then back extracted twice with methylene chloride. The combined organic layers were extracted once with water and dried over sodium sulfate. After concentration under vacuum, 2.27 g of the product was obtained as an orange foam which was carried on without further purification.

NMR data was as follows:

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ = 2.60 (br s, 2 H), 4.72 (s, 1 H), 5.34 (s, 2 H), 7.10-7.70 (m, 12 H), 7.91 (d, J = 7.60 Hz, 2 H).

IEX MS (M+1): 370.2

**Step C**      Synthesis of 3-(N'-(tert-Butoxycarbonyl)-L-alaninyl)amino-2,3-dihydro-1-(2-oxo-2-phenylethyl)-5-phenyl-1H-1,4-benzodiazepin-2-one

To a solution of HOBt-H<sub>2</sub>O (697 mg, 5.16 mmol), N,N-diisopropylethylamine (900 uL, 5.16 mmol) and N-t-BOC-L-alanine (975 mg, 5.16 mmol) in 20 mL of anhydrous THF at 0°C was added 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (EDCI; 986 mg, 5.16 mmol) in one portion. After stirring 5 min., a solution of 3-amino-2,3-dihydro-1-

-- 186 --

(2-oxo-2-phenylethyl)-5-phenyl-1H-1,4-benzodiazepin-2-one (2.0 g, 5.43 mmol) in 20 mL of anhydrous THF was added via syringe and the reaction mixture was warmed to room temperature and stirred overnight. The reaction mixture was diluted with 200 mL methylene chloride, extracted sequentially with 10 % citric acid, saturated sodium bicarbonate, water and brine and then dried over sodium sulfate. Purification of the crude product using silica gel chromatography (10%-30% ethyl acetate/methylene chloride) gave 2.59 g (93%) of a white foam.

NMR data was as follows:

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  = 1.30-1.60 (m, 12 H), 4.35 (br s, 1 H), 5.00-5.50 (m, 3 H), 5.65-5.70 (m, 1 H), 7.15-7.65 (m, 12 H), 7.70-7.80 (m, 1 H), 7.85-7.95 (m, 1 H).

IR (CHCl<sub>3</sub>): 1705.8, 1678.8, 1488.7, 1450.2, 1230.4, 1164.4 cm<sup>-1</sup>.

IEX MS (M+1): 541.2.

**Step D**      Synthesis of 3-(L-Alaninyl)amino-2,3-dihydro-1-(2-oxo-2-phenylethyl)-5-phenyl-1H-1,4-benzodiazepin-2-one

3-(N'-(*tert*-Butoxycarbonyl)-L-alaninyl)amino-2,3-dihydro-1-(2-oxo-2-phenylethyl)-5-phenyl-1H-1,4-benzodiazepin-2-one (2.5 g, 4.63 mmol) was added to 100 mL of a stirring solution of 15 % TFA/methylene chloride in one portion. After stirring 2 h, the reaction mixture was concentrated under reduced pressure and the residue was dissolved in 150 mL of methylene chloride. This solution was washed twice with saturated sodium bicarbonate, once with brine and then dried over sodium sulfate. Purification of the crude product using silica gel column chromatography (1-10% methanol/methylene chloride) gave 1.91 g (94%) of the title compound as a white foam.

NMR data was as follows:

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  = 1.30-1.50 (m, 3 H), 1.80-2.20 (br s, 2 H), 3.55-3.75 (m, 1 H), 5.20-5.45 (m, 2 H), 5.67 (t, *J* = 7.48 Hz, 1 H), 7.20-7.65 (m, 12 H), 7.90 (d, *J* = 7.7 Hz, 2 H), 8.80 (dd, *J*<sub>1</sub> = 25.09 Hz, *J*<sub>2</sub> = 8.33 Hz, 1 H).



-- 187 --

EX MS (M+1): 441.2.

**EXAMPLE C-AC**

5        Synthesis of 3-(L-Alaninyl)amino-2,3-dihydro-1-(4,4,4-trifluorobutyl)-5-phenyl-1H-1,4-benzodiazepin-2-one

Step A        Synthesis of 3-(Benzyloxycarbonyl)amino-2,3-dihydro-1-(4,4,4-trifluorobutyl)-5-phenyl-1H-1,4-benzodiazepin-2-one

To a solution of 3-(benzyloxycarbonyl)amino-2,3-dihydro-5-phenyl-1H-1,4-benzodiazepin-2-one (3.7 g, 9.61 mmol) in 40 mL of anhydrous DMF at 0°C was added potassium *tert*-butoxide (1.6 g, 14.4 mmol) in one portion. The reaction mixture was stirred 20 min. and 4,4,4-trifluoro-1-bromobutane (Lancaster; Windham, NH; 2.6 g, 13.4 mmol) was added. The reaction mixture was warmed to room temperature over 30 min. and then diluted with 100 mL of water and 200 mL of methylene chloride. The layers were separated. The organic layer was extracted with water and dried over sodium sulfate. Purification of the crude product by silica gel column chromatography (0-3 % ethyl acetate / methylene chloride) gave 1.52 g (32 %) of an off white foam.

NMR data was as follows:

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ = 1.50-2.10 (m, 4 H), 3.70-3.90 (m, 1 H), 4.35-4.55 (m, 1 H), 5.15 (s, 2 H), 5.33 (d, J = 8.47 Hz, 1 H), 6.67 (d, J = 8.40 Hz, 1 H), 7.2-7.70 (m, 14 H).

      IR (CHCl<sub>3</sub>): 1720.4, 1683.0, 1604.8, 1505.5, 1451.1, 1323.9, 1254.5, 1148.4 cm<sup>-1</sup>.

      IEX MS (M+1): 496.3.

25        Step B        Synthesis of 3-Amino-2,3-dihydro-1-(4,4,4-trifluorobutyl)-5-phenyl-1H-1,4-benzodiazepin-2-one

A solution of 3-(benzyloxycarbonyl)amino-2,3-dihydro-1-(4,4,4-trifluorobutyl)-5-phenyl-1H-1,4-benzodiazepin-2-one (1.42 g, 2.87 mmol) in 50 mL of anhydrous methylene chloride was cooled to 0°C under nitrogen. A stream of anhydrous HBr gas was slowly bubbled through the solution for 1 h.

-- 188 --

The bubbler was removed and the reaction was warmed to room temperature under nitrogen. After stirring for 1 h, the reaction was concentrated under vacuum and the residue was redissolved in 10 mL of methylene chloride. The crude HBr salt of the product was precipitated from solution using 90 mL of anhydrous ether and collected by filtration. After washing with ether, the HBr salt was dissolved in methylene chloride and saturated sodium bicarbonate. The layers were separated and the organic layer was extracted with saturated sodium bicarbonate. The combined aqueous layers were then back extracted twice with methylene chloride. The combined organic layers were extracted once with water and dried over sodium sulfate. After concentration under vacuum, 1.06 g (100%) of the product was obtained as a white foam which was carried on without further purification. NMR data was as follows:

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  = 1.60-2.10 (m, 4 H), 2.76 (br s, 2 H), 3.75-3.85 (m, 1 H), 4.40-4.60 (m, 2 H), 7.20-7.70 (m, 9 H).

IEX MS (M+1): 362.1.

**Step C**      Synthesis of 3-(N'-(*tert*-Butoxycarbonyl)-L-alaninyl)amino-2,3-dihydro-1-(4,4,4-trifluorobutyl)-5-phenyl-1H-1,4-benzodiazepin-2-one

To a solution of HOBt- $\text{H}_2\text{O}$  (373 mg, 2.76 mmol), N,N-diisopropylethylamine (481  $\mu\text{L}$ , 2.76 mmol) and N-t-BOC-L-alanine (522 mg, 2.76 mmol) in 10 mL of anhydrous THF at 0°C was added 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (EDCI; 527 mg, 2.76 mmol) in one portion. After stirring 5 min., a solution of 3-amino-2,3-dihydro-1-(4,4,4-trifluorobutyl)-5-phenyl-1H-1,4-benzodiazepin-2-one (1.05 g, 2.91 mmol) in 10 mL of anhydrous THF was added via syringe and the reaction mixture was warmed to room temperature and stirred overnight. The reaction mixture was diluted with 100 mL methylene chloride, extracted sequentially with 10% citric acid, saturated sodium bicarbonate, water and brine and then dried over sodium sulfate. Purification of the crude product using silica gel chromatography (10%-30% ethyl acetate/methylene chloride) gave 1.28 g (83%) of a white foam.

-- 189 --

NMR data was as follows:

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  = 1.40-2.10 (m, 16 H), 3.70-3.85 (m, 1 H), 4.30-4.55 (m, 2 H), 5.10 (br s, 1 H), 5.45-5.55 (m, 1 H), 7.25-7.80 (m, 10 H).

5 IR ( $\text{CDCl}_3$ ): 1676.6, 1605.2, 1488.6, 1450.9, 1393.2, 1338.7, 1324.9, 1253.8, 1150.4  $\text{cm}^{-1}$ .

IEX MS ( $M+1$ ): 533.1.

Step D      Synthesis of 3-(L-Alaninyl)amino-2,3-dihydro-1-(4,4,4-trifluorobutyl)-5-phenyl-1H-1,4-benzodiazepin-2-one

10 3-(N'-(*tert*-Butoxycarbonyl)-L-alaninyl)amino-2,3-dihydro-1-(4,4,4-trifluorobutyl)-5-phenyl-1H-1,4-benzodiazepin-2-one (1.21 g, 2.27 mmol) was added to 50 mL of a stirring solution of 15 % TFA / methylene chloride in one portion. After stirring 2 h, the reaction mixture was concentrated under reduced pressure and the residue was dissolved in 100 mL of methylene chloride. This  
15 solution was washed twice with saturated sodium bicarbonate, once with brine and then dried over sodium sulfate. Purification of the crude product using silica gel column chromatography (1-5% methanol / methylene chloride) gave 670 mg (68%) of a light pink foam.

NMR data was as follows:

20  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  = 1.43 (t,  $J$  = 7.0 Hz, 3 H), 1.60-2.20 (m, 7 H), 3.60-3.85 (m, 2 H), 4.35-4.55 (m, 1 H), 5.51 (dd,  $J_1$  = 8.36 Hz,  $J_2$  = 2.48 Hz, 1 H), 7.20-7.70 (m, 9 H), 8.80 (dd,  $J_1$  = 27.73 Hz,  $J_2$  = 8.34 Hz, 1 H).

IEX MS ( $M+1$ ): 433.2.

25

**EXAMPLE C-AE**

Synthesis of 3-[(L-Alaninyl)amino]-2,3-dihydro-1-methyl-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one

30 Step A      Synthesis of 3-Amino-2,3-dihydro-1-methyl-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one

-- 190 --

The title compound was synthesized as described in *Synth. Commun.*, **26(4)**, 721-727 (1996), incorporated herein by reference.

**Step B**      Synthesis of 3-[(N-tert-Butoxycarbonyl-L-alaninyl)amino]-2,3-dihydro-1-methyl-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one

5            A solution of L-Boc-alanine (1.74 g, 9.20 mmol), HOBt monohydrate (1.24 g, 9.20 mmol), diisopropylethylamine (1.6 mL, 9.20 mmol) and CH<sub>2</sub>Cl<sub>2</sub> (30 mL) was purged with nitrogen and cooled in an ice bath. To the cold solution was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.76 g, 9.20 mmol) followed by a solution of 3-amino-2,3-dihydro-1-methyl-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one (2.45 g, 9.20 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (15 mL).  
10           The cold bath was removed and the solution stirred overnight at room temperature. The reaction mixture was extracted with H<sub>2</sub>O, 0.1 N aq. citric acid, 5% aq. NaHCO<sub>3</sub>, and brine. The remaining CH<sub>2</sub>Cl<sub>2</sub> solution was dried (MgSO<sub>4</sub>) and concentrated to a tan foam. The title compound was crystallized from  
15           CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to give 3.47 g (86% yield) of white crystals, mp. 228-229°C.

Anal. Calcd for C<sub>23</sub>H<sub>27</sub>N<sub>5</sub>O<sub>4</sub>: C, 63.14; H, 6.22; N, 16.01. Found: C, 63.25; H, 6.15; N, 15.95. MS (FD<sup>+</sup>) 437 m/z.

**Step C**      Synthesis of 3-[(L-Alaninyl)amino]-2,3-dihydro-1-methyl-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one

20           A solution of 3-[(N-tert-butoxycarbonyl-L-alaninyl)amino]-2,3-dihydro-1-methyl-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one (3.42 g, 7.82 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (90 mL) was cooled in an ice bath and treated with TFA (13.2 mL, 172 mmol). The cold bath was removed and the solution stirred at room temperature for four hours. The reaction mixture was washed with 1 M aq. K<sub>2</sub>CO<sub>3</sub> and the aqueous  
25           portion back-extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined extracts were washed with H<sub>2</sub>O, dried (MgSO<sub>4</sub>) and concentrated to obtain 1.75 g (66% yield) of the title compound as an off-white foam. MS (IS<sup>+</sup>) 338 (m/e).

<sup>1</sup>HNMR (CDCl<sub>3</sub>): δ = 8.76-8.86 (1H, m), 8.63 (1H, m), 8.17 (1H, m), 7.82 (2H, m), 7.60 (1H, m), 7.41 (3H, m), 5.60 (1H, m), 3.63 (1H, m), 3.49  
30           (3H, s), 1.66 (2H, broad), 1.45 (3H, m).

-- 191 --

## EXAMPLE C-AF

Synthesis of 3-[(L-Alaninyl)aminol]-2,3-dihydro-1-(2-N,N-diethylaminoethyl)-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one

Step A      Synthesis of 3-Amino-2,3-dihydro-1-(2-N,N-diethylaminoethyl)-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one

The title compound was synthesized as described in *Synth. Commun.*, **26(4)**, 721-727 (1996), incorporated herein by reference.

Step B      Synthesis of 3-[(N-tert-Butoxycarbonyl-L-alaninyl)aminol]-2,3-dihydro-1-(2-N,N-diethylaminoethyl)-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one

A solution of L-Boc-alanine (1.80 g, 9.50 mmol), HOBt monohydrate (1.28 g, 9.50 mmol), diisopropylethylamine (1.65 mL, 9.50 mmol) and CH<sub>2</sub>Cl<sub>2</sub> (40 mL) was purged with nitrogen and cooled in an ice bath. To the cold solution was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.82 g, 9.50 mmol) followed by a solution of 3-amino-2,3-dihydro-1-(2-N,N-diethylaminoethyl)-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one (3.34 g, 9.50 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (25 mL). The cold bath was removed and the solution stirred overnight at room temperature. The reaction mixture was extracted with H<sub>2</sub>O, 5% aq. NaHCO<sub>3</sub>, and brine. The remaining CH<sub>2</sub>Cl<sub>2</sub> solution was dried (MgSO<sub>4</sub>) and concentrated to a tan foam. The title compound was isolated via column chromatography (2% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to 10% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to give 3.53 g (71% yield) of yellow foam.

MS (FD<sup>+</sup>) 522 (m/z).

<sup>1</sup>HNMR (CDCl<sub>3</sub>): δ = 8.62 (1H, d), 8.11 (1H, m), 7.80 (2H, m), 7.59 (2H, m), 7.32-7.45 (2H, m), 5.54 (1H, m), 5.02-5.18 (1H, m), 4.38 (1H, m), 4.20 (1H, m), 3.83 (1H, m), 2.62 (2H, t), 2.44 (4H, m), 1.40-1.56 (12H, m), 0.88 (6H, m).

Step C      Synthesis of 3-[(L-Alaninyl)aminol]-2,3-dihydro-1-(2-N,N-diethylaminoethyl)-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one

The title compound was synthesized using the procedure described in Example C-AE, Step C. A solution of 3-[(N-tert-butoxycarbonyl-L-

-- 192 --

alaninyl)amino]-2,3-dihydro-1-(2-N,N-diethylaminoethyl)-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one (3.52 g, 6.73 mmol) was treated with TFA (11.4 mL, 148 mmol) to give 2.61 g (92% yield) the title compound as a light yellow foam.

MS ( $\text{IS}^+$ ) 423 (m/e).

5  $^1\text{H}$ NMR ( $\text{CDCl}_3$ ):  $\delta$  = 8.78-8.93 (1H, m), 8.62 (1H, d), 8.11 (1H, m), 7.80 (2H, m), 7.58 (2H, m), 7.39 (2H, m), 5.58 (1H, m), 4.22 (1H, m), 3.88 (1H, m), 3.61 (1H, m), 2.67 (2H, t), 2.49 (4H, m), 1.73 (2H, broad), 1.42 (3H, m), 0.91 (6H, m).

10

#### EXAMPLE C-AG

##### Synthesis of 3-[(L-Alaninyl)amino]-2,3-dihydro-1-(3,3-dimethyl-2-oxobutyl)-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one

##### Step A Synthesis of 3-Amino-2,3-dihydro-1-(3,3-dimethyl-2-oxobutyl)-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one

15 The title compound was synthesized as described in *Synth. Commun.*, 26(4), 721-727 (1996), incorporated herein by reference.

##### Step B Synthesis of 3-[(N-tert-Butoxycarbonyl-L-alaninyl)amino]-2,3-dihydro-1-(3,3-dimethyl-2-oxobutyl)-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one

20 A solution of L-Boc-alanine (1.57 g, 8.33 mmol), HOBt monohydrate (1.13 g, 8.33 mmol), diisopropylethylamine (1.45 mL, 8.33 mmol) and  $\text{CH}_2\text{Cl}_2$  (40 mL) was purged with nitrogen and cooled in an ice bath. To the cold solution was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.60 g, 8.33 mmol) followed by a solution of 3-amino-2,3-dihydro-1-(3,3-dimethyl-2-oxobutyl)-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one (2.92 g, 8.33 mmol)  
25 dissolved in  $\text{CH}_2\text{Cl}_2$  (25 mL). The cold bath was removed and the solution stirred overnight at room temperature. The reaction mixture was extracted with  $\text{H}_2\text{O}$ , 0.1 N aq. citric acid, 5% aq.  $\text{NaHCO}_3$ , and brine. The remaining  $\text{CH}_2\text{Cl}_2$  solution was dried ( $\text{MgSO}_4$ ) and concentrated to a yellow foam. The title

-- 193 --

compound was isolated via column chromatography (20% EtOAc/hexanes to 60% EtOAc/hexanes) to give 4.19 g (96% yield) of light yellow foam.

MS (FD<sup>+</sup>) 521 (m/z).

<sup>1</sup>HNMR (CDCl<sub>3</sub>): δ = 8.65 (1H, t), 8.17 (1H, t), 7.90 (1H, t), 7.71-7.85 (1H, m), 7.54 (1H, m), 7.44 (1H, t), 7.37 (1H, d), 7.24-7.32 (1H, m), 7.14 (1H, m), 5.67 (1H, dd), 5.18 (1H, broad), 4.93-5.07 (1H, m), 4.50-4.64 (1H, m), 4.38 (1H, broad), 1.42-1.51 (12H, m), 1.26 (9H, d).

Step C      Synthesis of 3-[(L-Alaninyl)amino]-2,3-dihydro-1-(3,3-dimethyl-2-oxobutyl)-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one

The title compound was synthesized using the procedure described in Example C-AE, Step C. A solution of 3-[(N-tert-butoxycarbonyl-L-alaninyl)amino]-2,3-dihydro-1-(3,3-dimethyl-2-oxobutyl)-5-(2-pyridyl)-1H-1,4-benzodiazepin-2-one (4.18 g, 8.01 mmol) was treated with TFA (13.6 mL, 176 mmol) to give 3.14 g (93% yield) the title compound as an off-white foam.

MS (IS<sup>+</sup>) 422 (m/e).

<sup>1</sup>HNMR (CDCl<sub>3</sub>) δ 8.85-8.99 (1H, m), 8.68 (1H, d), 8.20 (1H, t), 7.87 (1H, t), 7.58 (1H, t), 7.42 (2H, m), 7.30 (1H, t), 7.17 (1H, d), 5.72 (1H, m), 5.08 (1H, d), 4.60 (1H, d), 3.66 (1H, m), 1.47 (3H, m), 1.28 (9H, m).

20

**EXAMPLE C-AH**

Synthesis of 3-[(L-Alaninyl)amino]-2,3-dihydro-1-methyl-5-(2-thiazyl)-1H-1,4-benzodiazepin-2-one

Step A      Synthesis of 3-Amino-2,3-dihydro-1-methyl-5-(2-thiazyl)-1H-1,4-benzodiazepin-2-one

The title compound was synthesized in a manner similar to the procedure described in *Synth. Commun.*, 26(4), 721-727 (1996), incorporated herein by reference, starting with 2-(2-aminobenzoyl)thiazole (prepared as described in *Tetrahedron*, 51(3), 773-786, (1995), incorporated herein by reference).

MS (IS<sup>+</sup>) 273 (m/e).

-- 194 --

<sup>1</sup>HNMR (CDCl<sub>3</sub>): δ = 7.83-7.94 (2H, m), 7.61 (1H, t), 7.50 (1H, d), 7.34 (2H, m), 4.60 (1H, s), 3.46 (3H, s), 1.97 (2H, broad).

**Step B**      Synthesis of 3-[(N-tert-Butoxycarbonyl-L-alaninyl)amino]-2,3-dihydro-1-methyl-5-(2-thiazyl)-1H-1,4-benzodiazepin-2-one

5            A solution of L-Boc-alanine (1.85 g, 9.77 mmol), HOBt monohydrate (1.32 g, 9.77 mmol), diisopropylethylamine (1.70 mL, 9.77 mmol) and CH<sub>2</sub>Cl<sub>2</sub> (30 mL) was purged with nitrogen and cooled in an ice bath. To the cold solution was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.87 g, 9.77 mmol) followed by a solution of 3-amino-2,3-dihydro-1-methyl-5-(2-thiazyl)-  
10    1H-1,4-benzodiazepin-2-one (2.66 g, 9.77 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL). The cold bath was removed and the solution stirred overnight at room temperature. The reaction mixture was extracted with H<sub>2</sub>O, 0.1 N aq. citric acid, 5% aq. NaHCO<sub>3</sub>, and brine. The remaining CH<sub>2</sub>Cl<sub>2</sub> solution was dried (MgSO<sub>4</sub>) and concentrated to a light yellow foam. The title compound was crystallized  
15    from EtOAc/hexane to give 3.22 g (74% yield) of white crystals, mp. 196-197°C.

Anal. Calcd for C<sub>21</sub>H<sub>25</sub>N<sub>5</sub>O<sub>4</sub>S: C, 56.87; H, 5.68; N, 15.79. Found: C, 56.74; H, 5.75; N, 15.55.

MS (IS<sup>+</sup>) 444 m/e.

20    **Step C**      Synthesis of 3-[(L-Alaninyl)amino]-2,3-dihydro-1-methyl-5-(2-thiazyl)-1H-1,4-benzodiazepin-2-one

The title compound was synthesized using the procedure described in Example C-AE, Step C.

**EXAMPLE C-AI**

25    Synthesis of 3-[(L-Alaninyl)amino]-2,3-dihydro-1-methyl-5-(thiophen-2-yl)-1H-1,4-benzodiazepin-2-one

**Step A**      Synthesis of 3-Amino-2,3-dihydro-1-methyl-5-(2-thiophen-2-yl)-1H-1,4-benzodiazepin-2-one

30    The title compound was synthesized in a manner similar to the procedure described in *Synth. Commun.*, 26(4), 721-727 (1996), incorporated herein by



-- 195 --

reference, starting with 2-(2-aminobenzoyl)thiophene (prepared as described in *Collect. Czech. Chem. Commun.*, 34(2), 468-478, (1969), incorporated herein by reference).

MS ( $\text{IS}^+$ ) 272 (m/e).

5  $^1\text{H}$ NMR ( $\text{CDCl}_3$ ):  $\delta$  = 7.68 (1H, d), 7.60 (1H, t), 7.48 (1H, m), 7.35 (2H, d), 7.28 (1H, m), 7.15 (1H, d), 7.05 (1H, d), 4.50 (1H, broad), 3.45 (3H, s), 2.26 (2H, broad).

Step B      Synthesis of 3-[(N-tert-Butoxycarbonyl-L-alaninyl)amino]-2,3-dihydro-1-methyl-5-(thiophen-2-yl)-1H-1,4-benzodiazepin-2-one

10      The title compound was synthesized in a manner similar to the procedure described in Example C-AH, Step B.

MS ( $\text{IS}^+$ ) 443 (m/e).

$^1\text{H}$ NMR ( $\text{CDCl}_3$ ):  $\delta$  = 7.69 (1H, d), 7.61 (2H, m), 7.48 (1H, d), 7.27-7.42 (2H, m), 7.18 (1H, m), 7.05 (1H, m), 5.51 (1H, d), 5.13 (1H, broad), 4.36  
15 (1H, broad), 3.44 (3H, s), 1.38-1.57 (12H, m).

Step C      Synthesis of 3-[(L-Alaninyl)amino]-2,3-dihydro-1-methyl-5-(thiophen-2-yl)-1H-1,4-benzodiazepin-2-one

The title compound was synthesized in a manner similar to the procedure described in Example C-AE, Step C.

20      MS ( $\text{IS}^+$ ) 343 (m/e).

$^1\text{H}$ NMR ( $\text{CDCl}_3$ ):  $\delta$  = 8.55 (1H, d), 7.68 (1H, d), 7.59 (1H, m), 7.48 (1H, d), 7.36 (1H, d), 7.31 (1H, d), 7.16 (1H, m), 7.04 (1H, t), 5.54 (1H, d), 3.58 (1H, m), 3.45 (3H, s), 1.41 (3H, d).

-- 196 --

## EXAMPLE 5-A

Synthesis of 3-(L-Alaninyl)-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1-benzazepin-2-one5      Step A      Preparation of 1-phenyl-1-[2-N-( $\alpha$ -isopropylthio)-N'-(benzyloxycarbonyl)-glycinyl]-aminophenyl]ethylene

A solution of  $\alpha$ -(isopropylthio)-N-(benzyloxycarbonyl)glycine (1 eq; prepared according to Zoller, V. and D. Ben-Ishai, *Tetrahedron* **1975**, *31*, 863, incorporated herein by reference) in dry THF was cooled to 0°C and treated with oxalyl chloride (1 eq.) and 3 drops of DMF. After stirring for 15 minutes at 0°C, the cooling bath was removed and stirring continued at ambient temperature for 40 minutes. The solution was re-cooled to 0°C. A solution of 1-phenyl-1-(2-aminophenyl)ethylene (0.9 eq.; Arienti, A. et al. *Tetrahedron* **1997**, *53*, 3795, incorporated herein by reference) and 4-methylmorpholine (2.0 eq.) in dry THF was added via cannulation to the acid chloride. The cooling bath was removed and the reaction stirred at ambient for 5 hours. The reaction was diluted with methylene chloride and washed with 0.5 M citric acid, saturated aqueous NaHCO<sub>3</sub>, and brine. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The residue was purified via flash chromatography eluting with CH<sub>2</sub>Cl<sub>2</sub> then 90:10 CH<sub>2</sub>Cl<sub>2</sub>/ethyl acetate giving a pale yellow oil.

20      C<sub>27</sub>H<sub>28</sub>N<sub>2</sub>O<sub>3</sub>S (MW = 460.60); mass spectroscopy (MH<sup>+</sup>) 461.4.  
Anal. Calcd for C<sub>27</sub>H<sub>28</sub>N<sub>2</sub>O<sub>3</sub>S: C, 70.41; H, 6.13; N, 6.08. Found: C, 70.42; H, 6.05; N, 6.05.

25      Step B      Preparation of 3-(benzyloxycarbonyl)-amino-2,3-dihydro-5-phenyl-1H-1-benzazepin-2-one

A solution of 1-phenyl-1-[2-N-( $\alpha$ -isopropylthio)-N'-(benzyloxycarbonyl)-glycinyl-aminophenyl]ethylene (1 eq) in acetonitrile under nitrogen was treated with mercury(II) chloride (1.0 equiv.; Aldrich). A white precipitate formed immediately after the mercury(II) chloride had dissolved. The mixture was heated to reflux for 2.5 hours; an additional 0.05 equiv. of mercury(II) chloride was added and refluxing continued for 1 hour. The reaction was cooled to

-- 197 --

ambient temperature and the mercury salts were filtered washing with methylene chloride. The filtrate was concentrated *in vacuo*; the resultant residue was taken up in methylene chloride and filtered to remove additional mercury salts. The filtrate was washed with water. The aqueous layer was back-extracted five times  
5 with methylene chloride. The combined organics were allowed to stand overnight; additional mercury salts were filtered. The filtrate was dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated *in vacuo*. The crude oil was purified via flash chromatography eluting with a gradient from  $\text{CH}_2\text{Cl}_2$  to 82:18  $\text{CH}_2\text{Cl}_2$ /ethyl acetate giving a tan solid. The solid could be further purified by trituration with  
10 diethyl ether which provided a white solid.

$\text{C}_{24}\text{H}_{20}\text{N}_2\text{O}_3$  (MW = 384.43); mass spectroscopy found (M+H) 385.1.

Anal. Calcd for  $\text{C}_{24}\text{H}_{20}\text{N}_2\text{O}_3 \cdot 0.5 \text{H}_2\text{O}$ : C, 73.27; H, 5.38; N, 7.12.

Found: C, 73.41; H, 5.13; N, 7.30.

15 Step C      Preparation of 3-(benzyloxycarbonyl)-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1-benzazepin-2-one

Following General Procedure 4-A and using 3-(benzyloxycarbonyl)-amino-2,3-dihydro-5-phenyl-1H-1-benzazepin-2-one, the title intermediate was prepared as a white solid.

$\text{C}_{25}\text{H}_{22}\text{N}_2\text{O}_3$  (MW = 398.4); mass spectroscopy found (M+H) 399.2.

20 Anal. Calcd for  $\text{C}_{25}\text{H}_{22}\text{N}_2\text{O}_3$ : C, 75.36; H, 5.57; N, 7.03. Found: C, 75.21; H, 5.57; N, 7.13.

Step D      Preparation of 3-amino-1,3-dihydro-1-methyl-5-phenyl-2H-1-benzazepin-2-one

Following General Procedure 4-B and using 3-(benzyloxycarbonyl)-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1-benzazepin-2-one, the title intermediate was  
25 prepared as an amber oil which was used immediately in Step E.

Step E      Preparation of 3-[N'-(tert-butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1-benzazepin-2-one

Following General Procedure D using N-Boc Alanine and 3-amino-1,3-dihydro-1-methyl-5-phenyl-2H-1-benzazepin-2-one, the title intermediate was  
30

-- 198 --

prepared as a white solid.

$C_{25}H_{29}N_3O_4$  (MW = 435.57); mass spectroscopy found (M+H) 436.3.

$^1H$  NMR (300 MHz,  $CDCl_3$ )  $\delta$  7.54 (2H, d, J=4.6 Hz), 7.52-7.25 (18H, m), 5.90 (1H, d, J=5.3 Hz), 5.88 (1H, d, J=5.3 Hz), 5.1 (1H, bs), 4.9 (1H, bs), 4.60 (2H, m), 4.31 (2H, m), 3.48 (6H, s), 1.48 (9H, s), 1.46 (9H, s), 1.43 (3H, d, J=4.5 Hz), 1.40 (3H, d, J=4.1 Hz).

**Step F**      Preparation of 3-(L-alaninyl)-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1-benzazepin-2-one

Following General Procedure 4-C using 3-[N'-(*tert*-butylcarbamate)-L-alaninyl]-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1-benzazepin-2-one, the title intermediate was prepared as a white foam. No further purification was necessary.

$C_{20}H_{21}N_3O_2$  (MW = 335.40); mass spectroscopy found (M+H) 336.2.

Anal. Calcd for  $C_{20}H_{21}N_3O_2$ : C, 71.62; H, 6.31; N, 12.53. Found: C, 71.78; H, 6.54; N, 12.22.

**EXAMPLE 5-B**

Synthesis of 3-(L-Alaninyl)-amino-2,3,4,5-tetrahydro-1-methyl-5-phenyl-1H-1-benzazepin-2-one

**20 Step A**      Preparation of 3-amino-2,3,4,5-tetrahydro-1-methyl-5-phenyl-2H-1-benzazepin-2-one

A solution of 3-(benzyloxycarbonyl)-amino-2,3-dihydro-1-methyl-5-phenyl-1H-1-benzazepin-2-one (Example 5-A, Steps A-C) in methanol under nitrogen was treated with 10% palladium on carbon (0.4 equiv.). The reaction vessel was placed under a balloon of hydrogen and stirred for 3 hours. The reaction flask was

-- 199 --

flushed well with nitrogen and the reaction mixture filtered through celite washing with  $\text{CH}_2\text{Cl}_2$ . The filtrate was concentrated to a white foam.

$\text{C}_{17}\text{H}_{18}\text{N}_2\text{O}$  (MW = 266.37); mass spectroscopy found (M+H) 267.1.

Anal. Calcd for  $\text{C}_{17}\text{H}_{18}\text{N}_2\text{O}$ : C, 76.66; H, 6.81; N, 10.52. Found: C, 76.56; H, 6.83; N, 10.38.

**Step B**      Preparation of 3-[N'-(*tert*-butylcarbamate)-L-alaninyl]-amino-2,3,4,5-tetrahydro-1-methyl-5-phenyl-1H-1-benzazepin-2-one

Following General Procedure D using N-Boc Alanine and 3-amino-2,3,4,5-tetrahydro-1-methyl-5-phenyl-2H-1-benzazepin-2-one, the title intermediate was prepared as a white foam.

$\text{C}_{25}\text{H}_{31}\text{N}_3\text{O}_4$  (MW = 437.59); mass spectroscopy found (M+H) 438.2.

Anal. Calcd for  $\text{C}_{25}\text{H}_{31}\text{N}_3\text{O}_4$ : C, 68.63; H, 7.14; N, 9.60. Found: C, 68.93; H, 7.13; N, 9.49.

**Step C**      Preparation of 3-(L-alaninyl)-amino-2,3,4,5-tetrahydro-1-methyl-5-phenyl-1H-1-benzazepin-2-one

Following General Procedure 4-C using 3-[N'-(*tert*-butylcarbamate)-L-alaninyl]-amino-2,3,4,5-tetrahydro-1-methyl-5-phenyl-1H-1-benzazepin-2-one, the title intermediate was prepared as a white foam. No further purification was necessary.

$\text{C}_{20}\text{H}_{23}\text{N}_3\text{O}_2$  (MW = 337.46); mass spectroscopy found (M+H) 338.2.

Anal. Calcd for  $\text{C}_{20}\text{H}_{23}\text{N}_3\text{O}_2$ : C, 71.19; H, 6.87; N, 12.45. Found: C, 71.38; H, 6.83; N, 12.51.

**EXAMPLE 5-C**

Synthesis of 3-(L-Alaninyl)-amino-2,3,4,5-tetrahydro-1-methyl-5-phenyl-1H-1-benzazepin-2-one

**Step A**      Preparation of 3-amino-2,3,4,5-tetrahydro-1-methyl-5-phenyl-2H-1-benzazepin-2-one

To a flask containing 300 mL of freshly condensed liquid ammonia at  $-70^\circ\text{C}$  was added lithium metal (4.1 equiv.). The dark blue slurry was warmed to  $-45^\circ\text{C}$  and treated with a pre-cooled solution of 3-(benzyloxycarbonyl)-amino-2,3-

-- 200 --

dihydro-1-methyl-5-phenyl-1H-1-benzazepin-2-one (Example 4-AC, Steps A-C) in 30 mL distilled THF. After 10 min a solution of tert-butanol (4.0 equiv.) in distilled THF was added. After an additional 10 min the reaction was quenched with ammonium chloride. The cooling bath was removed and the ammonia  
5 allowed to evaporate overnight. The contents were partitioned between CH<sub>2</sub>Cl<sub>2</sub> and saturated aqueous NaHCO<sub>3</sub>. The organic phase was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. It was purified via liquid chromatography eluting with a gradient from 99:1 to 90:10 CH<sub>2</sub>Cl<sub>2</sub>/MeOH.

C<sub>17</sub>H<sub>18</sub>N<sub>2</sub>O (MW = 266.37); mass spectroscopy found (M+H) 267.0.

10 **Step B**      Preparation of 3-[N'-(*tert*-butylcarbamate)-L-alaninyl]-amino-2,3,4,5-tetrahydro-1-methyl-5-phenyl-1H-1-benzazepin-2-one

Following General Procedure D using N-Boc-alanine and 3-amino-2,3,4,5-tetrahydro-1-methyl-5-phenyl-2H-1-benzazepin-2-one, the title intermediate was prepared as a white foam.

15      C<sub>25</sub>H<sub>31</sub>N<sub>3</sub>O<sub>4</sub> (MW = 437.59); mass spectroscopy found (M+H) 438.2.  
Anal. Calcd for C<sub>25</sub>H<sub>31</sub>N<sub>3</sub>O<sub>4</sub>: C, 68.63; H, 7.14; N, 9.60. Found: C, 68.70; H, 7.15; N, 9.54.

**Step C**      Preparation of 3-(L-alaninyl)-amino-2,3,4,5-tetrahydro-1-methyl-5-phenyl-1H-1-benzazepin-2-one

20      Following General Procedure 4-C using 3-[N'-(*tert*-butylcarbamate)-L-alaninyl]-amino-2,3,4,5-tetrahydro-1-methyl-5-phenyl-1H-1-benzazepin-2-one, the title intermediate was prepared as a white foam. No further purification was necessary.

25      C<sub>20</sub>H<sub>23</sub>N<sub>3</sub>O<sub>2</sub> (MW = 337.46); mass spectroscopy found (M+H) 338.2.  
Anal. Calcd for C<sub>20</sub>H<sub>23</sub>N<sub>3</sub>O<sub>2</sub>: C, 71.19; H, 6.87; N, 12.45. Found: C, 71.32; H, 6.57; N, 12.24.

-- 201 --

**E. Cyclic Ketone Derivatives****GENERAL PROCEDURE 6-A****Jones Oxidation Procedure**

- 5           The compound to be oxidized was stirred in acetone and the Jones reagent was added in portions until the starting material was consumed. The reaction mixture was quenched with isopropanol and the mixture was filtered through Celite and concentrated under reduced pressure. The residue was partitioned between ethyl acetate and water and the organic portion was dried over sodium
- 10   sulfate and then concentrated under reduced pressure. The crude product was purified by silica gel chromatography and/or recrystallization.

**GENERAL PROCEDURE 6-B****Swern Oxidation Procedure**

- 15           To a stirred mixture of oxalyl chloride (0.15 mL, 1.2 mmol) in 10 mL of dichloromethane cooled to -78°C was added DMSO (0.106 mL, 1.5 mmol) and the mixture was stirred for 10 minutes. A solution of the alcohol (0.1828 g, 0.60 mmol) in 20 mL of chloroform was added dropwise. The reaction mixture was stirred at -78°C for 2 hours, and then 0.5 mL (3.6 mmol) of triethylamine was
- 20   added. Stirring was continued for 1 hour and then the mixture was allowed to warm to room temperature and stirring was continued at ambient temperature overnight. The mixture was then diluted with 50 mL of dichloromethane, washed with brine (3x), dried over magnesium sulfate, filtered and evaporated to dryness to give the crude product which as typically purified by column chromatography.

25

-- 202 --

**F. Dibenzazepinone Derivatives and Related Compounds****GENERAL PROCEDURE 7-A****Preparation of 5-Amino-7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Derivatives**

5

**Step A**

Following General Procedure 5-A and using 5,7-dihydro-6H-dibenz[b,d]azepin-6-one and an alkyl halide, the 7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one was prepared.

10 **Step B**

The 7-alkyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (1 eq.) was dissolved in THF and isoamyl nitrite (1.2 eq.) was added. The mixture was cooled to 0°C in an ice bath. NaHMDS (1.1 eq., 1M in THF) was added dropwise. After stirring for 1 hour or until the reaction was complete, the mixture was concentrated then

15 acidified with 1N HCl and extracted with EtOAc. The organic portion was dried and concentrated to yield a crude product which was purified by silica gel chromatography.

**Step C**

The resulting oxime was dissolved in EtOH/NH<sub>3</sub> (20:1) and hydrogenated

20 in a bomb using Raney nickel and hydrogen (500 psi) at 100°C for 10 hours. The resulting mixture was filtered and concentrated to provide an oil which was purified by silica gel chromatography to yield the title compound.

**GENERAL PROCEDURE 7-B**25 **Preparation of Fluoro-substituted 5,7-dihydro-6H-dibenz[b,d]azepin-6-one Derivatives**

A modification of the procedure of Robin D. Clark et al. *Tetrahedron*, Vol. 49, No. 7, pp. 1351-1356, 1993, incorporated herein by reference, was used. Specifically, an appropriately substituted N-t-Boc-2-amino-2'-

30 methylbiphenyl was dissolved in THF and cooled to -78°C. s-Butyl lithium



-- 203 --

(1.3M in cyclohexane, 2.2 eq.) was added slowly so that the temperature remained below -65°C. The resulting mixture was allowed to warm to -25°C and was stirred at that temperature for 1 hour. The mixture was cooled to -78°C. Dry CO<sub>2</sub> was bubbled through the mixture for 30 seconds. The mixture was allowed to warm to ambient temperature then was carefully quenched with water. The mixture was concentrated under reduced pressure then was adjusted to pH 3 with 1N HCl. The mixture was extracted with EtOAc and the organic portion was dried and concentrated to yield a crude material. The crude material was dissolved in methanol and the solution was saturated with HCl. The mixture was heated at reflux for 12 hours then was allowed to cool. The mixture was concentrated to provide crude lactam which was purified by chromatography or crystallization.

#### GENERAL PROCEDURE 7-C

##### 15      Resolution of 5-Amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

In a round bottom flask was added the racemic freebase amine (1.0 eq.) in methanol followed by di-*p*-toluoyl-D-tartaric acid monohydrate (1.0 eq.). The mixture was concentrated *in vacuo* to a residue and redissolved in a moderate volume of methanol and allowed to stir at room temperature open to the atmosphere (8-72 hours). The solid was removed by filtration. The enantiomeric excess was determined by chiral HPLC (Chiracel ODR) using 15% acetonitrile and 85% H<sub>2</sub>O with 0.1% trifluoroacetic acid and a flow rate of 1.0 mL/min at 35°C. The resolved di-*p*-toluoyl-D-tartaric salt was then dissolved in EtOAc and saturated NaHCO<sub>3</sub> until pH 9-10 was reached. The layers were separated and the organic layer was washed again with saturated NaHCO<sub>3</sub>, H<sub>2</sub>O, and brine. The organic layer was dried over MgSO<sub>4</sub> and the drying agent was removed by filtration. The filtrate was concentrated *in vacuo*. The free amine was dissolved in MeOH and HCl (12M, 1.0 eq.) was added. The salt was concentrated *in vacuo*

-- 204 --

and the resulting film was triturated with EtOAc. The HCl salt was filtered and rinsed with EtOAc. The ee was determined by chiral HPLC.

### EXAMPLE 7-A

5                    Synthesis of 5-Amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

Step A            Synthesis of 7-Methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

A round bottom flask was charged with sodium hydride (0.295 g, 7.46 mmol) in 9.0 ml of DMF and treated with 5,7-dihydro-6H-dibenz[b,d]azepin-6-one (1.3 g, 6.22 mmol) (CAS # 20011-90-9, prepared as described in Brown, et al., Tetrahedron Letters, No. 8, 667-670, (1971) and references cited therein, which are incorporated herein by reference). After stirring at 60°C for 1 h, the solution was treated with methyl iodide (1.16 ml, 18.6 mmol) and stirring continued for 17 h with the exclusion of light. After cooling, the reaction was diluted with CH<sub>2</sub>Cl<sub>2</sub>/H<sub>2</sub>O, washed with NaHSO<sub>4</sub> solution, H<sub>2</sub>O, and dried over Na<sub>2</sub>SO<sub>4</sub>. Evaporation and flash chromatography (SiO<sub>2</sub>, CHCl<sub>3</sub>) gave 0.885 g (63%) of the title compound as a colorless solid.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>): δ = 7.62 (d, 2H), 7.26-7.47 (m, 6H), 3.51 (m, 2H), 3.32 (s, 3H).

C<sub>15</sub>H<sub>13</sub>NO (MW = 223.27); mass spectroscopy (MH<sup>+</sup>) 223.

Anal. Calcd for C<sub>15</sub>H<sub>13</sub>NO; C, 80.69 H, 5.87 N, 6.27. Found: C, 80.11 H, 5.95 N, 6.23.

25                    Step B            Synthesis of 7-Methyl-5-oximo-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

The compound isolated above (0.700 g, 3.14 mmol) was dissolved in 20 ml of toluene and treated with butyl nitrite (0.733 ml, 6.28 mmol). The reaction temperature was lowered to 0°C and the solution was treated with KHMDS (9.42 ml, 0.5 M) under N<sub>2</sub> atmosphere. After stirring for 1 h the reaction was quenched with a saturated solution of NaHSO<sub>4</sub>, diluted with CH<sub>2</sub>Cl<sub>2</sub> and

-- 205 --

separated. The organic layer was dried over  $\text{Na}_2\text{SO}_4$  and the title compound purified by chromatography ( $\text{SiO}_2$ , 98:2  $\text{CHCl}_3/\text{MeOH}$ ) giving 0.59 g (80 %) as a colorless solid.

$\text{C}_{15}\text{H}_{12}\text{N}_2\text{O}_2$  (MW = 252.275); mass spectroscopy (MH+) 252.

5        Anal. Calcd for  $\text{C}_{15}\text{H}_{12}\text{N}_2\text{O}_2$ ; C, 71.42 H, 4.79 N, 11.10. Found: C, 71.24 H, 4.69 N, 10.87.

**Step C**        Synthesis of 5-Amino-7-Methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

10        The oxime isolated above (0.99 g, 3.92 mmol) was hydrogenated in a Parr apparatus at 35 psi over 10 % Pd/C (0.46 g) in 3A ethanol. After 32 h the reaction mixture was filtered through a plug of celite, the filtrate evaporated to a foam and treated with a saturated solution of HCl (g) in  $\text{Et}_2\text{O}$ . The resulting colorless solid was filtered, rinsed with cold  $\text{Et}_2\text{O}$  and vacuum dried to give 0.66 g (61 %) of the title compound.

15        NMR data was as follows:

$^1\text{H}$ -nmr (DMSO-d<sub>6</sub>):  $\delta$  = 9.11 (bs, 3H), 7.78-7.41(m, 8H), 4.83 (s, 1H), 3.25 (s, 3H).

$\text{C}_{15}\text{H}_{14}\text{N}_2\text{O} \cdot \text{HCl}$  (MW = 274.753); mass spectroscopy (MH+ free base) 238.

20        Anal. Calcd for  $\text{C}_{15}\text{H}_{14}\text{N}_2\text{O} \cdot \text{HCl}$ ; C, 65.57 H, 5.50 N, 10.19 Found: C, 65.27 H, 5.67 N, 10.13.

**EXAMPLE 7-B**

25        Synthesis of (S)- and (R)-5-(L-Alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

**Step A**        Synthesis of (S)- and (R)-5-(N-Boc-L-Alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

30        Boc-L-Alanine (0.429 g, 2.26 mmol) (Aldrich) was dissolved in THF and treated with HOBt (0.305 g, 2.26 mmol), and 5-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (0.45 g, 1.89 mmol) (Example 7-A). The temperature

-- 206 --

was lowered to 0°C and the reaction mixture treated with EDC (0.449 g, 2.26 mmol) (Alrich) and stirred 17 hours under N<sub>2</sub>. The reaction mixture was evaporated, the residue diluted with EtOAc/H<sub>2</sub>O, washed 1.0 N HCl, sat. NaHCO<sub>3</sub>, brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The diastereomers were separated on a  
5 Chiralcel OD column using 10% IPA/heptane at 1.5 ml/minute.

Isomer 1: Retention time 3.37 minutes.

NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>): δ = 7.62-7.33 (m, 9H), 5.26 (d, 1H), 5.08 (m, 1H),  
4.34 (m, 1H), 3.35 (s, 3H), 1.49 (s, 9H), 1.40 (d, 3H).

10 Optical Rotation: [α]<sub>20</sub> = - 96 @ 589 nm (c = 1, MeOH).

C<sub>23</sub>H<sub>27</sub>N<sub>3</sub>O<sub>4</sub> (MW = 409.489); mass spectroscopy (MH<sup>+</sup>) 409.

Anal. Calcd for C<sub>23</sub>H<sub>27</sub>N<sub>3</sub>O<sub>4</sub>; C, 67.46 H, 6.64 N, 10.26. Found: C, 68.42 H, 7.02 N, 9.81.

Isomer 2: Retention time 6.08 minutes.

15 NMR data was as follows:

<sup>1</sup>H-nmr (CDCl<sub>3</sub>): δ = 7.74 (bd, 1H), 7.62-7.32 (m, 8H), 5.28 (d, 1H),  
4.99 (m, 1H), 4.36 (m, 1H), 3.35 (s, 3H), 1.49 (s, 9H), 1.46 (d, 3H).

Optical Rotation: [α]<sub>20</sub> = 69 @ 589 nm (c = 1, MeOH).

C<sub>23</sub>H<sub>27</sub>N<sub>3</sub>O<sub>4</sub> (MW = 409.489); mass spectroscopy (MH<sup>+</sup>) 409.

20 Anal. Calcd for C<sub>23</sub>H<sub>27</sub>N<sub>3</sub>O<sub>4</sub>; C, 67.46 H, 6.64 N, 10.26. Found: C, 67.40 H, 6.62 N, 10.02

Step B      Synthesis of (S)- and (R)-5-(L-Alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one Hydrochloride

The compounds isolated in Step A (each isomer separately) were dissolved  
25 in dioxane and treated with excess HCl (g). After stirring for 17 hours, the title compounds were isolated as colorless solids after evaporation and vacuum drying.

Isomer 1:

C<sub>18</sub>H<sub>19</sub>N<sub>3</sub>O<sub>2</sub>·HCl (MW = 345.832); mass spectroscopy (MH<sup>+</sup> free base) 309.

Optical Rotation: [α]<sub>20</sub> = - 55 @ 589 nm (c = 1, MeOH).

-- 207 --

Isomer 2:

 $C_{18}H_{19}N_3O_2 \cdot HCl$  (MW = 345.832); mass spectroscopy (MH+ free base) 309.Optical Rotation:  $[\alpha]_{20} = 80 @ 589 \text{ nm}$  (c = 1, MeOH).

5

**GENERAL PROCEDURE 8-G**N-Alkylation of Amides or LactamsUsing Sodium Hydride or Potassium tert-Butoxide

To a slurry of sodium hydride or potassium tert-butoxide (1.1 eq) in 15 mL of dry DMF was added the appropriate amide (0.0042 moles) as a solution in 10 mL of DMF. The alkyl iodide was then added and a thick slurry resulted. The reaction became less thick as time elapsed and when complete by TLC the reaction had become homogeneous. The reaction mixture was poured over ice and extracted into ethyl acetate. The organic layer was washed with water, followed by brine. The organic layer was then dried over sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by HPLC (LC 2000), eluting with an ethyl acetate/hexane system.

15

**GENERAL PROCEDURE 8-I**N-Alkylation of Amides or Lactams Using Cesium Carbonate

To a solution of the amide or lactam in DMF was added cesium carbonate (1.05 eq) and an alkyl iodide (1.1 eq). The mixture was allowed to stir overnight at room temperature and then the reaction mixture was diluted with ethyl acetate and washed with water, followed by brine. The organic layer was dried over sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by HPLC (LC 2000), eluting with an ethyl acetate/hexane system.

20

25

**GENERAL PROCEDURE 8-J**BOC Removal Procedure

-- 208 --

To an N-Boc protected compound was added  $\text{CH}_2\text{Cl}_2/\text{TFA}$  (4:1) at room temperature. The reaction mixture was stirred at room temperature for 3 hours and then concentrated. The residue was extracted into dichloromethane and washed with water, saturated sodium bicarbonate, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to give the free amine.

### GENERAL PROCEDURE AA

#### Reductive Amination

To a solution of the arylamine in ethanol in a hydrogenation flask was added 1 equivalent of the 2-oxocarboxylic acid ester (e.g., pyruvate ester), followed by 10% palladium on carbon (25 weight percent based on the arylamine). The reaction was hydrogenated at 20 psi  $\text{H}_2$  on a Parr shaker until complete reaction was indicated by tlc (30 minutes to 16 hours). The reaction mixture was then filtered through a pad of Celite 545 (available from Aldrich Chemical Company, Inc.) and stripped free of solvent on a rotary evaporator. The crude product residue was then further purified via chromatography.

### GENERAL PROCEDURE AB

#### First Transesterification Technique

A solution of 1-5 equivalents of the desired alcohol was added to 1 equivalent of sodium hydride in toluene. After off-gassing had ceased, the compound to be transesterified, dissolved in toluene, was added. After 0.5 hours, the reaction was either heated to  $40^\circ\text{C}$  and placed under house vacuum ( $\sim 20$  mmHg), or nitrogen was bubbled through the solution while it was heated at  $90^\circ\text{C}$ . The reaction was followed by tlc, and when the reaction was complete the solution was cooled and quenched with water or 1M HCl, and in smaller scale reactions diluted with ethyl acetate. The organic phase was extracted with saturated aqueous  $\text{NaHCO}_3$ , then washed with saturated aqueous NaCl and dried over  $\text{MgSO}_4$ . The solution was stripped free of solvent on a rotary evaporator,

-- 209 --

and the crude product residue was then further purified by chromatography. Alternatively, the reaction mixture was worked-up by evaporation of the solvents and direct chromatography of the crude mixture.

This procedure is particularly useful in the case of costly and/or high  
5 boiling alcohols.

### GENERAL PROCEDURE AC

#### Second Transesterification Technique

The compound to be transesterified was placed in a large excess of the  
10 desired alcohol. A catalytic amount of dry NaH was added, and the reaction was followed by tlc until the presence of starting material was no longer detected. The reaction was quenched with a few milliliters of 1N HCl, and after a few minutes of stirring saturated aqueous NaHCO<sub>3</sub> was added. The organic phase was washed with saturated aqueous NaCl and dried over MgSO<sub>4</sub>. The solution was stripped  
15 free of solvent on a rotary evaporator, and the crude product residue was then further purified by chromatography.

### GENERAL PROCEDURE AD

#### Third Transesterification Technique

20 The compound to be transesterified was placed in a large excess of the desired alcohol. A catalytic amount of dry NaH was added, and the reaction was followed by tlc until the presence of starting material was no longer detected. The reaction was quenched with a few milliliters of 1N HCl, and after a few minutes of stirring saturated aqueous NaHCO<sub>3</sub> was added. The volume of the reaction  
25 mixture was reduced on a rotary evaporator until the excess alcohol was removed and then the remaining residue was taken up in ethyl acetate and additional water was added. The organic phase was washed with saturated aqueous NaCl and dried over MgSO<sub>4</sub>. The solution was stripped free of solvent on a rotary evaporator, and the crude product residue was then further purified by chromatography.

-- 210 --

This procedure is particularly employed in the case of low boiling, inexpensive alcohols, miscible with water.

### GENERAL PROCEDURE AE

#### 5                    O-Alkylation Technique

To a carboxylic acid compound (prepared, for example, by reductive amination via General Procedure AA to provide for the *N*-aryl amino acid ester, followed by hydrolysis via Procedure AF) in DMF was added 1.5 equivalents  $K_2CO_3$ , followed by 1 equivalent of alkylating agent (e.g., *tert*-butyl  
10 bromoacetate). The reaction was stirred at room temperature for 2 hours, then was quenched with water and extracted into ethyl acetate. The organic phase was washed with saturated aqueous  $NaHCO_3$ , water, and saturated aqueous  $NaCl$ , and was then dried over  $MgSO_4$ . The solution was stripped free of solvent on a rotary evaporator to yield the crude product.

15

### GENERAL PROCEDURE AF

#### Ester Hydrolysis to Free Acid

To a carboxylic ester compound (prepared, for example, by reductive amination via General Procedure AA to provide for the *N*-aryl amino acid ester) in  
20 a 1:1 mixture of  $CH_3OH/H_2O$  was added 2-5 equivalents of  $K_2CO_3$ . The mixture was heated to  $50^\circ C$  for 0.5 to 1.5 hours until tlc showed complete reaction. The reaction was cooled to room temperature and the methanol was removed on a rotary evaporator. The pH of the remaining aqueous solution was adjusted to  $\sim 2$ , and ethyl acetate was added to extract the product. The organic phase was then  
25 washed with saturated aqueous  $NaCl$  and dried over  $MgSO_4$ . The solution was stripped free of solvent on a rotary evaporator to yield the crude product.

### GENERAL PROCEDURE AG

#### *N*-Heteroarylation of Alanine



-- 211 --

A solution of 1.1 equivalents of L-alanine and 2 equivalents NaOH in DMSO was stirred at room temperature for 1 hour, then 1 equivalent of 2-chlorobenzothiazole was added. The mixture was heated to 100°C for 4 hours, then cooled to room temperature and poured onto ice. The pH of the resulting aqueous solution was adjusted to ~2, and the precipitated solid was removed by filtration. This solid was then dissolved in 1N NaOH and the resulting solution was filtered through a pad of Celite 545. The pH of the filtrate was adjusted to ~2, and the white precipitate was removed by filtration and washed with water to yield the crude product.

10

#### GENERAL PROCEDURE AH

##### EDC Coupling

To a 1:1 mixture of the desired acid and alcohol in  $\text{CH}_2\text{Cl}_2$  at 0°C was added 1.5 equivalents triethylamine, followed by 2.0 equivalents hydroxybenzotriazole monohydrate, then 1.25 equivalents of ethyl-3-(3-dimethylamino)-propyl carbodiimide HCl (EDC). The reaction was stirred overnight at room temperature, then transferred to a separatory funnel and washed with water, saturated aqueous  $\text{NaHCO}_3$ , 1N HCl, and saturated aqueous NaCl, and was then dried over  $\text{MgSO}_4$ . The solution was stripped free of solvent on a rotary evaporator to yield the crude product.

20

#### GENERAL PROCEDURE AI

##### Oxime or Amine Coupling Technique

The trichlorophenyl ester (1 eq) of a carboxylic acid was stirred in DMF or THF. The oxime or amine (1.2 eq) was added and the mixture was stirred at ambient temperature for 1-4 hours. In cases where the hydrochloride salt form of an amine was used, a suitable base such as *N,N*-diisopropylethylamine (1.2 eq) was also added. The resulting mixture was concentrated under reduced pressure

25

-- 212 --

to yield a crude product which was used without purification or was purified by silica gel chromatography and/or crystallization.

#### GENERAL PROCEDURE AJ

5

##### Alkylation Technique

The amine (1 eq), the  $\alpha$ -bromo ester (1.1 eq) and a suitable base (such as triethylamine) (2 eq) were stirred in chloroform. The resulting solution was heated at reflux for 4-12 hours. After cooling, the mixture was diluted with chloroform and washed with water. The organic portion was dried (sodium sulfate) and concentrated under reduced pressure. The crude product was purified by silica gel chromatography.

#### GENERAL PROCEDURE AK

##### Oxime or Alcohol Coupling Technique

15

The carboxylic acid (1 eq) was stirred in a suitable solvent (such as THF, dioxane or DMF). An alcohol or oxime (1-5 eq) was added. EDC hydrochloride (1.2 eq) and hydroxybenzotriazole hydrate (1 eq) were added. A suitable base (such as 4-methylmorpholine or triethylamine) (0-1 eq) was added. A catalytic amount (0.1 eq) of 4-dimethylaminopyridine was added. The mixture was stirred at ambient temperature and under a dry atmosphere of nitrogen. After 20 hours, the mixture was concentrated under reduced pressure. The resulting concentrate was partitioned between ethyl acetate and water. The organic portion was separated and washed with aqueous sodium bicarbonate and brine. The organic portion was dried (sodium sulfate) and concentrated under reduced pressure. The crude product was used without purification or was purified by silica gel chromatography and/or crystallization.

-- 213 --

**GENERAL PROCEDURE AL****EDC Coupling**

The carboxylic acid was dissolved in methylene chloride. The amino acid (1 eq.), N-methylmorpholine (5 eq.) and hydroxybenzotriazole monohydrate (1.2 eq.) were added in sequence. A cooling bath was applied to the round bottomed flask until the solution reached 0°C. At that time, 1.2 eq. of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) was added. The solution was allowed to stir overnight and come to room temperature under nitrogen pressure. The reaction mixture was worked up by washing the organic phase with saturated aqueous sodium carbonate, 0.1M citric acid, and brine before drying with sodium sulfate. The solvents were then removed to yield crude product. Pure products were obtained by flash chromatography in an appropriate solvent.

**GENERAL PROCEDURE AM****Triflate Displacement**

To a 0°C solution of *iso*-butyl R-(+)-lactate in CH<sub>2</sub>Cl<sub>2</sub> was added 1.1 equivalents of trifluoromethanesulfonic anhydride. After stirring at room temperature for 20 min, 1.1 equivalents of 2,6-lutidine was added and stirring was continued for 10 min. This solution was then transferred to a flask containing 1 equivalent the arylamine and 1 equivalent *N,N*-diisopropylethylamine in CH<sub>2</sub>Cl<sub>2</sub> or CH<sub>3</sub>NO<sub>2</sub> at 0°C. The reaction was held overnight at room temperature and then stripped free of solvent on a rotary evaporator. The residue was dissolved in ethyl acetate, washed with 5% citric acid, followed by saturated aqueous NaCl, dried over magnesium sulfate or sodium sulfate and then the solution was stripped free of solvent on a rotary evaporator to yield the crude product, which was then purified by chromatography.

-- 214 --

## GENERAL PROCEDURE AN

BOC Removal

The BOC-protected compound was added to a 1:1 mixture of  $\text{CH}_2\text{Cl}_2$  and trifluoroacetic acid, and was stirred until tlc indicated complete conversion, typically 2h. The solution was then stripped to dryness and the residue was taken up in ethyl acetate and extracted with dilute HCl. The acid reaction was neutralized and extracted with ethyl acetate. The organic phase was washed with saturated aqueous NaCl and dried over  $\text{MgSO}_4$ . The solution was stripped free of solvent on a rotary evaporator to yield the product.

10

## GENERAL PROCEDURE AO

Synthesis of Pyruvate Esters

To a mixture of pyruvic acid (8.8 g, 0.1 mol) (Aldrich) in 100 mL of benzene was added *iso*-butanol (14.82 g, 0.2 mol) and a catalytic amount of *p*-toluenesulfonic acid. The mixture was then refluxed using a Dean Stark apparatus. After 4 hours, the reaction appeared to be complete with the isolation of 1.8 g (0.1 mol) of water. The benzene and *iso*-butanol were removed on a rotary evaporator. The residue (14 g, 0.1 mol), which was primarily the pyruvate *iso*-butyl ester by nmr [ $^1\text{H-Nmr}$  ( $\text{CDCl}_3$ ):  $\delta = 4.0$  (d, 2H), 2.5 (s, 3H), 2.0 (m, 1H), 1.0 (d, 6H)], was used without further purification. By substituting other alcohols in place of *iso*-butanol (e.g., ethanol, isopropanol, *n*-butanol, benzyl alcohol and the like), other esters of pyruvic acid can be prepared in a similar manner.

20

25

## GENERAL PROCEDURE AP

Aromatic Nucleophilic Substitution of Fluorobenzenes

A mixture of 1.82 g (10 mmol) of D,L-alanine *iso*-butyl ester hydrochloride, the fluorobenzene (10 mmol) and 3 g of anhydrous potassium carbonate in 10 mL of DMSO was stirred at  $120^\circ\text{C}$  for 2-5 hours. The reaction

-- 215 --

mixture was then cooled to room temperature and diluted with 100 mL of ethyl acetate. The ethyl acetate extract was washed with water (3x), dried over  $\text{MgSO}_4$  and evaporated to dryness to afford the crude product, which was further purified by column chromatography.

5

#### GENERAL PROCEDURE AQ

##### Fourth Transesterification Technique

The ester to be transesterified was dissolved in a large excess of the alcohol and 0.3 equivalents of titanium(IV) isopropoxide (Aldrich) was added.

10 The reaction was followed by tlc until complete and then the volatiles were removed at reduced pressure. The resulting crude material was then chromatographed to obtain the desired product.

#### GENERAL PROCEDURE AR

15

##### Synthesis on *N*-BOC Anilines

To a solution of the aniline in THF was added dropwise 1 equivalent of di-*tert*-butyl dicarbonate (Aldrich) in THF and then 1.5 equivalents of 10N aqueous sodium hydroxide at 0°C. After stirring at room temperature for 16 hours, or heating at 80°C for 3 hours, if needed, the reaction mixture was diluted with ether  
20 and washed with  $\text{NaHCO}_3$ , brine, dried over sodium sulfate and potassium carbonate, concentrated at reduced pressure and chromatographed to afford the *N*-BOC aniline.

#### GENERAL PROCEDURE AS

25

##### Oxime Ester Formation

The trichlorophenyl ester (1 eq.) was stirred in DMF or THF. The oxime (1.2 eq.) was added and the mixture was stirred at ambient temperature for 1 to 4 hours. The resulting mixture was concentrated under reduced pressure and the residue was purified by silica gel chromatography and/or crystallization.

#### IV. Epoxide Preparation

## GENERAL PROCEDURE EA

### First Epoxide Opening Procedure

5 A solution of 5-(S)-[L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one hydrochloride (1 eq., Ex. 7B) in methanol (0.1 M) was treated with diisopropylethylamine (1.5-2 eq., Aldrich) and the appropriate epoxide (1-3 eq.). The resulting mixture was stirred at room temperature for 96 hr, then concentrated *in vacuo*. The resulting oil was dissolved in methylene chloride and washed with water. The aqueous layer was re-extracted with methylene chloride then the combined organic extracts were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The crude product was purified by flash chromatography.

## GENERAL PROCEDURE EB

### Second Epoxide Opening Procedure

A solution of 5-(S)-[L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one hydrochloride (1 eq., Ex. 7B) in methanol (0.1 M) was treated with diisopropylethylamine (1-3 eq., Aldrich) and the appropriate epoxide (1.5-2 eq.). The resulting mixture was heated to 50 °C for 48 hr, then cooled to room temperature and concentrated *in vacuo*. The resulting oil was dissolved in methylene chloride and washed with water. The aqueous layer was re-extracted with methylene chloride then the combined organic extracts were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The crude product was purified by flash chromatography.

-- 217 --

**EXAMPLE EB**Synthesis of N-(2(R/S)-hydroxy-2-(3,5-difluorophenyl)ethyl)-L-alanine

Following General Procedure EB and reacting 2-(3,5-difluorophenyl)-1,2-epoxyethane (Example EC1) and L-alanine methyl ester hydrochloride (Bachem),  
5 the title compound was isolated after flash chromatography with 3:1 to 1:1 (gradient) hexanes/ethyl acetate.

$C_{12}H_{15}F_2NO_3$  (MW = 259.25); mass spectroscopy (MH<sup>+</sup>) 260.2

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250 MHz,  $\delta$ ) 7.04-6.75 (m, 3H), 4.78-4.64 (m, 1H),  
3.71 (s, 3H), 3.49-3.29 (m, 1H), 2.84-2.60 (m, 2H), 1.31 (d, J = 5.6Hz, 3H).

10

**GENERAL PROCEDURE EC**Preparation of Epoxides

A solution of the appropriate aldehyde (1 eq.) in 1:1 methylene chloride/50% aq. sodium hydroxide (0.15M) was treated with  
15 tetrabutylammonium iodide (0.1 eq., Aldrich) and trimethylsulfonium iodide (1.2 eq., Aldrich). The resulting mixture was heated to 50 °C for 24 hours, then cooled to room temperature and washed with water. The aqueous layer was re-extracted with methylene chloride then the combined organic extracts were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The crude product was purified by flash  
20 chromatography.

**EXAMPLE EC1**Synthesis of 2-(3,5-difluorophenyl)-1,2-epoxyethane

Following General Procedure EC and using 3,5-difluorobenzaldehyde  
25 (Aldrich), the title compound was isolated after flash chromatography with 20:1 pentane/diethylether.

$C_8H_6F_2O$  (MW = 156.13); mass spectroscopy 156.0

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz,  $\delta$ ), 6.84-6.70 (m, 3H), 3.84-3.82 (m, 1H),  
3.16-3.12 (m, 1H), 2.73-2.70 (m, 1H).

-- 218 --

**EXAMPLE EC2**Synthesis of 2-(3,4,5-trifluorophenyl)-1,2-epoxyethane

Following General Procedure EC and using 3,4,5-trifluorobenzaldehyde (Strem), the title compound was isolated after flash chromatography with 10:1  
5 pentane/diethylether.

$C_8H_5F_3O$  (MW = 174.12); mass spectroscopy 174.1

$^1H$  NMR ( $CDCl_3$ , 300 MHz,  $\delta$ ) 6.94-6.89 (m, 2H), 3.81-3.79 (m, 1H),  
3.15-3.12 (m, 1H), 2.69-2.67 (m, 1H).

10

**EXAMPLE EC3**Synthesis of 2-(3,4,5-trifluorophenyl)-1,2-epoxyethane

Following General Procedure EC and using 4-(trifluoromethyl)-benzaldehyde (Aldrich), the title compound was isolated after flash  
chromatography with 10:1 pentane/diethylether.

15

$C_9H_7F_3O$  (MW = 188.15); mass spectroscopy 188.0

$^1H$  NMR ( $CDCl_3$ , 300 MHz,  $\delta$ ) 7.61 (d, J = 7.9Hz, 2H), 7.40 (d, J =  
7.9Hz, 2H), 3.93-3.90 (m, 1H), 3.20-3.17 (m, 1H), 2.78-2.75 (m, 1H).

**EXAMPLE EC4**

20

Synthesis of 2-(3,5-bis-(trifluoromethyl)phenyl)-1,2-epoxyethane

Following General Procedure EC and using 3,5-bis-(trifluoromethyl)-benzaldehyde (Lancaster), the title compound was isolated after flash  
chromatography with 10:1 pentane/diethylether.

$C_{10}H_6F_6O$  (MW = 256.15); mass spectroscopy 257

25

$^1H$  NMR ( $CDCl_3$ , 300 MHz,  $\delta$ ) 7.82 (s, 1H), 7.74 (s, 2H), 4.00-3.98 (m,  
1H), 3.25-3.22 (m, 1H), 2.80-2.77 (m, 1H).



-- 219 --

**GENERAL PROCEDURE ED****Deprotection of SEM protected alcohol**

An appropriate SEM protected alcohol was stirred in 5% HCl in methanol at room temperature for 17 hours. The reaction solution was poured into aqueous saturated NaHCO<sub>3</sub> and extracted with dichloromethane. The organic layer was concentrated *in vacuo* to afford the de-protected product.

**V. Synthesis of Intermediates****10 Intermediate A**

Synthesis of 5-(S)-(N'-(3-methyl-2-(S)-(2-trimethylsilylethoxy)-methoxybutyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

**15 Step A Preparation of Methyl-2-(S)-(2-trimethylsilylethoxy)-methoxy-3-methylbutanoate**

A solution of methyl-2-(S)-hydroxy-3-methylbutanoate (1.0 eq; J. Org. Chem. (1997), 62(7), 2292-2297, incorporated herein by reference) in dichloromethane was stirred with diisopropylethylamine (2.0 eq) under nitrogen. 2-Trimethylsilylethoxymethyl chloride (3 eq; Aldrich) was added dropwise via syringe. The solution was stirred at room temperature for 17 hours. The solution was washed with 1.0N HCl, and the water layer was back extracted with dichloromethane. The organics were combined, dried over Na<sub>2</sub>SO<sub>4</sub>, and the solvents removed *in vacuo*. The residue was purified via radial chromatography using 20% ethyl acetate in hexanes affording the title intermediate as a clear oil.

25 <sup>1</sup>H NMR data was as follows:

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 4.71 (2H s), δ 3.89 (1H d, J=5.7 Hz), δ 3.72 (3H s), δ 3.72-3.57 (2H m), δ 2.10-2.03 (1H m), δ 0.97-0.86 (8H m), δ 0.10 (9H s).

C<sub>12</sub>H<sub>26</sub>O<sub>4</sub>Si (MW = 262.42); mass spectroscopy (M+) 262.

-- 220 --

Step B      Preparation of 3-methyl-2-(S)-2-(trimethylsilyl)ethoxy-methoxybutanal

A solution of methyl-2-(S)-(2-trimethylsilylethoxy)methoxybutyl-3-methylbutanoate (1.0 eq) in toluene was cooled to -78 °C under a nitrogen atmosphere. DIBAL-H (Aldrich, 1.0M in toluene, 1.5eq) was added dropwise at -78 °C. The reaction was stirred at -78 °C for 20 minutes, and methanol was added to quench the reaction. The mixture was stirred for an additional 30 minutes at -78 °C, and a 10% aqueous solution of Rochelle's salt was added. The reaction was allowed to warm to room temperature, and stirred for 17 hours. The layers were separated, and the organic layer was concentrated *in vacuo*. The residue was purified by radial chromatography using 20% ethyl acetate in hexanes, affording the title intermediate as a clear oil.

<sup>1</sup>H NMR data was as follows:

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.61 (1H s), δ 4.79-4.72 (1H m), δ 4.69-4.63 (2H m), δ 3.71-3.56 (2H m), δ 2.16-2.01 (1H m), δ 1.23 (2H br s), δ 1.20-0.82 (6H m), δ 0.03 (9H s).

C<sub>11</sub>H<sub>24</sub>O<sub>3</sub>Si (MW = 232.40); mass spectroscopy (MH<sup>+</sup>) 233.

Step C      Preparation of N-(3-methyl-2-(S)-(2-trimethylsilylethoxy)-methoxybutyl)-L-alanine, methyl ester

3-methyl-2-(S)-(2-trimethylsilylethoxy)methoxybutanal (1.0 eq) was stirred in methanol at room temperature with L-Alanine, methyl ester (0.8 eq; Bachem). Sodium cyanoborohydride (0.5 eq) was added to the solution, and the reaction was stirred for 17 hours. The reaction was made alkaline by addition of aqueous saturated NaHCO<sub>3</sub> solution. The mixture was extracted with dichloromethane, dried over sodium sulfate, and concentrated *in vacuo*. The residue was purified via radial chromatography using 20% ethyl acetate in hexanes to afford the title intermediate as a colorless oil.

C<sub>15</sub>H<sub>33</sub>NO<sub>4</sub>Si (MW = 319.50); mass spectroscopy (MH<sup>+</sup>) 320

-- 221 --

Anal. Calcd for  $C_{15}H_{33}NO_4Si$ : C 56.39, H 10.41, N 4.38. Found:  
C56.15, H 10.14, N 4.42.

**Step D**      Preparation of 5-(S)-(N'-(3-methyl-2-(S)-(2-trimethylsilylethyloxy)-methoxybutyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

5

Following General Procedure II-A, Method B and using N-(3-methyl-2-(S)-(2-trimethylsilylethyloxy)methoxybutyl)-L-alanine, methyl ester, the acid was prepared. The residue was used, following General Procedure D, and using  
10 5-(S)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one hydrochloride (Example 7A) to prepare the title intermediate.

$C_{29}H_{43}N_3O_4Si$  (MW = 525.77), mass spectroscopy (MH<sup>+</sup>) 526.

Anal. Calcd. for  $C_{29}H_{43}N_3O_4Si$ : C 66.25, H 8.24, N 7.99. Found: C  
15 66.10, H 8.13, N 8.13.

#### Intermediate B

Synthesis of 3-(N'-(3-methyl-2-(S)-(2-trimethylsilylethyloxy)methoxybutyl)-L-Alaninyl)-amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

20 Following General Procedure II-A, Method B and using N-(3-methyl-2-(S)-(2-trimethylsilylethyloxy)methoxybutyl)-L-alanine, methyl ester (Intermediate A, Step C: 1 eq), the free acid was prepared. The residue was used, following General Procedure D, and using 3-amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine (CAS No. 131604-75-6) to prepare the title  
25 intermediate.

$C_{30}H_{44}N_4O_5Si$  (MW = 568.79), mass spectroscopy (MH<sup>+</sup>) 569.

Anal Calcd. for  $C_{30}H_{44}N_4O_5Si$ : C 63.35, H 7.80, N 9.85. Found: C  
63.24, H 7.48, N 9.89.

-- 222 --

**Intermediate C**Synthesis of N-(2-(R/S)-3,5-difluorophenyl-2-hydroxyethyl)-L-alanine

Following General Procedure II-A, Method B. modified by using 3:1 butanol/toluene for extraction of the product in place of ethyl acetate, and using  
5 N-(2-(R/S)-3,5-difluorophenyl-2-hydroxyethyl)-L-alanine (Example EB) the title intermediate was prepared.

<sup>1</sup>H NMR data was as follows:

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 7.04-7.01 (2H m), δ 6.89-6.79 (1H m), δ 4.97-4.83 (1H m), δ 4.00 (1H q, J = 7.0 Hz), δ 3.27-3.08 (2H m), δ 1.54 (3H d,  
10 J = 2.8 Hz)

C<sub>11</sub>H<sub>13</sub>F<sub>2</sub>NO<sub>3</sub> (MW = 245.23); mass spectroscopy (MH<sup>+</sup>) 246.

**EXAMPLE 6**

15 Synthesis of 5-(S)-(N'-(2-(3,5-difluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

A solution of 3,5-difluorophenylacetaldehyde (CAS 109346-94-3; Elliott, Michael et al. *Pestic. Sci.* (1987), 18(4), 239-44, incorporated herein by reference; 1.4 eq) and 5-(S)-[L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one hydrochloride (1 eq., Example 7B) with sodium  
20 cyanoborohydride (0.6 eq., Aldrich) in methanol was stirred at room temperature for 1.5 hours. The solvent was removed *in vacuo*, and the residue dissolved in a mixture of 1:1 aqueous saturated NaHCO<sub>3</sub>/diethyl ether. The layers were separated, and the aqueous layer was back-extracted with diethyl ether. The organic layers were combined, and concentrated *in vacuo*. The residue was  
25 purified via flash column chromatography using 50% ethyl acetate in hexanes to afford the title compound.

C<sub>26</sub>H<sub>25</sub>F<sub>2</sub>N<sub>3</sub>O<sub>2</sub> (MW = 449.50); mass spectroscopy (MH<sup>+</sup>) 450.

Anal. Calcd for C<sub>26</sub>H<sub>25</sub>F<sub>2</sub>N<sub>3</sub>O<sub>2</sub>; C 69.47, H 5.61, N 9.35, F 8.45.

Found: C 69.55, H 5.63, N 9.21, F 8.30.

-- 223 --

## EXAMPLES 7 and 8

Synthesis of 5(S)-(N'-(2(R)-hydroxy-2-(3,5-difluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

5

and

5(S)-(N'-(2(S)-hydroxy-2-(3,5-difluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

Following General Procedure EB and using 2-(3,5-difluorophenyl)-1,2-epoxyethane (Example EC1), the title compounds were isolated as pure diastereomers after flash chromatography with 1:1 hexanes/ethyl acetate.

10

Ex. 7 (less polar by TLC):  $C_{26}H_{25}F_2N_3O_3$  (MW = 465.50); mass spectroscopy (MH+) 466.2

Anal. Calcd for  $C_{26}H_{25}F_2N_3O_3$ : C, 67.09; H, 5.41; N, 9.03. Found: C, 67.06; H, 5.67; N, 8.75.

15

Ex. 8 (more polar by TLC):  $C_{26}H_{25}F_2N_3O_3$  (MW = 465.50); mass spectroscopy (MH+) 466.2

Anal. Calcd for  $C_{26}H_{25}F_2N_3O_3$ : C, 67.09; H, 5.41; N, 9.03. Found: C, 67.04; H, 5.46; N, 8.89.

20

## EXAMPLE 9

Synthesis of 5(S)-(N'-(2(R/S)-hydroxy-3-isopropoxyloxy-propyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

Following General Procedure EB and using isopropyl glycidyl ether (Aldrich), the title compound was isolated as a mixture of diastereomers after flash chromatography with 95:5 ethyl acetate/methanol.

25

$C_{24}H_{31}N_3O_4$  (MW = 425.53); mass spectroscopy (MH+) 426.2

Anal. Calcd for  $C_{24}H_{31}N_3O_4$ : C, 67.74; H, 7.34; N, 9.87. Found: C, 67.90; H, 7.40; N, 9.60.

30

## EXAMPLE 10

Synthesis of 5(S)-(N'-(2(R/S)-hydroxy-butyl)-L-alaninyl)-amino-7-methyl-

-- 224 --

5,7-dihydro-6H-dibenz[b,d]azepin-6-one

Following General Procedure EB and using 1,2-epoxybutane (Aldrich), the title compound was isolated as a mixture of diastereomers after flash chromatography with 9:1 ethyl acetate/methanol.

- 5             $C_{22}H_{27}N_3O_3$  (MW = 381.48); mass spectroscopy (MH+) 382.4  
            Anal. Calcd for  $C_{22}H_{27}N_3O_3$ : C, 69.27; H, 7.13; N, 11.02. Found: C, 69.51; H, 7.14; N, 10.76.

**EXAMPLE 11**

- 10        Synthesis of 5(S)-(N'-(2(R/S)-3-dihydroxy-propyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

Following General Procedure EB and using glycidol (Aldrich), the title compound was isolated as a mixture of diastereomers after flash chromatography with 7:3 ethyl acetate/methanol.

- 15             $C_{21}H_{25}N_3O_4$  (MW = 383.45); mass spectroscopy (MH+) 384.3  
            Anal. Calcd for  $C_{21}H_{25}N_3O_4$ : C, 65.78; H, 6.57; N, 10.96. Found: C, 64.13; H, 6.17; N, 9.84.

**EXAMPLE 12**

- 20        Synthesis of 5(S)-(N'-(2(R/S)-hydroxy-3-morpholinopropyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

Following General Procedure EB and using 3-morpholino-1,2-epoxypropane (Transworld), the title compound was isolated as a mixture of diastereomers after flash chromatography with 9:1 ethyl acetate/methanol.

- 25             $C_{25}H_{32}N_4O_4$  (MW = 452.56); mass spectroscopy (MH+) 453.2  
            Anal. Calcd for  $C_{25}H_{32}N_4O_4$ : C, 66.35; H, 7.13; N, 12.38. Found: C, 65.45; H, 7.21; N, 11.37.

**EXAMPLE 13**

- 30        Synthesis of 5(S)-(N'-(2(R/S)-hydroxy-tetradecyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

-- 225 --

Following General Procedure EB and using 1,2-epoxytetradecane (Aldrich), the title compound was isolated as a mixture of diastereomers after flash chromatography with 100% ethyl acetate.

$C_{32}H_{47}N_3O_3$  (MW = 521.75); mass spectroscopy (MH+) 522.4

5     Anal. Calcd for  $C_{32}H_{47}N_3O_3$ : C, 73.67; H, 9.08; N, 8.05. Found: C, 73.77; H, 8.16; N, 8.11.

#### EXAMPLE 14

10     Synthesis of 5(S)-(N'-(2(R/S)-hydroxy-octyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

Following General Procedure EB and using 1,2-epoxyoctane (Aldrich), the title compound was isolated as a mixture of diastereomers after flash chromatography with 100% ethyl acetate.

$C_{26}H_{35}N_3O_3$  (MW = 437.58); mass spectroscopy (MH+) 438.2

15     Anal. Calcd for  $C_{26}H_{35}N_3O_3$ : C, 71.37; H, 8.06; N, 9.60. Found: C, 67.61; H, 7.28; N, 8.93.

#### EXAMPLES 15 and 16

20     Synthesis of 5(S)-(N'-(2(R)-hydroxy-2-(3,4,5-trifluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one  
and

5(S)-(N'-(2(S)-hydroxy-2-(3,4,5-trifluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

25     Following General Procedure EB and using 2-(3,4,5-trifluorophenyl)-1,2-epoxyethane (Example EC2), the title compounds were isolated as pure diastereomers after flash chromatography with 100% ethyl acetate.

Ex. 15 (less polar by TLC):  $C_{26}H_{24}F_3N_3O_3$  (MW = 483.49); mass spectroscopy (MH+) 484.5

30     Anal. Calcd for  $C_{26}H_{24}F_3N_3O_3$ : C, 64.59; H, 5.00; N, 8.69. Found: C, 64.60; H, 4.99; N, 8.76.

-- 226 --

Ex. 16 (more polar by TLC):  $C_{26}H_{24}F_3N_3O_3$  (MW = 483.49); mass spectroscopy (MH+) 484.5

Anal. Calcd for  $C_{26}H_{24}F_3N_3O_3$ : C, 64.59; H, 5.00; N, 8.69. Found: C, 64.71; H, 4.93; N, 8.74.

5

**EXAMPLES 17 and 18**

Synthesis of 5(S)-(N'-(2(R)-hydroxy-2-(4-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one  
and

10 5(S)-(N'-(2(S)-hydroxy-2-(4-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

Following General Procedure EB and using 2-(4-(trifluoromethyl)phenyl)-1,2-epoxyethane (Example EC3), the title compounds were isolated as pure diastereomers after flash chromatography with 100% ethyl acetate.

15 Ex. 17 (less polar by TLC):  $C_{27}H_{26}F_3N_3O_3$  (MW = 497.52); mass spectroscopy (MH+) 498.3

Anal. Calcd for  $C_{27}H_{26}F_3N_3O_3$ : C, 65.18; H, 5.27; N, 8.45. Found: C, 64.60; H, 5.31; N, 8.32.

20 Ex. 18 (more polar by TLC):  $C_{27}H_{26}F_3N_3O_3$  (MW = 497.52); mass spectroscopy (MH+) 498.3

Anal. Calcd for  $C_{27}H_{26}F_3N_3O_3$ : C, 65.18; H, 5.27; N, 8.45. Found: C, 66.12; H, 5.46; N, 8.77.

**EXAMPLES 19 and 20**

25 Synthesis of 5(S)-(N'-(2(R)-hydroxy-2-(3,5-bis-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one  
and

30 5(S)-(N'-(2(S)-hydroxy-2-(3,5-bis-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

Following General Procedure EB and using 2-(3,5-bis-(trifluoromethyl)phenyl)-1,2-epoxyethane (Example EC4), the title compounds



were isolated as pure diastereomers after flash chromatography with 100% ethyl acetate.

5      Anal. Calcd for  $C_{28}H_{25}F_6N_3O_3$ : C, 59.47; H, 4.46; N, 7.43. Found: C, 61.24; H, 4.63; N, 7.71.

Anal. Calcd for  $C_{28}H_{25}F_6N_3O_3$ : C, 59.47; H, 4.46; N, 7.43. Found: C, 59.51; H, 4.40; N, 7.26.

Synthesis of 5(S)-(N'-(2(R)-hydroxy-2-(3,3,3-trifluoro)-propyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one  
and  
5(S)-(N'-(2(S)-hydroxy-2-(3,3,3-trifluoro)-propyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

Following General Procedure EA and using 3,3,3-trifluoro-1,2-  
20 epoxyp propane (Lancaster), the title compounds were isolated as pure  
diastereomers after flash chromatography with 100% ethyl acetate.

Ex. 21 (less polar by TLC):  $C_{21}H_{22}F_3N_3O_3$  (MW = 421.42); mass spectroscopy (MH+) 422.1

Anal. Calcd for  $C_{21}H_{22}F_3N_3O_3$ : C, 59.85; H, 5.26; N, 9.97. Found: C, 59.86; H, 5.54; N, 9.69.

Ex. 22 (more polar by TLC):  $C_{21}H_{22}F_3N_3O_3$  (MW = 421.42); mass spectroscopy (MH+) 422.1

Anal. Calcd for  $C_{21}H_{22}F_3N_3O_3$ : C, 59.85; H, 5.26; N, 9.97. Found: C, 59.42; H, 5.37; N, 9.85.

-- 228 --

## EXAMPLE 23

Synthesis of 5-(S)-(N'-(3-methyl-2-butanone)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

5           A solution of 5-(S)-[L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one hydrochloride (Example 7B, 1.0 eq) in dichloromethane was stirred at 0° C with DIEA (2 eq). Bromo-3-methyl-2-butanone (2 eq; Org. Syn., 1988, Coll. Vol6, p193, incorporated herein by reference) in dichloromethane was added dropwise via addition funnel. The reaction mixture  
10           was allowed to warm to room temperature, and stirring was continued for 17 hours. The stirring was stopped, and the solvents removed *in vacuo* to yield a yellow oil. This oil was purified by radial chromatography on silica gel using ethyl acetate as the eluent to afford title compound.

<sup>1</sup>H NMR data was as follows:

15           <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.64 (1H d, J=7.8 Hz), δ 7.59-7.22 (8H m), δ 5.30 (1H d, J=7.8 Hz), δ 3.81 (1H d, J=19.1 Hz), δ 3.39 (1H d, J=19.1 Hz), δ 3.29 (3H s), δ 3.07 (1H q, J=6.8 Hz), δ 2.63-2.54 (1H m) δ 1.22 (3H d, J=7.1 Hz), δ 1.15-1.02 (6H m).

C<sub>23</sub>H<sub>27</sub>N<sub>3</sub>O<sub>3</sub> (MW = 393.48); mass spectroscopy (MH<sup>+</sup>) 394.

20

## EXAMPLE 24

Synthesis of 5-(S)-(N'-(3-methyl-2-(S)-hydroxybutyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

25           Following General Procedure ED and using 5-(S)-(N'-(3-methyl-2-(S)-2-trimethylsilylethoxy)methoxybutyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (Intermediate A), the title compound was prepared.  
C<sub>23</sub>H<sub>29</sub>N<sub>3</sub>O<sub>3</sub> (MW = 395.51); mass spectroscopy (MH<sup>+</sup>) 396.

Anal. Calcd. for C<sub>23</sub>H<sub>29</sub>N<sub>3</sub>O<sub>3</sub>; C 69.85, H 7.39, N 10.62. Found: C 69.74, H 7.45, N 10.52.

30

-- 229 --

**EXAMPLE 25**Synthesis of 3-(N'-(3-methyl-2-(S)-2-(hydroxybutyl)-L-Alaninyl)-amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

Following General Procedure ED above, and using 3-(N'-(3-methyl-2-(S)-  
 5 2-(trimethylsilyl)ethoxymethoxybutyl)-L-Alaninyl)-amino-2,4-dioxo-1-methyl-  
 5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine (Intermediate B), the title  
 compound was prepared.

$C_{24}H_{30}N_4O_4$  (MW = 438.53); mass spectroscopy (MH+) 439.

Anal. Calcd. for  $C_{24}H_{30}N_4O_4$ ; C 65.73, H 6.90, N 12.78. Found: C  
 10 63.76, H 6.81, N 12.22.

**EXAMPLE 26**

Synthesis of 3-(N'-(2-(R/S)-3,5-difluorophenyl-2-hydroxyethyl)-L-alaninyl)-  
 amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-  
 15 1H-1,5-benzodiazepine

Following General Procedure D and using 3-amino-2,4-dioxo-1-methyl-5-  
 phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine (CAS No. 131604-75-6) and N-  
 (2-(R/S)-3,5-difluorophenyl-2-hydroxyethyl)-L-alanine (Intermediate C), the title  
 compound was prepared.

20  $C_{27}H_{26}F_2N_4O_4$  (MW = 508.52); mass spectroscopy (MH+) 509.

Anal. Calcd. for  $C_{27}H_{26}F_2N_4O_4$  C 63.77, H 5.15, N 11.02. Found: C  
 63.57, H 5.65, N 9.74.

**EXAMPLE 27**

25 Synthesis of 5-(S)-{N'-[(1RS, 2SR)-1,2,3,4-tetrahydro-1-hydroxy-2-  
 naphthyl]alaninyl}-amino-7-methyl-5,7-dihydro-  
 6H-dibenz[b,d]azepin-6-one

Step A Synthesis of 1,2,3,4-tetrahydro-naphthalene-1,2-dione-2-oxime

To the mixture of t-BuOK (1.2 eq.) in THF at 0 °C was added a solution  
 30 of isoamyl nitrite (1.2 eq.) and  $\alpha$ -tetralone (Aldrich) (1.0 eq.) in THF dropwise.  
 The reaction mixture was warmed to RT and stirred at RT for 2 h. The reaction  
 mixture was stripped to dryness, mixed with ice-cooled HCl (1M), extracted with

-- 230 --

EtOAc, washed with brine, dried over  $\text{MgSO}_4$ , filtered and concentrated. The residue was purified with flash chromatography (silica gel, 2:1 hexane/EtOAc) to give the title compound.

$\text{C}_{10}\text{H}_9\text{NO}_2$  (MW = 175.186); mass spectroscopy ( $\text{MH}^+$ ) 176.

5        Anal. Calcd for  $\text{C}_{10}\text{H}_9\text{NO}_2$ : C, 68.56 H, 5.18 N, 8.00; Found: C, 68.81 H, 5.20 N 7.98.

Step B        Synthesis of (1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl)-amine

The compound made in Step A above (1.0 eq.) was dissolved in THF, cooled to 0 °C. LAH(2.0 eq.) in THF (1M) was added slowly. The reaction  
10        mixture was allowed to warm to RT. The reaction mixture was stirred at RT overnight. The reaction mixture was cooled in an ice-bath, worked up with EtOH and saturated potassium sodium tartrate solution, extracted with  $\text{CH}_2\text{Cl}_2$ , washed with brine, and dried over  $\text{Na}_2\text{SO}_4$ . The residue was purified by acid/base extractions to give the title compound.

15         $\text{C}_{10}\text{H}_{13}\text{NO}$  (MW = 163.219); mass spectroscopy ( $\text{MH}^+$ ) 164.

Step C        Synthesis of N'-(1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl)-alanine ethyl ester

Following General Procedure AM using Ethyl-D-lactate (Fluka) and the compound made in B above, the title compound was prepared.

20         $\text{C}_{15}\text{H}_{21}\text{NO}_3$  (MW = 263.335); mass spectroscopy ( $\text{MH}^+$ ) 264.

Step D        Synthesis of N-(3a,4,5,9b-tetrahydronaphth[2,1-d]oxazol-2(3H)-one)-alanine ethyl ester

To a solution of the compound made in C above (1 eq.) in  $\text{CH}_2\text{Cl}_2$  under  $\text{N}_2$  at 0 °C,  $\text{Et}_3\text{N}$  (1 eq.) and carbonyldiimidazole (1 eq.) in  $\text{CH}_2\text{Cl}_2$  were added  
25        and the mixture was stirred for 3h. Another 1eq. of carbonyldiimidazole in  $\text{CH}_2\text{Cl}_2$  was added and stirred at 0 °C for 1h. The reaction mixture was stirred at RT for 3h. Water was added, stirred for 15 min, extracted with  $\text{CH}_2\text{Cl}_2$ , washed with water, dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. The residue was purified with flash chromatography (silica gel, 2:1 hexane/EtOAc) to give the title  
30        compound.

-- 231 --

$C_{16}H_{19}NO_4$  (MW = 289.329); mass spectroscopy ( $MH^+$ ) 290.

**Step E**      Synthesis of N-(3a,4,5,9b-tetrahydronaphth[2,1-d]oxazol-2(3H)-one)-alanine

Following General Procedure II-A, Method B using the compound made in

5    D above, the title compound was prepared.

$C_{14}H_{15}NO_4$  (MW = 261.276); mass spectroscopy ( $MH^+$ ) 262.

**Step F**      Synthesis of 5-(S)-{N'-[(3a RS, 9a SR)-3a,4,5,9b-tetrahydronaphth[2,1-d]oxazol-2(3H)-one]-alaninyl}-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

10      Following General Procedure D using the compound made in E above and 5-(S)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one hydrochloride (Example 7A), the title compound was prepared, which was purified by flash chromatography (silica gel, 5% MeOH/ $CH_2Cl_2$ ).

$C_{29}H_{27}N_3O_4$  (MW = 481.549); mass spectroscopy ( $MH^+$ ) 482.

15      Anal. Calcd for  $C_{29}H_{27}N_3O_4$ : C, 72.33 H, 5.65 N, 8.73; Found: C, 72.44 H, 5.90 N 8.55.

**Step G**      Synthesis of 5-S-{N'-[(1RS, 2SR)-1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl]alaninyl}amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

20      The compound made in F above was heated under reflux with 2 N NaOH in EtOH/ $H_2O$  (2:1) for 20 min. The reaction mixture was stripped to dryness, extracted with EtOAc, washed with water, and dried over  $Na_2SO_4$ . Evaporation and SCX column purification [5% MeOH(7N  $NH_3$ )/ $CH_2Cl_2$ ] gave the title compound as a mixture of four diastereomers.

25       $C_{28}H_{29}N_3O_3$  (MW = 455.555); mass spectroscopy ( $MH^+$ ) 456.

Anal. Calcd for  $C_{28}H_{29}N_3O_3$ : C, 73.82 H, 6.42 N, 9.22; Found: C, 73.87 H, 6.30 N 9.46.

Four isomers were isolated by HPLC (reversed phase,  $H_2O/CH_3CN$  with 0.1% TFA), SCX column purification [5% MeOH(7N  $NH_3$ )/ $CH_2Cl_2$ ] and  
30    chromatography [silica gel, 5% MeOH(7N  $NH_3$ )/ $CH_2Cl_2$ ]:

-- 232 --

Isomer 1:  $^1\text{H-nmr}(\text{CDCl}_3)$ :  $\delta = 8.20$  [d, 1H, (-CONH-)]

Isomer 2:  $^1\text{H-nmr}(\text{CDCl}_3)$ :  $\delta = 8.38$  [d, 1H, (-CONH-)]

Isomer 3:  $^1\text{H-nmr}(\text{CDCl}_3)$ :  $\delta = 8.60$  [d, 1H, (-CONH-)]

Isomer 4:  $^1\text{H-nmr}(\text{CDCl}_3)$ :  $\delta = 9.03$  [d, 1H, (-CONH-)]

5

**EXAMPLE 28**

Synthesis of 5-(S)-{N'-[(1RS, 2RS)-1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl]alaninyl}-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

10 Step A Synthesis of N-[(1RS, 2RS)-1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl]-alanine ethyl ester

Following General Procedure AM using Ethyl-D-lactate (Fluka) and N-[(1RS, 2RS)-1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl]-amine (Liebigs Ann. Chem. 1992, pp273, incorporated herein by reference), the title compound was

15 prepared.  $\text{C}_{15}\text{H}_{21}\text{NO}_3$  (MW = 263.335); mass spectroscopy ( $\text{MH}^+$ ) 264.

Step B Synthesis of N-[(1RS, 2RS)-1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl]-alanine lithium salt

Following General Procedure II-A, Method B, and using the compound made in A above except without work-up, the title compound was obtained by

20 stripping the reaction mixture to dryness.

$\text{C}_{13}\text{H}_{16}\text{NO}_3\text{Li}$  (MW = 241.241); mass spectroscopy ( $\text{MH}^+$ ) 242.

Step C Synthesis of 5-(S)-{N'-[(1RS, 2RS)-1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl]-alaninyl}-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

25 Following General Procedure D using the compound made in B above and 5-(S)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one hydrochloride (Example 7A), the title compound was prepared as a mixture of three diastereomers, which were separated with HPLC (reversed phase, 0.1% TFA in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ ) and SCX column [5% MeOH (7N  $\text{NH}_3$ )/ $\text{CH}_2\text{Cl}_2$ ] to give the

30 following isomers:

Isomer 1:  $^1\text{H-nmr}(\text{CDCl}_3)$ :  $\delta = 8.70$  [d, 1H, (-CONH-)];  $\text{C}_{28}\text{H}_{29}\text{N}_3\text{O}_3$

-- 233 --

(MW = 455.555); mass spectroscopy ( $MH^+$ ) 456.

Isomer 2:  $^1H$ -nmr( $CDCl_3$ ):  $\delta$  = 9.31 [d, 1H, (-CONH-)];  $C_{28}H_{29}N_3O_3$   
(MW = 455.555); mass spectroscopy ( $MH^+$ ) 456; Anal. Calcd for  
 $C_{28}H_{29}N_3O_3$ : C, 73.82 H, 6.42 N, 9.22; Found: C, 74.05 H, 6.51 N 8.96.

5     A mixture of isomer 2 and isomer 3:  $^1H$ -nmr( $CDCl_3$ ):  $\delta$  = 9.31  
[d, (-CONH-)], 8.95 [d, (-CONH-)];  $C_{28}H_{29}N_3O_3$  (MW = 455.555); mass  
spectroscopy ( $MH^+$ ) 456; Anal. Calcd for  $C_{28}H_{29}N_3O_3$  with 1/4  $H_2O$ : C, 73.09  
H, 6.48 N, 9.14; Found: C, 73.08 H, 6.77 N 8.79.

10

**EXAMPLE 29**

Synthesis of 5-(S)-[N'-(2- $\alpha$ -tetralone)-L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one hydrochloride

Following General Procedure 6-A using 5-(S)-{N'-[(1RS, 2RS)-1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl]-L-alaninyl}-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one and a mixture of isomer 2 and 3 from Example 29, the free base of the title compound was prepared as a mixture of two diastereomers, which were treated with HCl/dioxane respectively to give isomer 1 and isomer 2 of the hydrochlorides.

15     Isomer 1: Exact mass spectroscopy: calc. for  $C_{28}H_{28}N_3O_3$  = 454.2131.  
20     Found 454.2135; Anal. Calcd for  $C_{28}H_{27}N_3O_3$  with 1/2 dioxane: C, 67.46 H, 6.05 N, 7.87; Found: C, 67.06 H, 5.74 N 7.49.

Isomer 2: Exact mass spectroscopy: calc. for  $C_{28}H_{28}N_3O_3$  =  
454.2131, Found 454.2126; Anal. Calcd for  $C_{28}H_{29}N_3O_3$  with 1 dioxane: C,  
66.48 H, 6.29 N, 7.27; Found: C, 66.32 H, 5.97 N 7.42.

25

**EXAMPLE 30**

Synthesis of 5-(S)-[N'-(1,2,3,4-tetrahydro-2-naphthyl)-L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

A mixture of 5-(S)-(L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one (Example 7-B) (1.00 eq.),  $\beta$ -tetralone (Aldrich) (1.00 eq.)

30

-- 234 --

and titanium(IV)isopropoxide (Aldrich) (1.25 eq.) was stirred at RT for 1h and then diluted with MeOH. NaBH<sub>3</sub>CN (0.67 eq.) was added and stirred at RT overnight. Water was added, diluted with EtOAc, filtered, and concentrated to give a residue, which was purified by SCX column [3 % MeOH(7N  
5 NH<sub>3</sub>)/CH<sub>2</sub>Cl<sub>2</sub>] and flash chromatography [silica gel, 3 % MeOH(7N NH<sub>3</sub>)/CH<sub>2</sub>Cl<sub>2</sub>] to give the title compound.

C<sub>28</sub>H<sub>29</sub>N<sub>3</sub>O<sub>2</sub> (MW = 439.556); mass spectroscopy (MH<sup>+</sup>) 440.

<sup>1</sup>H-nmr(CDCl<sub>3</sub>): δ = 9.50(m, 1H), 7.60-7.11 (m, 12H), 5.28-5.20 (m, 1H), 3.39-2.86 (m, 8H), 2.15-2.10 (m, 1H), 1.56-1.21 (m, 6H).

10

#### EXAMPLE 31

##### Synthesis of 5-(S)-{N'-[(1RS, 2SR)-1-hydroxy-2-cyclohexyl]alaninyl}-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

Following the procedure in Example 28 using cis-2-aminocyclohexanol  
15 hydrochloride (ACROS), the title compound was prepared.

Exact mass spectroscopy: Calc. for C<sub>24</sub>H<sub>30</sub>N<sub>3</sub>O<sub>3</sub>: 408.2287; Found 408.2283.

Anal. Calcd for C<sub>24</sub>H<sub>29</sub>N<sub>3</sub>O<sub>3</sub>: C, 70.74 H, 7.17 N, 10.31; Found: C, 70.72 H, 7.23 N 10.41.

20

#### EXAMPLE 32

##### Synthesis of 5-(S)-[N'-(4-methylpentyl)-L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

To a solution of 5-(S)-(L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-  
25 dibenz[b,d]azepin-6-one (Example 7-B) (2.00eq.) in MeOH was added a few drops of HCl-MeOH, 4-methyl-1-pentanal (1.00eq.) (made by following the procedure described in Tetrahedron Letter, No.31, 1975, pp2647, incorporated herein by reference) and molecular sieves. NaBH<sub>3</sub>CN (0.67eq.) was added. The pH of the reaction mixture was maintained at 5-6 by adding HCl-MeOH. The reaction  
30 mixture was stirred at RT overnight. The reaction mixture was basified, extracted



-- 235 --

with EtOAc, dried over  $\text{Na}_2\text{SO}_4$ . Concentration and flash chromatography [silica gel, 5% MeOH (7 N  $\text{NH}_3$ )/ $\text{CH}_2\text{Cl}_2$ ] gave the title compound.

Exact mass spectroscopy: Calc. for  $\text{C}_{24}\text{H}_{32}\text{N}_3\text{O}_2$ : 394.2495; Found 394.2499.

Anal. Calcd for  $\text{C}_{24}\text{H}_{31}\text{N}_3\text{O}_2$  with 4/5  $\text{H}_2\text{O}$ : C, 70.66 H, 8.07 N, 10.30;

5 Found: C, 70.25 H, 7.58 N 10.64.

### EXAMPLE 33

#### Cellular Screen for the Detection of Inhibitors of $\beta$ -Amyloid Production

A compound of formula I above was assayed for its ability to inhibit  $\beta$ -  
10 amyloid production in a cell line possessing the Swedish mutation. This screening  
assay employed cells (K293 = human kidney cell line) which were stably  
transfected with the gene for amyloid precursor protein 751 (APP751) containing  
the double mutation Lys<sub>651</sub>Met<sub>652</sub> to Asn<sub>651</sub>Leu<sub>652</sub> (APP751 numbering) in the  
manner described in International Patent Application Publication No. 94/10569<sup>8</sup>  
15 and Citron et al.<sup>16</sup>. This mutation is commonly called the Swedish mutation and  
the cells, designated as "293 751 SWE", were plated in Corning 96-well plates at  
2-4 x 10<sup>4</sup> cells per well in Dulbecco's minimal essential media (Sigma, St. Louis,  
MO) plus 10% fetal bovine serum. Cell number is important in order to achieve  
 $\beta$ -amyloid ELISA results within the linear range of the assay (~0.2 to 2.5 ng per  
20 mL).

Following overnight incubation at 37°C in an incubator equilibrated with  
10% carbon dioxide, media were removed and replaced with 200  $\mu\text{L}$  of a  
compound of formula I (drug) containing media per well for a two hour  
pretreatment period and cells were incubated as above. Drug stocks were  
25 prepared in 100% dimethyl sulfoxide such that at the final drug concentration used  
in the treatment, the concentration of dimethyl sulfoxide did not exceed 0.5% and,  
in fact, usually equaled 0.1%.

At the end of the pretreatment period, the media were again removed and  
replaced with fresh drug containing media as above and cells were incubated for

-- 236 --

an additional two hours. After treatment, plates were centrifuged in a Beckman GPR at 1200 rpm for five minutes at room temperature to pellet cellular debris from the conditioned media. From each well, 100  $\mu$ L of conditioned media or appropriate dilutions thereof were transferred into an ELISA plate precoated with antibody 266 [P. Seubert, *Nature* (1992) 359:325-327]<sup>17</sup> against amino acids 13-28 of  $\beta$ -amyloid peptide as described in International Patent Application Publication No. 94/10569<sup>8</sup> and stored at 4°C overnight. An ELISA assay employing labelled antibody 3D6 [P. Seubert, *Nature* (1992) 359:325-327]<sup>17</sup> against amino acids 1-5 of  $\beta$ -amyloid peptide was run the next day to measure the amount of  $\beta$ -amyloid peptide produced.

Cytotoxic effects of the compounds were measured by a modification of the method of Hansen, et al.<sup>18</sup>. To the cells remaining in the tissue culture plate was added 25  $\mu$ L of a 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (Sigma, St. Louis, MO) stock solution (5 mg/mL) to a final concentration of 1 mg/mL. Cells were incubated at 37°C for one hour, and cellular activity was stopped by the addition of an equal volume of MTT lysis buffer (20% w/v sodium dodecylsulfate in 50% dimethylformamide, pH 4.7). Complete extraction was achieved by overnight shaking at room temperature. The difference in the OD<sub>562nm</sub> and the OD<sub>650nm</sub> was measured in a Molecular Device's UV<sub>max</sub> microplate reader as an indicator of the cellular viability.

The results of the  $\beta$ -amyloid peptide ELISA were fit to a standard curve and expressed as ng/mL  $\beta$ -amyloid peptide. In order to normalize for cytotoxicity, these results were divided by the MTT results and expressed as a percentage of the results from a drug free control. All results are the mean and standard deviation of at least six replicate assays.

The test compound was assayed for  $\beta$ -amyloid peptide production inhibition activity in cells using this assay. The results of this assay demonstrate that the compounds of formula I inhibit  $\beta$ -amyloid peptide production by at least 30% as compared to the control when employed at 10  $\mu$ g/mL.

-- 237 --

**EXAMPLE 34****In Vivo Suppression of  $\beta$ -Amyloid Release and/or Synthesis**

This example illustrates how the compounds of this invention could be tested for *in vivo* suppression of  $\beta$ -amyloid release and/or synthesis. For these experiments, 3 to 4 month old PDAPP mice are used [Games et al., (1995) *Nature* 373:523-527].<sup>19</sup> Depending upon which compound is being tested, the compound is usually formulated at between 1 and 10 mg/mL. Because of the low solubility factors of the compounds, they may be formulated with various vehicles, such as corn oil (Safeway, South San Francisco, CA); 10% ethanol in corn oil; 2-hydroxypropyl- $\beta$ -cyclodextrin (Research Biochemicals International, Natick MA); and carboxy-methyl-cellulose (Sigma Chemical Co., St. Louis MO).

The mice are dosed subcutaneously with a 26 gauge needle and 3 hours later the animals are euthanized via CO<sub>2</sub> narcosis and blood is taken by cardiac puncture using a 1 cc 25G 5/8" tuberculin syringe/needle coated with solution of 0.5 M EDTA, pH 8.0. The blood is placed in a Becton-Dickinson vacutainer tube containing EDTA and spun down for 15 minutes at 1500 xg at 5°C. The brains of the mice are then removed and the cortex and hippocampus are dissected out and placed on ice.

**1. Brain Assay**

To prepare hippocampal and cortical tissue for enzyme-linked immunosorbent assays (ELISAs) each brain region is homogenized in 10 volumes of ice cold guanidine buffer (5.0 M guanidine-HCl, 50 mM Tris-HCl, pH 8.0) using a Kontes motorized pestle (Fisher, Pittsburgh PA). The homogenates are gently rocked on a rotating platform for three to four hours at room temperature and stored at -20°C prior to quantitation of  $\beta$ -amyloid.

The brain homogenates are diluted 1:10 with ice-cold casein buffer [0.25% casein, phosphate buffered saline (PBS), 0.05% sodium azide, 20  $\mu$ g/ml aprotinin, 5 mM EDTA, pH 8.0, 10  $\mu$ g/ml leupeptin], thereby reducing the final

-- 238 --

concentration of guanidine to 0.5 M, before centrifugation at 16,000 xg for 20 minutes at 4°C. Samples are further diluted, if necessary, to achieve an optimal range for the ELISA measurements by the addition of casein buffer with 0.5 M guanidine hydrochloride added. The  $\beta$ -amyloid standards (1-40 or 1-42 amino acids) were prepared such that the final composition equaled 0.5 M guanidine in the presence of 0.1% bovine serum albumin (BSA).

The total  $\beta$ -amyloid sandwich ELISA, quantitating both  $\beta$ -amyloid (aa 1-40) and  $\beta$ -amyloid (aa 1-42) consists of two monoclonal antibodies (mAb) to  $\beta$ -amyloid. The capture antibody, 266 [P. Seubert, *Nature* (1992) 359:325-327]<sup>17</sup>, is specific to amino acids 13 - 28 of  $\beta$ -amyloid. The antibody 3D6 [Johnson-Wood et al., *PNAS USA* (1997) 94:1550-1555]<sup>20</sup>, which is specific to amino acids 1-5 of  $\beta$ -amyloid, is biotinylated and served as the reporter antibody in the assay. The 3D6 biotinylation procedure employs the manufacturer's (Pierce, Rockford IL) protocol for NHS-biotin labeling of immunoglobulins except that 100 mM sodium bicarbonate, pH 8.5 buffer is used. The 3D6 antibody does not recognize secreted amyloid precursor protein (APP) or full-length APP but detects only  $\beta$ -amyloid species with an amino terminal aspartic acid. The assay has a lower limit of sensitivity of ~50 pg/ml (11 pM) and shows no cross-reactivity to the endogenous murine  $\beta$ -amyloid peptide at concentrations up to 1 ng/ml.

The configuration of the sandwich ELISA quantitating the level of  $\beta$ -amyloid (aa 1-42) employs the mAb 21F12 [Johnson-Wood et al., *PNAS USA* (1997) 94:1550-1555]<sup>20</sup> (which recognizes amino acids 33-42 of  $\beta$ -amyloid) as the capture antibody. Biotinylated 3D6 is also the reporter antibody in this assay which has a lower limit of sensitivity of ~125 pg/ml (28 pM).

The 266 and 21F12 capture mAbs are coated at 10  $\mu$ g/ml into 96 well immunoassay plates (Costar, Cambridge MA) overnight at room temperature. The plates are then aspirated and blocked with 0.25% human serum albumin in PBS buffer for at least 1 hour at room temperature, then stored desiccated at 4°C until use. The plates are rehydrated with wash buffer (Tris-buffered saline, 0.05%

-- 239 --

Tween 20) prior to use. The samples and standards are added to the plates and incubated overnight at 4°C. The plates are washed  $\geq 3$  times with wash buffer between each step of the assay. The biotinylated 3D6, diluted to 0.5  $\mu\text{g/ml}$  in casein incubation buffer (0.25 % casein, PBS, 0.05 % Tween 20, pH 7.4) is  
5 incubated in the well for 1 hour at room temperature. Avidin-HRP (Vector, Burlingame CA) diluted 1:4000 in casein incubation buffer is added to the wells for 1 hour at room temperature. The colorimetric substrate, Slow TMB-ELISA (Pierce, Cambridge MA), is added and allowed to react for 15 minutes, after which the enzymatic reaction is stopped with addition of 2 N  $\text{H}_2\text{SO}_4$ . Reaction  
10 product is quantified using a Molecular Devices Vmax (Molecular Devices, Menlo Park CA) measuring the difference in absorbance at 450 nm and 650 nm.

## 2. Blood Assay

The EDTA plasma is diluted 1:1 in specimen diluent (0.2 gm/l sodium  
15 phosphate. $\text{H}_2\text{O}$  (monobasic), 2.16 gm/l sodium phosphate. $7\text{H}_2\text{O}$  (dibasic), 0.5gm/l thimerosal, 8.5 gm/l sodium chloride, 0.5 ml Triton X-405, 6.0 g/l globulin-free bovine serum albumin; and water). The samples and standards in specimen diluent are assayed using the total  $\beta$ -amyloid assay (266 capture/3D6 reporter) described above for the brain assay except the specimen diluent was used  
20 instead of the casein diluents described.

Formulations other than those described above can also be used for oral delivery and intravenous delivery to a mammal. For oral delivery, the compound can be mixed with either 100% corn oil or, alternatively, in a solution containing 80% corn oil, 19.5% oleic acid and 0.5% labrafil. The compound can be mixed  
25 with the above solutions in concentrations ranging from 1 mg/mL to 10 mg/mL. The compound in solution is preferably administered orally to the mammal at a dose volume of 5 mL/kg of body weight. For IV delivery, the compound is preferably mixed with a solution of 3% ethanol, 3% solutol HS-15 and 94% saline. The compound is preferably mixed with the above solution in

-- 240 --


concentrations ranging from 0.25 mg/mL to 5 mg/mL. The compound in solution is preferably administered by IV to the mammal at a dose volume of 2 mL/kg of body weight.

From the foregoing description, various modifications and changes in the composition and method will occur to those skilled in the art. All such  
5 modifications coming within the scope of the appended claims are intended to be included therein.


--241--

**WHAT IS CLAIMED IS:**


1. A method for inhibiting  $\beta$ -amyloid peptide release and/or its synthesis in a cell which method comprises administering to such a cell an amount of a compound or a mixture of compounds effective in inhibiting the cellular
- 5 release and/or synthesis of  $\beta$ -amyloid peptide wherein said compounds are represented by formula I:



wherein:


 $W$  is a cyclic group selected from the group consisting of:

--242--





--243--



wherein

ring *A*, together with the atoms to which it is attached, forms a carbocyclic or heterocyclic ring selected from the group consisting of aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and

5 heterocyclic;


ring *B*, together with the atoms to which it is attached, forms a carbocyclic or heterocyclic ring selected from the group consisting of aryl, cycloalkyl,

--244--

substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

ring C, together with the atoms to which it is attached, forms a heteroaryl or heterocyclic ring;

5 Y is represented by the formula:



R<sup>1</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkenyl, substituted alkynyl, substituted cycloalkyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic;

15 R<sup>2</sup> is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclic;

R<sup>3</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, acyl, aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

20

each R<sup>4</sup> is independently selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

25

R<sup>5</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, aryl, aryloxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted

--245--

cycloalkenyl, substituted amino, heteroaryl, heterocyclic, thioalkoxy and substituted thioalkoxy;

Q is oxygen, sulfur, -S(O)-, -S(O)<sub>2</sub>-, -C(O)- or -C(S)- ;

5 Z is represented by the formula -T-CX'X"V-, wherein T is selected from the group consisting of a bond covalently linking R<sup>1</sup> to -CX'X"-, oxygen, sulfur and -NR<sup>6</sup>, wherein R<sup>6</sup> is hydrogen, acyl, alkyl, aryl or heteroaryl,

X' is hydrogen, hydroxy or fluoro,

X" is hydrogen, hydroxy or fluoro, or X' and X" together form an oxo group, and

10 V is selected from the group consisting of alkylene or substituted alkylene, or R<sup>1</sup> and Z together form an aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic or substituted heterocyclic;

X is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, acyl, aryl, 15 cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted alkenyl, heteroaryl and heterocyclic; or X and one of R<sup>4</sup> and the atoms to which they are attached form a double bond;

t is an integer from 0 to 2;


f is an integer from 0 to 2;

20 n is an integer equal to 1 or 2;


and pharmaceutically acceptable salts thereof provided that neither of X' and X" can be hydroxy or fluoro when T is other than a covalent bond linking R<sup>1</sup> to -CX'X"-.

25 2. A method for preventing the onset of Alzheimer's disease in a human patient at risk for developing Alzheimer's disease which method comprises administering to said patient a pharmaceutical composition comprising a pharmaceutically inert carrier and an effective amount of a compound or a mixture of compounds of formula I:

--246--



wherein:


*W* is a cyclic group selected from the group consisting of:

5


10

15

--247--



--248--



wherein

ring *A*, together with the atoms to which it is attached, forms a carbocyclic or heterocyclic ring selected from the group consisting of aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and

5 heterocyclic;


ring *B*, together with the atoms to which it is attached, forms a carbocyclic or heterocyclic ring selected from the group consisting of aryl, cycloalkyl,

--249--

substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

ring C, together with the atoms to which it is attached, forms a heteroaryl or heterocyclic ring;

5 Y is represented by the formula:



R<sup>1</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkenyl, substituted alkynyl, substituted cycloalkyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic;

15 R<sup>2</sup> is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclic;

R<sup>3</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, acyl, aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

20

each R<sup>4</sup> is independently selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

25

R<sup>5</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, aryl, aryloxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted

--250--

cycloalkenyl, substituted amino, heteroaryl, heterocyclic, thioalkoxy and substituted thioalkoxy;

Q is oxygen, sulfur, -S(O)-, -S(O)<sub>2</sub>-, -C(O)- or -C(S)- ;

5 Z is represented by the formula -T-CX'X''V-, wherein T is selected from the group consisting of a bond covalently linking R<sup>1</sup> to -CX'X''-, oxygen, sulfur and -NR<sup>6</sup>, wherein R<sup>6</sup> is hydrogen, acyl, alkyl, aryl or heteroaryl,

X' is hydrogen, hydroxy or fluoro,

X'' is hydrogen, hydroxy or fluoro, or X' and X'' together form an oxo group, and

10 V is selected from the group consisting of alkylene or substituted alkylene, or R<sup>1</sup> and Z together form an aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic or substituted heterocyclic;

15 X is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, acyl, aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted alkenyl, heteroaryl and heterocyclic; or X and one of R<sup>4</sup> and the atoms to which they are attached form a double bond;

t is an integer from 0 to 2;

f is an integer from 0 to 2;


20 n is an integer equal to 1 or 2;

and pharmaceutically acceptable salts thereof provided that neither of X' and X'' can be hydroxy or fluoro when T is other than a covalent bond linking R<sup>1</sup> to -CX'X''-.


25 3. A method for treating a human patient with Alzheimer's disease in order to inhibit further deterioration in the condition of that patient which method comprises administering to said patient a pharmaceutical composition comprising a pharmaceutically inert carrier and an effective amount of a compound or a mixture of compounds of formula I:



--251--



wherein:












 $W$  is a cyclic group selected from the group consisting of:

5


10

15

--252--



--253--



wherein

ring *A*, together with the atoms to which it is attached, forms a carbocyclic or heterocyclic ring selected from the group consisting of aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and

5 heterocyclic;


ring *B*, together with the atoms to which it is attached, forms a carbocyclic or heterocyclic ring selected from the group consisting of aryl, cycloalkyl,

--254--

substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

ring C, together with the atoms to which it is attached, forms a heteroaryl or heterocyclic ring;

5 Y is represented by the formula:



10

$\text{R}^1$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkenyl, substituted alkynyl, substituted cycloalkyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic;

15  $\text{R}^2$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclic;

$\text{R}^3$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, acyl, aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

20 each  $\text{R}^4$  is independently selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

25  $\text{R}^5$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, aryl, aryloxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted

--255--

cycloalkenyl, substituted amino, heteroaryl, heterocyclic, thioalkoxy and substituted thioalkoxy;

Q is oxygen, sulfur, -S(O)-, -S(O)<sub>2</sub>-, -C(O)- or -C(S)- ;

5 Z is represented by the formula -T-CX'X"V-, wherein T is selected from the group consisting of a bond covalently linking R<sup>1</sup> to -CX'X"-, oxygen, sulfur and -NR<sup>6</sup>-, wherein R<sup>6</sup> is hydrogen, acyl, alkyl, aryl or heteroaryl,

X' is hydrogen, hydroxy or fluoro,

X" is hydrogen, hydroxy or fluoro, or X' and X" together form an oxo group, and

10 V is selected from the group consisting of alkylene or substituted alkylene, or R<sup>1</sup> and Z together form an aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic or substituted heterocyclic;

X is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, acyl, aryl, 15 cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted alkenyl, heteroaryl and heterocyclic; or X and one of R<sup>4</sup> and the atoms to which they are attached form a double bond;

t is an integer from 0 to 2;

f is an integer from 0 to 2;

20 n is an integer equal to 1 or 2;

and pharmaceutically acceptable salts thereof provided that neither of X' and X" can be hydroxy or fluoro when T is other than a covalent bond linking R<sup>1</sup> to -CX'X"-.

25 4. The method according to any of Claims 1, 2 or 3 wherein n is 1.

5. The method according to any of Claims 1, 2 or 3 wherein rings A and B are preferably independently selected from the group consisting of aryl, cycloalkyl, cycloalkenyl, heteroaryl and heterocyclic.

--256--

6. The method according to Claim 5 wherein rings *A* and *B* are independently aryl.

7. The method according to any of Claims 1, 2 or 3 wherein  $R^1$  is  
5 selected from the group consisting of phenyl, 1-naphthyl, 2-naphthyl, 2-chlorophenyl, 2-fluorophenyl, 2-bromophenyl, 2-hydroxyphenyl, 2-nitrophenyl, 2-methylphenyl, 2-methoxyphenyl, 2-phenoxyphenyl, 2-trifluoromethylphenyl, 4-fluorophenyl, 4-chlorophenyl, 4-bromophenyl, 4-nitrophenyl, 4-methylphenyl, 4-hydroxyphenyl, 4-methoxyphenyl, 4-ethoxyphenyl, 4-butoxyphenyl, 4-*iso*-  
10 propylphenyl, 4-phenoxyphenyl, 4-trifluoromethylphenyl, 4-hydroxymethylphenyl, 3-methoxyphenyl, 3-hydroxyphenyl, 3-nitrophenyl, 3-fluorophenyl, 3-chlorophenyl, 3-bromophenyl, 3-phenoxyphenyl, 3-thiomethoxyphenyl, 3-methylphenyl, 3-trifluoromethylphenyl, 2,3-dichlorophenyl, 2,3-difluorophenyl, 2,4-dichlorophenyl, 2,5-dimethoxyphenyl, 3,4-dichlorophenyl, 3,4-  
15 difluorophenyl, 3,4-methylenedioxyphenyl, 3,4-dimethoxyphenyl, 3,5-difluorophenyl, 3,5-dichlorophenyl, 3,5-di-(trifluoromethyl)phenyl, 3,5-dimethoxyphenyl, 2,4-difluorophenyl, 2,6-difluorophenyl, 3,4,5-trifluorophenyl, 3,4,5-trimethoxyphenyl, 3,4,5-tri-(trifluoromethyl)phenyl, 2,4,6-trifluorophenyl, 2,4,6-trimethylphenyl, 2,4,6-tri-(trifluoromethyl)phenyl, 2,3,5-trifluorophenyl,  
20 2,4,5-trifluorophenyl, 2,5-difluorophenyl, 2-fluoro-3-trifluoromethylphenyl, 4-fluoro-2-trifluoromethylphenyl, 2-fluoro-4-trifluoromethylphenyl, 4-benzyloxyphenyl, 2-chloro-6-fluorophenyl, 2-fluoro-6-chlorophenyl, 2,3,4,5,6-pentafluorophenyl, 2,5-dimethylphenyl, 4-phenylphenyl, adamantyl, benzyl, 2-phenylethyl, 3-phenyl-*n*-propyl, 4-phenyl-*n*-butyl, methyl, ethyl, *n*-propyl, *iso*-  
25 propyl, *iso*-butyl, *sec*-butyl, *tert*-butyl, *n*-pentyl, *iso*-valeryl, *n*-hexyl, cyclopropyl, cyclobutyl, cyclohexyl, cyclopentyl, cyclopent-1-enyl, cyclopent-2-enyl, cyclohex-1-enyl, -CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>-cyclobutyl, -CH<sub>2</sub>-cyclohexyl, -CH<sub>2</sub>-cyclopentyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclobutyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclohexyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclopentyl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, fluoropyridyls (including 5-

--257--

fluoropyridin-3-yl), chloropyridyls (including 5-chloropyridin-3-yl), thien-2-yl,  
 thien-3-yl, benzothiazol-4-yl, 2-phenylbenzoxazol-5-yl, furan-2-yl, benzofuran-2-  
 yl, thionaphthen-2-yl, thionaphthen-3-yl, thionaphthen-4-yl, 2-chlorothiophen-5-yl,  
 3-methylisoxazol-5-yl, 2-(thiophenyl)thien-5-yl, 6-methoxythionaphthen-2-yl, 3-  
 5 phenyl-1,2,4-thioxadiazol-5-yl, 2-phenyloxazol-4-yl, indol-3-yl, 1-phenyl-tetraol-  
 5-yl, allyl, 2-(cyclohexyl)ethyl,  $(\text{CH}_3)_2\text{CH}=\text{CHCH}_2\text{CH}_2\text{CH}(\text{CH}_3)-$ ,  $\phi\text{C}(\text{O})\text{CH}_2-$ ,  
 thien-2-yl-methyl, 2-(thien-2-yl)ethyl, 3-(thien-2-yl)-*n*-propyl, 2-(4-  
 nitrophenyl)ethyl, 2-(4-methoxyphenyl)ethyl, norboran-2-yl, (4-  
 methoxyphenyl)methyl, (2-methoxyphenyl)methyl, (3-methoxyphenyl)methyl, (3-  
 10 hydroxyphenyl)methyl, (4-hydroxyphenyl)methyl, (4-methylphenyl)methyl, (4-  
 fluorophenyl)methyl, (4-fluorophenoxy)methyl, (2,4-dichlorophenoxy)ethyl, (4-  
 chlorophenyl)methyl, (2-chlorophenyl)methyl, (1-phenyl)ethyl, (1-(*p*-  
 chlorophenyl)ethyl, (1-trifluoromethyl)ethyl, (4-methoxyphenyl)ethyl,  
 $\text{CH}_3\text{OC}(\text{O})\text{CH}_2-$ , benzylthiomethyl, 5-(methoxycarbonyl)-*n*-pentyl, 3-  
 15 (methoxycarbonyl)-*n*-propyl, indan-2-yl, (2-methylbenzofuran-3-yl),  
 methoxymethyl,  $\text{CH}_3\text{CH}=\text{CH}-$ ,  $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}-$ , (4-chlorophenyl) $\text{C}(\text{O})\text{CH}_2-$ ,  
 4-(fluorophenyl)- $\text{NHC}(\text{O})\text{CH}_2-$ , 1-phenyl-*n*-butyl,  $(\phi)_2\text{CHNHC}(\text{O})\text{CH}_2\text{CH}_2-$ ,  
 $(\text{CH}_3)_2\text{NC}(\text{O})\text{CH}_2-$ ,  $(\phi)_2\text{CHNHC}(\text{O})\text{CH}_2\text{CH}_2-$ , methylcarbonylmethyl, (2,4-  
 dimethylphenyl) $\text{C}(\text{O})\text{CH}_2-$ , 4-methoxyphenyl- $\text{C}(\text{O})\text{CH}_2-$ , phenyl- $\text{C}(\text{O})\text{CH}_2-$ ,  
 20  $\text{CH}_3\text{C}(\text{O})\text{N}(\phi)-$ , ethenyl, methylthiomethyl,  $(\text{CH}_3)_3\text{CNHC}(\text{O})\text{CH}_2-$ , 4-  
 fluorophenyl- $\text{C}(\text{O})\text{CH}_2-$ , diphenylmethyl, phenoxymethyl, 3,4-  
 methylenedioxyphenyl- $\text{CH}_2-$ , benzo[b]thiophen-3-yl,  $(\text{CH}_3)_3\text{COC}(\text{O})\text{NHCH}_2-$ ,  
*trans*-styryl,  $\text{H}_2\text{NC}(\text{O})\text{CH}_2\text{CH}_2-$ , 2-trifluoromethylphenyl- $\text{C}(\text{O})\text{CH}_2$ ,  
 $\phi\text{C}(\text{O})\text{NHCH}(\phi)\text{CH}_2-$ , mesityl,  $\text{CH}_3\text{CH}(\text{=NHOH})\text{CH}_2-$ , 4- $\text{CH}_3-\phi-$   
 25  $\text{NHC}(\text{O})\text{CH}_2\text{CH}_2-$ ,  $\phi\text{C}(\text{O})\text{CH}(\phi)\text{CH}_2-$ ,  $(\text{CH}_3)_2\text{CHC}(\text{O})\text{NHCH}(\phi)-$ ,  
 $\text{CH}_3\text{CH}_2\text{OCH}_2-$ ,  $\text{CH}_3\text{OC}(\text{O})\text{CH}(\text{CH}_3)(\text{CH}_2)_3-$ , 2,2,2-trifluoroethyl, 1-  
 (trifluoromethyl)ethyl, 2- $\text{CH}_3$ -benzofuran-3-yl, 2-(2,4-dichlorophenoxy)ethyl,  
 $\phi\text{SO}_2\text{CH}_2-$ , 3-cyclohexyl-*n*-propyl,  $\text{CF}_3\text{CH}_2\text{CH}_2\text{CH}_2-$  and *N*-pyrrolidinyl.

--258--

8. The method according to any of Claims 1, 2 or 3 wherein each  $R^2$  is independently selected from the group consisting of alkyl, substituted alkyl, alkenyl, cycloalkyl, aryl, heteroaryl and heterocyclic.

- 5           9. The method according to Claim 8 wherein  $R^2$  is selected from the group consisting of methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *sec*-butyl, *tert*-butyl,  $-\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)_2$ , 2-methyl-*n*-butyl, 6-fluoro-*n*-hexyl, phenyl, benzyl, cyclohexyl, cyclopentyl, cycloheptyl, allyl, *iso*-but-2-enyl, 3-methylpentyl,  $-\text{CH}_2$ -cyclopropyl,  $-\text{CH}_2$ -cyclohexyl,  $-\text{CH}_2\text{CH}_2$ -cyclopropyl, 10  $-\text{CH}_2\text{CH}_2$ -cyclohexyl,  $-\text{CH}_2$ -indol-3-yl, *p*-(phenyl)phenyl, *o*-fluorophenyl, *m*-fluorophenyl, *p*-fluorophenyl, *m*-methoxyphenyl, *p*-methoxyphenyl, phenethyl, benzyl, *m*-hydroxybenzyl, *p*-hydroxybenzyl, *p*-nitrobenzyl, *m*-trifluoromethylphenyl, *p*-( $\text{CH}_3$ )<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O-benzyl, *p*-( $\text{CH}_3$ )<sub>3</sub>COC(O)CH<sub>2</sub>O-benzyl, *p*-(HOOCCH<sub>2</sub>O)-benzyl, 2-aminopyridin-6-yl, *p*-(*N*-morpholino-CH<sub>2</sub>CH<sub>2</sub>O)-benzyl,  $-\text{CH}_2\text{CH}_2\text{C}(\text{O})\text{NH}_2$ ,  $-\text{CH}_2$ -imidazol-4-yl, 15  $-\text{CH}_2$ -(3-tetrahydrofuranyl),  $-\text{CH}_2$ -thiophen-2-yl,  $-\text{CH}_2$ -(1-methyl)cyclopropyl,  $-\text{CH}_2$ -thiophen-3-yl, thiophen-3-yl, thiophen-2-yl,  $-\text{CH}_2$ -C(O)O-*t*-butyl,  $-\text{CH}_2$ -C(CH<sub>3</sub>)<sub>3</sub>,  $-\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)_2$ , 2-methylcyclopentyl, cyclohex-2-enyl,  $-\text{CH}[\text{CH}(\text{CH}_3)_2]\text{COOCH}_3$ ,  $-\text{CH}_2\text{CH}_2\text{N}(\text{CH}_3)_2$ ,  $-\text{CH}_2\text{C}(\text{CH}_3)=\text{CH}_2$ , 20  $-\text{CH}_2\text{CH}=\text{CHCH}_3$  (cis and trans),  $-\text{CH}_2\text{OH}$ ,  $-\text{CH}(\text{OH})\text{CH}_3$ ,  $-\text{CH}(\text{O}-t\text{-butyl})\text{CH}_3$ ,  $-\text{CH}_2\text{OCH}_3$ ,  $-(\text{CH}_2)_4\text{NH-Boc}$ ,  $-(\text{CH}_2)_4\text{NH}_2$ ,  $-\text{CH}_2$ -pyridyl, pyridyl,  $-\text{CH}_2$ -naphthyl,  $-\text{CH}_2$ -(*N*-morpholino), *p*-(*N*-morpholino-CH<sub>2</sub>CH<sub>2</sub>O)-benzyl, benzo[b]thiophen-2-yl, 5-chlorobenzo[b]thiophen-2-yl, 4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl, benzo[b]thiophen-3-yl, 5-chlorobenzo[b]thiophen-3-yl, benzo[b]thiophen-5-yl, 6-methoxynaphth-2-yl, 25  $-\text{CH}_2\text{CH}_2\text{SCH}_3$ , thien-2-yl and thien-3-yl.



-259-

10. The method according to any of Claims 1, 2 or 3 wherein R<sup>3</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl and cycloalkyl.

5 11. The method according to Claim 10 wherein R<sup>3</sup> is selected from the group consisting of hydrogen, methyl, 2-methypropyl, hexyl, methoxycarbonylmethyl, 3,3-dimethyl-2-oxobutyl, 4-phenylbutyl, cyclopropylmethyl, 2,2,2-trifluoroethyl, and cyclohexyl.

10 12. The method according to any of Claims 1, 2 or 3 wherein R<sup>4</sup> is hydrogen, alkyl or substituted alkyl.

13. The method according to any of Claims 1, 2 or 3 wherein R<sup>5</sup> is selected from the group consisting of alkyl, substituted alkyl, phenyl, substituted  
15 phenyl, cycloalkyl, heteroaryl and heterocyclic.

14. The method according to any of Claims 1, 2 or 3 wherein said compound is selected from the group consisting of:

20 5-(S)-(N'-(2-(3,5-difluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

5-(S)-(N'-(2(R)-hydroxy-2-(3,5-difluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

25 5-(S)-(N'-(2(S)-hydroxy-2-(3,5-difluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

5-(S)-(N'-(2(R/S)-hydroxy-3-isopropoxypropyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

30 5-(S)-(N'-(2(R/S)-hydroxy-butyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

35 5-(S)-(N'-(2(R/S)-3-dihydroxy-propyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

--260--

- 5(S)-(N'-(2(R/S)-hydroxy-3-morpholinopropyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5 5(S)-(N'-(2(R/S)-hydroxy-tetradecyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(R/S)-hydroxy-octyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 10 5(S)-(N'-(2(R)-hydroxy-2-(3,4,5-trifluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(S)-hydroxy-2-(3,4,5-trifluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 15 5(S)-(N'-(2(R)-hydroxy-2-(4-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(S)-hydroxy-2-(4-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 20 5(S)-(N'-(2(R)-hydroxy-2-(3,5-bis-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(S)-hydroxy-2-(3,5-bis-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 25 5(S)-(N'-(2(R)-hydroxy-2-(3,3,3-trifluoro)-propyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(S)-hydroxy-2-(3,3,3-trifluoro)-propyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 30 5(S)-(N'-(3-methyl-2-butanone)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(3-methyl-2-(S)-hydroxybutyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 35 3-(N'-(3-methyl-2-(S)-2-hydroxybutyl)-L-Alaninyl)-amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine
- 40 3-(N'-(2(R/S)-3,5-difluorophenyl-2-hydroxyethyl)-L-alaninyl)-amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

--261--

5-[N'-(S)-2-(4-methylpentyl)amino-3-methylbutyryl-L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

5 5-(S)-{N'-[(1RS, 2SR)-1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl]alaninyl}-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

10 5-(S)-{N'-[(1RS, 2RS)-1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl]alaninyl}-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one


5-(S)-[N'-(2- $\alpha$ -tetralone)-L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one hydrochloride

15 5-(S)-[N'-(1,2,3,4-tetrahydro-2-naphthyl)-L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

20 5-(S)-{N'-[(1RS, 2SR)-1-hydroxy-2-cyclohexyl]alaninyl}-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one


5-(S)-[N'-(4-methylpentyl)-L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

25 15. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a pharmaceutically effective amount of a compound of formula I:




wherein:


W is a cyclic group selected from the group consisting of:



--262--



--263--



wherein

ring *A*, together with the atoms to which it is attached, forms a carbocyclic or heterocyclic ring selected from the group consisting of aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and

5 heterocyclic;


ring *B*, together with the atoms to which it is attached, forms a carbocyclic or heterocyclic ring selected from the group consisting of aryl, cycloalkyl,

--264--

substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

ring C, together with the atoms to which it is attached, forms a heteroaryl or heterocyclic ring;

5 Y is represented by the formula:



10

$\text{R}^1$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkenyl, substituted alkynyl, substituted cycloalkyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic;

15  $\text{R}^2$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclic;

$\text{R}^3$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, acyl, aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

20 each  $\text{R}^4$  is independently selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

25  $\text{R}^5$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, aryl, aryloxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted

--265--

cycloalkenyl, substituted amino, heteroaryl, heterocyclic, thioalkoxy and substituted thioalkoxy;

Q is oxygen, sulfur, -S(O)-, -S(O)<sub>2</sub>-, -C(O)- or -C(S)- ;

5 Z is represented by the formula -T-CX'X"V-, wherein T is selected from the group consisting of a bond covalently linking R<sup>1</sup> to -CX'X"-, oxygen, sulfur and -NR<sup>6</sup>-, wherein R<sup>6</sup> is hydrogen, acyl, alkyl, aryl or heteroaryl,

X' is hydrogen, hydroxy or fluoro,

X" is hydrogen, hydroxy or fluoro, or X' and X" together form an oxo group, and

10 V is selected from the group consisting of alkylene or substituted alkylene, or R<sup>1</sup> and Z together form an aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic or substituted heterocyclic;

X is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, acyl, aryl, 15 cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted alkenyl, heteroaryl and heterocyclic; or X and one of R<sup>4</sup> and the atoms to which they are attached form a double bond:

t is an integer from 0 to 2;

f is an integer from 0 to 2;

20 n is an integer equal to 1 or 2;

and pharmaceutically acceptable salts thereof provided that neither of X' and X" can be hydroxy or fluoro when T is other than a covalent bond linking R<sup>1</sup> to -CX'X"-.

25 16. The pharmaceutical composition according to Claim 15 wherein n is 1.

--266--

17. The pharmaceutical composition according to Claim 15 wherein rings *A* and *B* are preferably independently selected from the group consisting of aryl, cycloalkyl, cycloalkenyl, heteroaryl and heterocyclic.

5 18. The pharmaceutical composition according to Claim 17 wherein rings *A* and *B* are independently aryl.

19. The pharmaceutical composition according to Claim 15 wherein R<sup>1</sup> is selected from the group consisting of phenyl, 1-naphthyl, 2-naphthyl, 2-chlorophenyl, 2-fluorophenyl, 2-bromophenyl, 2-hydroxyphenyl, 2-nitrophenyl, 2-methylphenyl, 2-methoxyphenyl, 2-phenoxyphenyl, 2-trifluoromethylphenyl, 4-fluorophenyl, 4-chlorophenyl, 4-bromophenyl, 4-nitrophenyl, 4-methylphenyl, 4-hydroxyphenyl, 4-methoxyphenyl, 4-ethoxyphenyl, 4-butoxyphenyl, 4-*iso*-propylphenyl, 4-phenoxyphenyl, 4-trifluoromethylphenyl, 4-hydroxymethylphenyl, 15 3-methoxyphenyl, 3-hydroxyphenyl, 3-nitrophenyl, 3-fluorophenyl, 3-chlorophenyl, 3-bromophenyl, 3-phenoxyphenyl, 3-thiomethoxyphenyl, 3-methylphenyl, 3-trifluoromethylphenyl, 2,3-dichlorophenyl, 2,3-difluorophenyl, 2,4-dichlorophenyl, 2,5-dimethoxyphenyl, 3,4-dichlorophenyl, 3,4-difluorophenyl, 3,4-methylenedioxyphenyl, 3,4-dimethoxyphenyl, 3,5-difluorophenyl, 3,5-dichlorophenyl, 3,5-di-(trifluoromethyl)phenyl, 3,5-dimethoxyphenyl, 2,4-difluorophenyl, 2,6-difluorophenyl, 3,4,5-trifluorophenyl, 3,4,5-trimethoxyphenyl, 3,4,5-tri-(trifluoromethyl)phenyl, 2,4,6-trifluorophenyl, 2,4,6-trimethylphenyl, 2,4,6-tri-(trifluoromethyl)phenyl, 2,3,5-trifluorophenyl, 2,4,5-trifluorophenyl, 2,5-difluorophenyl, 4-fluoro-2-trifluoromethylphenyl, 2-fluoro-4-trifluoromethylphenyl, 4-benzyloxyphenyl, 2-chloro-6-fluorophenyl, 2-fluoro-6-chlorophenyl, 2,3,4,5,6-pentafluorophenyl, 2,5-dimethylphenyl, 4-phenylphenyl, 2-fluoro-3-trifluoromethylphenyl, adamantyl, benzyl, 2-phenylethyl, 3-phenyl-*n*-propyl, 4-phenyl-*n*-butyl, methyl, ethyl, *n*-propyl, *iso*-propyl, *iso*-butyl, *sec*-butyl, *tert*-butyl, *n*-pentyl, *iso*-valeryl, *n*-hexyl, cyclopropyl, 20 25



--267--

cyclobutyl, cyclohexyl, cyclopentyl, cyclopent-1-enyl, cyclopent-2-enyl, cyclohex-1-enyl, -CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>-cyclobutyl, -CH<sub>2</sub>-cyclohexyl, -CH<sub>2</sub>-cyclopentyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclobutyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclohexyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclopentyl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, fluoropyridyls (including 5-fluoropyridin-3-yl), chloropyridyls (including 5-chloropyridin-3-yl), thien-2-yl, thien-3-yl, benzothiazol-4-yl, 2-phenylbenzoxazol-5-yl, furan-2-yl, benzofuran-2-yl, thionaphthen-2-yl, thionaphthen-3-yl, thionaphthen-4-yl, 2-chlorothiophen-5-yl, 3-methylisoxazol-5-yl, 2-(thiophenyl)thien-5-yl, 6-methoxythionaphthen-2-yl, 3-phenyl-1,2,4-thioxadiazol-5-yl, 2-phenyloxazol-4-yl, indol-3-yl, 1-phenyl-tetraol-5-yl, allyl, 2-(cyclohexyl)ethyl, (CH<sub>3</sub>)<sub>2</sub>CH=CHCH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)-,  $\phi$ C(O)CH<sub>2</sub>-, thien-2-yl-methyl, 2-(thien-2-yl)ethyl, 3-(thien-2-yl)-*n*-propyl, 2-(4-nitrophenyl)ethyl, 2-(4-methoxyphenyl)ethyl, norboran-2-yl, (4-methoxyphenyl)methyl, (2-methoxyphenyl)methyl, (3-methoxyphenyl)methyl, (3-hydroxyphenyl)methyl, (4-hydroxyphenyl)methyl, (4-methylphenyl)methyl, (4-fluorophenyl)methyl, (4-fluorophenoxy)methyl, (2,4-dichlorophenoxy)ethyl, (4-chlorophenyl)methyl, (2-chlorophenyl)methyl, (1-phenyl)ethyl, (1-(*p*-chlorophenyl)ethyl, (1-trifluoromethyl)ethyl, (4-methoxyphenyl)ethyl, CH<sub>3</sub>OC(O)CH<sub>2</sub>-, benzylthiomethyl, 5-(methoxycarbonyl)-*n*-pentyl, 3-(methoxycarbonyl)-*n*-propyl, indan-2-yl, (2-methylbenzofuran-3-yl), methoxymethyl, CH<sub>3</sub>CH=CH-, CH<sub>3</sub>CH<sub>2</sub>CH=CH-, (4-chlorophenyl)C(O)CH<sub>2</sub>-, 4-(fluorophenyl)-NHC(O)CH<sub>2</sub>-, 1-phenyl-*n*-butyl, ( $\phi$ )<sub>2</sub>CHNHC(O)CH<sub>2</sub>CH<sub>2</sub>-, (CH<sub>3</sub>)<sub>2</sub>NC(O)CH<sub>2</sub>-, ( $\phi$ )<sub>2</sub>CHNHC(O)CH<sub>2</sub>CH<sub>2</sub>-, methylcarbonylmethyl, (2,4-dimethylphenyl)C(O)CH<sub>2</sub>-, 4-methoxyphenyl-C(O)CH<sub>2</sub>-, phenyl-C(O)CH<sub>2</sub>-, CH<sub>3</sub>C(O)N( $\phi$ )-, ethenyl, methylthiomethyl, (CH<sub>3</sub>)<sub>3</sub>CNHC(O)CH<sub>2</sub>-, 4-fluorophenyl-C(O)CH<sub>2</sub>-, diphenylmethyl, phenoxymethyl, 3,4-methylenedioxyphenyl-CH<sub>2</sub>-, benzo[*b*]thiophen-3-yl, (CH<sub>3</sub>)<sub>3</sub>COC(O)NHCH<sub>2</sub>-, *trans*-styryl, H<sub>2</sub>NC(O)CH<sub>2</sub>CH<sub>2</sub>-, 2-trifluoromethylphenyl-C(O)CH<sub>2</sub>-,  $\phi$ C(O)NHCH( $\phi$ )CH<sub>2</sub>-, mesityl, CH<sub>3</sub>CH(=NHOH)CH<sub>2</sub>-, 4-CH<sub>3</sub>- $\phi$ -NHC(O)CH<sub>2</sub>CH<sub>2</sub>-,  $\phi$ C(O)CH( $\phi$ )CH<sub>2</sub>-, (CH<sub>3</sub>)<sub>2</sub>CHC(O)NHCH( $\phi$ )-,

--268--

CH<sub>3</sub>CH<sub>2</sub>OCH<sub>2</sub>-, CH<sub>3</sub>OC(O)CH(CH<sub>3</sub>)(CH<sub>2</sub>)<sub>3</sub>-, 2,2,2-trifluoroethyl, 1-(trifluoromethyl)ethyl, 2-CH<sub>3</sub>-benzofuran-3-yl, 2-(2,4-dichlorophenoxy)ethyl,  $\phi$ SO<sub>2</sub>CH<sub>2</sub>-, 3-cyclohexyl-*n*-propyl, CF<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>- and N-pyrrolidinyl.

- 5            20.    The pharmaceutical composition according to Claim 15 wherein each R<sup>2</sup> is independently selected from the group consisting of alkyl, substituted alkyl, alkenyl, cycloalkyl, aryl, heteroaryl and heterocyclic.
- 10            21.    The pharmaceutical composition according to Claim 20 wherein R<sup>2</sup> is selected from the group consisting of methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *sec*-butyl, *tert*-butyl, -CH<sub>2</sub>CH(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, 2-methyl-*n*-butyl, 6-fluoro-*n*-hexyl, phenyl, benzyl, cyclohexyl, cyclopentyl, cycloheptyl, allyl, *iso*-but-2-enyl, 3-methylpentyl, -CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>-cyclohexyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclohexyl, -CH<sub>2</sub>-indol-3-yl, *p*-(phenyl)phenyl, *o*-fluorophenyl, *m*-fluorophenyl, *p*-fluorophenyl, *m*-methoxyphenyl, *p*-methoxyphenyl, phenethyl, benzyl, *m*-hydroxybenzyl, *p*-hydroxybenzyl, *p*-nitrobenzyl, *m*-trifluoromethylphenyl, *p*-(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O-benzyl, *p*-(CH<sub>3</sub>)<sub>3</sub>COC(O)CH<sub>2</sub>O-benzyl, *p*-(HOOCCH<sub>2</sub>O)-benzyl, 2-aminopyridin-6-yl, *p*-(N-morpholino-CH<sub>2</sub>CH<sub>2</sub>O)-benzyl, -CH<sub>2</sub>CH<sub>2</sub>C(O)NH<sub>2</sub>, -CH<sub>2</sub>-imidazol-4-yl, 20 -CH<sub>2</sub>-(3-tetrahydrofuran-2-yl), -CH<sub>2</sub>-thiophen-2-yl, -CH<sub>2</sub>-(1-methyl)cyclopropyl, -CH<sub>2</sub>-thiophen-3-yl, thiophen-3-yl, thiophen-2-yl, -CH<sub>2</sub>-C(O)O-*t*-butyl, -CH<sub>2</sub>-C(CH<sub>3</sub>)<sub>3</sub>, -CH<sub>2</sub>CH(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, 2-methylcyclopentyl, cyclohex-2-enyl, -CH[CH(CH<sub>3</sub>)<sub>2</sub>]COOCH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>C(CH<sub>3</sub>)=CH<sub>2</sub>, -CH<sub>2</sub>CH=CHCH<sub>3</sub> (cis and trans), -CH<sub>2</sub>OH, -CH(OH)CH<sub>3</sub>, -CH(O-*t*-butyl)CH<sub>3</sub>, 25 -CH<sub>2</sub>OCH<sub>3</sub>, -(CH<sub>2</sub>)<sub>4</sub>NH-Boc, -(CH<sub>2</sub>)<sub>4</sub>NH<sub>2</sub>, -CH<sub>2</sub>-pyridyl, pyridyl, -CH<sub>2</sub>-naphthyl, -CH<sub>2</sub>-(N-morpholino), *p*-(N-morpholino-CH<sub>2</sub>CH<sub>2</sub>O)-benzyl, benzo[b]thiophen-2-yl, 5-chlorobenzo[b]thiophen-2-yl, 4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl, benzo[b]thiophen-3-yl, 5-chlorobenzo[b]thiophen-3-yl, benzo[b]thiophen-5-yl, 6-methoxynaphth-2-yl,

--269--

-CH<sub>2</sub>CH<sub>2</sub>SCH<sub>3</sub>, thien-2-yl and thien-3-yl.

22. The pharmaceutical composition according to Claim 15 wherein R<sup>3</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl and cycloalkyl.

23. The pharmaceutical composition according to Claim 22 wherein R<sup>3</sup> is selected from the group consisting of hydrogen, methyl, 2-methypropyl, hexyl, methoxycarbonylmethyl, 3,3-dimethyl-2-oxobutyl, 4-phenylbutyl, cyclopropylmethyl, 2,2,2-trifluoroethyl, and cyclohexyl.

24. The pharmaceutical composition according to Claim 15 wherein R<sup>4</sup> is hydrogen, alkyl or substituted alkyl.

25. The pharmaceutical composition according to Claim 15 wherein R<sup>5</sup> is selected from the group consisting of alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, heteroaryl and heterocyclic.

26. The pharmaceutical composition according to Claim 15 wherein said compound is selected from the group consisting of:

5-(S)-(N'-(2-(3,5-difluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

5(S)-(N'-(2(R)-hydroxy-2-(3,5-difluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

5(S)-(N'-(2(S)-hydroxy-2-(3,5-difluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

5(S)-(N'-(2(R/S)-hydroxy-3-isopropoxy-propyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

5(S)-(N'-(2(R/S)-hydroxy-butyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

--270--

- 5(S)-(N'-(2(R/S)-3-dihydroxy-propyl)-L-alaninyl)-amino-7-methyl-  
5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5 5(S)-(N'-(2(R/S)-hydroxy-3-morpholinopropyl)-L-alaninyl)-amino-  
7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(R/S)-hydroxy-tetradecyl)-L-alaninyl)-amino-  
7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 10 5(S)-(N'-(2(R/S)-hydroxy-octyl)-L-alaninyl)-amino-7-methyl-  
5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(R)-hydroxy-2-(3,4,5-trifluorophenyl)ethyl)-L-alaninyl)-amino-  
7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 15 5(S)-(N'-(2(S)-hydroxy-2-(3,4,5-trifluorophenyl)ethyl)-L-alaninyl)-amino-  
7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(R)-hydroxy-2-(4-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-  
amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 20 5(S)-(N'-(2(S)-hydroxy-2-(4-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-  
amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 25 5(S)-(N'-(2(R)-hydroxy-2-(3,5-bis-(trifluoromethyl)phenyl)ethyl)-L-  
alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(S)-hydroxy-2-(3,5-bis-(trifluoromethyl)phenyl)ethyl)-L-  
alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 30 5(S)-(N'-(2(R)-hydroxy-2-(3,3,3-trifluoro)-propyl)-L-alaninyl)-amino-7-  
methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(S)-hydroxy-2-(3,3,3-trifluoro)-propyl)-L-alaninyl)-amino-7-  
methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 35 5(S)-(N'-(3-methyl-2-butanone)-L-alaninyl)-amino-7-methyl-  
5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 40 5(S)-(N'-(3-methyl-2-(S)-hydroxybutyl)-L-alaninyl)-amino-  
7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 3-(N'-(3-methyl-2-(S)-2-hydroxybutyl)-L-Alaninyl)-amino-2,4-dioxo-1-  
methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

--271--

3-(N'-(2-(R/S)-3,5-difluorophenyl-2-hydroxyethyl)-L-alaninyl)-amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

5 5-[N'-(S)-2-(4-methylpentyl)amino-3-methylbutyryl-L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

10 5-(S)-{N'-[(1RS, 2SR)-1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl]alaninyl}-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

15 5-(S)-{N'-[(1RS, 2RS)-1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl]alaninyl}-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one


20 5-(S)-[N'-(2- $\alpha$ -tetralone)-L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one hydrochloride

25 5-(S)-[N'-(1,2,3,4-tetrahydro-2-naphthyl)-L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

5-(S)-{N'-[(1RS, 2SR)-1-hydroxy-2-cyclohexyl]alaninyl}-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one


5-(S)-[N'-(4-methylpentyl)-L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

27. A compound of formula I:




wherein:


W is a cyclic group selected from the group consisting of:



--272--



--273--



wherein

ring **A**, together with the atoms to which it is attached, forms a carbocyclic or heterocyclic ring selected from the group consisting of aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and

5 heterocyclic;


ring **B**, together with the atoms to which it is attached, forms a carbocyclic or heterocyclic ring selected from the group consisting of aryl, cycloalkyl,

--274--

substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

ring C, together with the atoms to which it is attached, forms a heteroaryl or heterocyclic ring;

5 Y is represented by the formula:



10

$\text{R}^1$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkenyl, substituted alkynyl, substituted cycloalkyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic;

$\text{R}^2$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclic;

$\text{R}^3$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, acyl, aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

each  $\text{R}^4$  is independently selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl and heterocyclic;

$\text{R}^5$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, aryl, aryloxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, substituted amino, heteroaryl, heterocyclic, thioalkoxy and substituted thioalkoxy;



--275--

Q is oxygen, sulfur, -S(O)-, -S(O)<sub>2</sub>-, -C(O)- or -C(S)- ;

Z is represented by the formula -T-CX'X"V-, wherein T is selected from the group consisting of a bond covalently linking R<sup>1</sup> to -CX'X"-, oxygen, sulfur and -NR<sup>6</sup>-, wherein R<sup>6</sup> is hydrogen, acyl, alkyl, aryl or heteroaryl,

5 X' is hydrogen, hydroxy or fluoro,

X" is hydrogen, hydroxy or fluoro, or X' and X" together form an oxo group, and

V is selected from the group consisting of alkylene or substituted alkylene, or R<sup>1</sup> and Z together form an aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic or substituted heterocyclic;

10 X is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, acyl, aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted alkenyl, heteroaryl and heterocyclic; or X and one of R<sup>4</sup> and the atoms to which they are attached  
15 form a double bond;

t is an integer from 0 to 2;

f is an integer from 0 to 2;

n is an integer equal to 1 or 2;

and pharmaceutically acceptable salts thereof provided that neither of X' and X"  
20 can be hydroxy or fluoro when T is other than a covalent bond linking R<sup>1</sup> to -CX'X"-.

28. The compound according to Claim 27 wherein n is 1.

25 29. The compound according to Claim 27 wherein rings A and B are preferably independently selected from the group consisting of aryl, cycloalkyl, cycloalkenyl, heteroaryl and heterocyclic.

-276-

30. The compound according to Claim 29 wherein rings *A* and *B* are independently aryl.

31. The compound according to Claim 27 wherein R<sup>1</sup> is selected from

5 the group consisting of phenyl, 1-naphthyl, 2-naphthyl, 2-chlorophenyl, 2-fluorophenyl, 2-bromophenyl, 2-hydroxyphenyl, 2-nitrophenyl, 2-methylphenyl, 2-methoxyphenyl, 2-phenoxyphenyl, 2-trifluoromethylphenyl, 4-fluorophenyl, 4-chlorophenyl, 4-bromophenyl, 4-nitrophenyl, 4-methylphenyl, 4-hydroxyphenyl, 4-methoxyphenyl, 4-ethoxyphenyl, 4-butoxyphenyl, 4-*iso*-propylphenyl, 4-

10 phenoxyphenyl, 4-trifluoromethylphenyl, 4-hydroxymethylphenyl, 3-methoxyphenyl, 3-hydroxyphenyl, 3-nitrophenyl, 3-fluorophenyl, 3-chlorophenyl, 3-bromophenyl, 3-phenoxyphenyl, 3-thiomethoxyphenyl, 3-methylphenyl, 3-trifluoromethylphenyl, 2,3-dichlorophenyl, 2,3-difluorophenyl, 2,4-dichlorophenyl, 2,5-dimethoxyphenyl, 3,4-dichlorophenyl, 3,4-difluorophenyl,

15 3,4-methylenedioxyphenyl, 3,4-dimethoxyphenyl, 3,5-difluorophenyl, 3,5-dichlorophenyl, 3,5-di-(trifluoromethyl)phenyl, 3,5-dimethoxyphenyl, 2,4-difluorophenyl, 2,6-difluorophenyl, 3,4,5-trifluorophenyl, 3,4,5-trimethoxyphenyl, 3,4,5-tri-(trifluoromethyl)phenyl, 2,4,6-trifluorophenyl, 2,4,6-trimethylphenyl, 2,4,6-tri-(trifluoromethyl)phenyl, 2,3,5-trifluorophenyl, 2,4,5-

20 trifluorophenyl, 2,5-difluorophenyl, 2-fluoro-3-trifluoromethylphenyl, 4-fluoro-2-trifluoromethylphenyl, 2-fluoro-4-trifluoromethylphenyl, 4-benzyloxyphenyl, 2-chloro-6-fluorophenyl, 2-fluoro-6-chlorophenyl, 2,3,4,5,6-pentafluorophenyl, 2,5-dimethylphenyl, 4-phenylphenyl, adamantyl, benzyl, 2-phenylethyl, 3-phenyl-*n*-propyl, 4-phenyl-*n*-butyl, methyl, ethyl, *n*-propyl, *iso*-propyl, *iso*-butyl, *sec*-butyl,

25 *tert*-butyl, *n*-pentyl, *iso*-valeryl, *n*-hexyl, cyclopropyl, cyclobutyl, cyclohexyl, cyclopentyl, cyclopent-1-enyl, cyclopent-2-enyl, cyclohex-1-enyl, -CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>-cyclobutyl, -CH<sub>2</sub>-cyclohexyl, -CH<sub>2</sub>-cyclopentyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclopropyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclobutyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclohexyl, -CH<sub>2</sub>CH<sub>2</sub>-cyclopentyl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, fluoropyridyls (including 5-fluoropyridin-

--277--

- 3-yl), chloropyridyls (including 5-chloropyridin-3-yl), thien-2-yl, thien-3-yl, benzothiazol-4-yl, 2-phenylbenzoxazol-5-yl, furan-2-yl, benzofuran-2-yl, thionaphthen-2-yl, thionaphthen-3-yl, thionaphthen-4-yl, 2-chlorothiophen-5-yl, 3-methylisoxazol-5-yl, 2-(thiophenyl)thien-5-yl, 6-methoxythionaphthen-2-yl, 3-phenyl-1,2,4-thioxadiazol-5-yl, 2-phenyloxazol-4-yl, indol-3-yl, 1-phenyl-tetraol-5-yl, allyl, 2-(cyclohexyl)ethyl,  $(\text{CH}_3)_2\text{CH}=\text{CHCH}_2\text{CH}_2\text{CH}(\text{CH}_3)-$ ,  $\phi\text{C}(\text{O})\text{CH}_2-$ , thien-2-yl-methyl, 2-(thien-2-yl)ethyl, 3-(thien-2-yl)-*n*-propyl, 2-(4-nitrophenyl)ethyl, 2-(4-methoxyphenyl)ethyl, norboran-2-yl, (4-methoxyphenyl)methyl, (2-methoxyphenyl)methyl, (3-methoxyphenyl)methyl, (3-hydroxyphenyl)methyl, (4-hydroxyphenyl)methyl, (4-methylphenyl)methyl, (4-fluorophenyl)methyl, (4-fluorophenoxy)methyl, (2,4-dichlorophenoxy)ethyl, (4-chlorophenyl)methyl, (2-chlorophenyl)methyl, (1-phenyl)ethyl, (1-(*p*-chlorophenyl)ethyl, (1-trifluoromethyl)ethyl, (4-methoxyphenyl)ethyl,  $\text{CH}_3\text{OC}(\text{O})\text{CH}_2-$ , benzylthiomethyl, 5-(methoxycarbonyl)-*n*-pentyl, 3-(methoxycarbonyl)-*n*-propyl, indan-2-yl, (2-methylbenzofuran-3-yl), methoxymethyl,  $\text{CH}_3\text{CH}=\text{CH}-$ ,  $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}-$ , (4-chlorophenyl) $\text{C}(\text{O})\text{CH}_2-$ , 4-(fluorophenyl)- $\text{NHC}(\text{O})\text{CH}_2-$ , 1-phenyl-*n*-butyl,  $(\phi)_2\text{CHNHC}(\text{O})\text{CH}_2\text{CH}_2-$ ,  $(\text{CH}_3)_2\text{NC}(\text{O})\text{CH}_2-$ ,  $(\phi)_2\text{CHNHC}(\text{O})\text{CH}_2\text{CH}_2-$ , methylcarbonylmethyl, (2,4-dimethylphenyl) $\text{C}(\text{O})\text{CH}_2-$ , 4-methoxyphenyl- $\text{C}(\text{O})\text{CH}_2-$ , phenyl- $\text{C}(\text{O})\text{CH}_2-$ ,  $\text{CH}_3\text{C}(\text{O})\text{N}(\phi)-$ , ethenyl, methylthiomethyl,  $(\text{CH}_3)_3\text{CNHC}(\text{O})\text{CH}_2-$ , 4-fluorophenyl- $\text{C}(\text{O})\text{CH}_2-$ , diphenylmethyl, phenoxymethyl, 3,4-methylenedioxyphenyl- $\text{CH}_2-$ , benzo[b]thiophen-3-yl,  $(\text{CH}_3)_3\text{COC}(\text{O})\text{NHCH}_2-$ , *trans*-styryl,  $\text{H}_2\text{NC}(\text{O})\text{CH}_2\text{CH}_2-$ , 2-trifluoromethylphenyl- $\text{C}(\text{O})\text{CH}_2$ ,  $\phi\text{C}(\text{O})\text{NHCH}(\phi)\text{CH}_2-$ , mesityl,  $\text{CH}_3\text{CH}(=\text{NHOH})\text{CH}_2-$ , 4- $\text{CH}_3-\phi-\text{NHC}(\text{O})\text{CH}_2\text{CH}_2-$ ,  $\phi\text{C}(\text{O})\text{CH}(\phi)\text{CH}_2-$ ,  $(\text{CH}_3)_2\text{CHC}(\text{O})\text{NHCH}(\phi)-$ ,  $\text{CH}_3\text{CH}_2\text{OCH}_2-$ ,  $\text{CH}_3\text{OC}(\text{O})\text{CH}(\text{CH}_3)(\text{CH}_2)_3-$ , 2,2,2-trifluoroethyl, 1-(trifluoromethyl)ethyl, 2- $\text{CH}_3$ -benzofuran-3-yl, 2-(2,4-dichlorophenoxy)ethyl,  $\phi\text{SO}_2\text{CH}_2-$ , 3-cyclohexyl-*n*-propyl,  $\text{CF}_3\text{CH}_2\text{CH}_2\text{CH}_2-$  and *N*-pyrrolidinyl.

--278--

32. The compound according to Claim 27 wherein each  $R^2$  is independently selected from the group consisting of alkyl, substituted alkyl, alkenyl, cycloalkyl, aryl, heteroaryl and heterocyclic.

5           33. The compound according to Claim 32 wherein  $R^2$  is selected from the group consisting of methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *sec*-butyl, *tert*-butyl,  $-\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)_2$ , 2-methyl-*n*-butyl, 6-fluoro-*n*-hexyl, phenyl, benzyl, cyclohexyl, cyclopentyl, cycloheptyl, allyl, *iso*-but-2-enyl, 3-methylpentyl,  $-\text{CH}_2$ -cyclopropyl,  $-\text{CH}_2$ -cyclohexyl,  $-\text{CH}_2\text{CH}_2$ -cyclopropyl, -  
 10  $\text{CH}_2\text{CH}_2$ -cyclohexyl,  $-\text{CH}_2$ -indol-3-yl, *p*-(phenyl)phenyl, *o*-fluorophenyl, *m*-fluorophenyl, *p*-fluorophenyl, *m*-methoxyphenyl, *p*-methoxyphenyl, phenethyl, benzyl, *m*-hydroxybenzyl, *p*-hydroxybenzyl, *p*-nitrobenzyl, *m*-trifluoromethylphenyl, *p*-( $\text{CH}_3$ )<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O-benzyl, *p*-( $\text{CH}_3$ )<sub>3</sub>COC(O)CH<sub>2</sub>O-benzyl, *p*-(HOOCCH<sub>2</sub>O)-benzyl, 2-aminopyridin-6-yl, *p*-  
 15 (N-morpholino-CH<sub>2</sub>CH<sub>2</sub>O)-benzyl,  $-\text{CH}_2\text{CH}_2\text{C}(\text{O})\text{NH}_2$ ,  $-\text{CH}_2$ -imidazol-4-yl,  $-\text{CH}_2$ -(3-tetrahydrofuranyl),  $-\text{CH}_2$ -thiophen-2-yl,  $-\text{CH}_2$ -(1-methyl)cyclopropyl,  $-\text{CH}_2$ -thiophen-3-yl, thiophen-3-yl, thiophen-2-yl,  $-\text{CH}_2$ -C(O)O-*t*-butyl,  $-\text{CH}_2$ -C(CH<sub>3</sub>)<sub>3</sub>,  $-\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)_2$ , 2-methylcyclopentyl, cyclohex-2-enyl, -  
 20  $\text{CH}[\text{CH}(\text{CH}_3)_2]\text{COOCH}_3$ ,  $-\text{CH}_2\text{CH}_2\text{N}(\text{CH}_3)_2$ ,  $-\text{CH}_2\text{C}(\text{CH}_3)=\text{CH}_2$ , -  
 $\text{CH}_2\text{CH}=\text{CHCH}_3$  (cis and trans),  $-\text{CH}_2\text{OH}$ ,  $-\text{CH}(\text{OH})\text{CH}_3$ ,  $-\text{CH}(\text{O}-t\text{-butyl})\text{CH}_3$ , -  
 $\text{CH}_2\text{OCH}_3$ ,  $-(\text{CH}_2)_4\text{NH-Boc}$ ,  $-(\text{CH}_2)_4\text{NH}_2$ ,  $-\text{CH}_2$ -pyridyl, pyridyl,  $-\text{CH}_2$ -naphthyl,  $-\text{CH}_2$ -(N-morpholino), *p*-(N-morpholino-CH<sub>2</sub>CH<sub>2</sub>O)-benzyl, benzo[b]thiophen-2-yl, 5-chlorobenzo[b]thiophen-2-yl, 4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl, benzo[b]thiophen-3-yl, 5-  
 25 chlorobenzo[b]thiophen-3-yl, benzo[b]thiophen-5-yl, 6-methoxynaphth-2-yl,  $-\text{CH}_2\text{CH}_2\text{SCH}_3$ , thien-2-yl and thien-3-yl.

34. The compound according to Claim 27 wherein  $R^3$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl and cycloalkyl.

--279--

35. The compound according to Claim 34 wherein R<sup>3</sup> is selected from the group consisting of hydrogen, methyl, 2-methypropyl, hexyl, methoxycarbonylmethyl, 3,3-dimethyl-2-oxobutyl, 4-phenylbutyl,  
5 cyclopropylmethyl, 2,2,2-trifluoroethyl, and cyclohexyl.

36. The compound according to Claim 27 wherein R<sup>4</sup> is hydrogen, alkyl or substituted alkyl.

10 37. The compound according to Claim 27 wherein R<sup>5</sup> is selected from the group consisting of alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, heteroaryl and heterocyclic.

38. The compound according to Claim 27 wherein said compound is  
15 selected from the group consisting of:

5-(S)-(N'-(2-(3,5-difluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-  
5,7-dihydro-6H-dibenz[b,d]azepin-6-one

20 5(S)-(N'-(2(R)-hydroxy-2-(3,5-difluorophenyl)ethyl)-L-alaninyl)-amino-  
7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

5(S)-(N'-(2(S)-hydroxy-2-(3,5-difluorophenyl)ethyl)-L-alaninyl)-amino-  
7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

25 5(S)-(N'-(2(R/S)-hydroxy-3-isopropoxy-propyl)-L-alaninyl)-amino-  
7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

30 5(S)-(N'-(2(R/S)-hydroxy-butyl)-L-alaninyl)-amino-7-methyl-  
5,7-dihydro-6H-dibenz[b,d]azepin-6-one

5(S)-(N'-(2(R/S)-3-dihydroxy-propyl)-L-alaninyl)-amino-7-methyl-  
5,7-dihydro-6H-dibenz[b,d]azepin-6-one

35 5(S)-(N'-(2(R/S)-hydroxy-3-morpholinopropyl)-L-alaninyl)-amino-  
7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

5(S)-(N'-(2(R/S)-hydroxy-tetradecyl)-L-alaninyl)-amino-

--280--

- 7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5 5(S)-(N'-(2(R/S)-hydroxy-octyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(R)-hydroxy-2-(3,4,5-trifluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 10 5(S)-(N'-(2(S)-hydroxy-2-(3,4,5-trifluorophenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(R)-hydroxy-2-(4-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 15 5(S)-(N'-(2(S)-hydroxy-2-(4-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(R)-hydroxy-2-(3,5-bis-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 20 5(S)-(N'-(2(S)-hydroxy-2-(3,5-bis-(trifluoromethyl)phenyl)ethyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5(S)-(N'-(2(R)-hydroxy-2-(3,3,3-trifluoro)-propyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 25 5(S)-(N'-(2(S)-hydroxy-2-(3,3,3-trifluoro)-propyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 30 5-(S)-(N'-(3-methyl-2-butanone)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5-(S)-(N'-(3-methyl-2-(S)-hydroxybutyl)-L-alaninyl)-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 35 3-(N'-(3-methyl-2-(S)-2-hydroxybutyl)-L-Alaninyl)-amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine
- 40 3-(N'-(2(R/S)-3,5-difluorophenyl-2-hydroxyethyl)-L-alaninyl)-amino-2,4-dioxo-1-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine
- 5-[N'-(S)-2-(4-methylpentyl)amino-3-methylbutyryl-L-alaninyl]-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one

--281--

- 5-(S)-{N'-[(1RS, 2SR)-1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl]  
alaninyl}-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5 5-(S)-{N'-[(1RS, 2RS)-1,2,3,4-tetrahydro-1-hydroxy-2-naphthyl]  
alaninyl}-amino-7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5-(S)-[N'-(2- $\alpha$ -tetralone)-L-alaninyl]-amino-7-methyl-  
5,7-dihydro-6H-dibenz[b,d]azepin-6-one hydrochloride
- 10 5-(S)-[N'-(1,2,3,4-tetrahydro-2-naphthyl)-L-alaninyl]-amino-7-methyl-  
5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 5-(S)-{N'-[(1RS, 2SR)-1-hydroxy-2-cyclohexyl]alaninyl}-amino-  
7-methyl-5,7-dihydro-6H-dibenz[b,d]azepin-6-one
- 15 5-(S)-[N'-(4-methylpentyl)-L-alaninyl]-amino-7-methyl-  
5,7-dihydro-6H-dibenz[b,d]azepin-6-one.

# INTERNATIONAL SEARCH REPORT

Int lional Application No

PCT/US 99/14096

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D223/18 A61K31/55 C07D471/06 C07D243/14 C07D409/12  
C07D243/12 C07D223/16 C07D267/14 C07D281/10 C07D471/04  
C07D471/08 C07D401/04 C07D417/04 C07D409/04 C07D243/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 16410 A (MERCK & CO., INC.) 9 May 1997 (1997-05-09) example 5 ---	1
X	WO 92 16524 A (MERCK & CO., INC.) 1 October 1992 (1992-10-01) example 11 ---	1
A	WO 98 22430 A (ATHENA NEUROSCIENCES, INC.) 28 May 1998 (1998-05-28) claims 1,49 ---	1,15
A	WO 98 22494 A (ATHENA NEUROSCIENCES, INC.) 28 May 1998 (1998-05-28) claims 1,61 ---	1,15
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*A\* document member of the same patent family

Date of the actual completion of the international search

1 October 1999

Date of mailing of the international search report

20/10/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H



# INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/US 99/14096

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D487/04 //(C07D471/06,223:00,221:00),(C07D471/04,223:00,221:00),(C07D471/08,223:00,221:00),(C07D487/04,249:00,243:00)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 98 38177 A (ATHENA NEUROSCIENCES, INC.) 3 September 1998 (1998-09-03) * page 286-287: example 8E *	1



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

1 October 1999

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/ 14096

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 2-14  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 2-14  
are directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/14096

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9716410 A	09-05-1997	AU 7525796 A US 5817757 A	22-05-1997 06-10-1998
WO 9216524 A	01-10-1992	US 5206235 A AT 142206 T AU 653992 B AU 1301292 A BG 61448 B BG 98112 A CA 2063185 A CN 1066070 A,B DE 69213318 D DE 69213318 T EP 0513974 A FI 921183 A HU 66796 A IE 76459 B IL 101206 A JP 2103149 C JP 6172316 A JP 8000814 B MX 9201226 A NZ 241958 A US 5310737 A	27-04-1993 15-09-1996 20-10-1994 24-09-1992 29-08-1997 27-05-1994 21-09-1992 11-11-1992 10-10-1996 10-04-1997 19-11-1992 21-09-1992 28-12-1994 22-10-1997 18-02-1997 22-10-1996 21-06-1994 10-01-1996 01-10-1992 27-04-1995 10-05-1993
WO 9822430 A	28-05-1998	AU 5249898 A AU 5267798 A NO 992381 A NO 992463 A WO 9822433 A	10-06-1998 10-06-1998 21-07-1999 19-07-1999 28-05-1998
WO 9822494 A	28-05-1998	AU 5356198 A EP 0942924 A HR 970627 A NO 992368 A	10-06-1998 22-09-1999 31-08-1998 21-06-1999
WO 9838177 A	03-09-1998	AU 6662298 A	18-09-1998